

Formula Name KETOPROFEN ANHYDROUS GEL UNGUATOR 10% GEL

Initials AP **Quantity** 60 **MLS** **Schedule** 1 **Days to Exp** 180 **Date of Formula** 7/10/2025
Cost \$10.19 **Cost Per Unit** \$0.17 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
KETOPROFEN USP	PWD	6.0000	GMS	☐	\$4.03		SOLUBILITY: WATER(I), ALCOHOL(FS)
PROPYLENE GLYCOL USP	LIQU	18.0000	MLS	☐	\$0.50		SOLUBILITY:N/A
VERSAPRO (ANHYDROUS GEL BASE)	GEL	30.7500	GMS	☐	\$5.66		NONE

Therapy**Description**

Off-white/opaque colored cream, uniform consistency, no odor

Classification(s) Nonaqueous

Disease State(s)

Instructions**A. Calculations/Special Considerations:**

Specific weight of final preparation averages 0.91g/mL

Use of the 500mL jar requires the stir rod with an 80mm diameter blade and a 210mm length rod

Use of the 300mL jar requires the stir rod with an 80mm diameter blade and a 165mm length rod

Use of the 200mL jar requires the stir rod with a 57mm diameter blade and a 200mm length rod

Use of the 100mL jar requires the stir rod with a 57mm diameter blade and a 153.5mm length rod

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula, 60mL syringe, amber bags, unguator/mixer, unguator jar, unguator stir rod

C. Method of Preparation:

1. Calculate required amounts and fill out compounding and control usage logs

2. Measure out ingredients, prepare large unguator jars by pulling back the spindle attachment. Remove spindle by turning counter-clockwise

3. Triturate Ketoprofen in mortar.

4. Wet step 3 with propylene glycol

5. Add approximately 50% of base to appropriate sized mixing jar, then mixture from step 4 to mixing jar, then add remaining base to mixing jar

6. Attach appropriate sized stir bar to lid.

7. Attach lid to mixing jar and expel additional air from jar and place lid on jar.

8. Connect rod to unguator with a clockwise quarter turn, then attach jar to oscillator arm with counter-clockwise turns until securely attached.

9. Begin mixing process, settings at 3:00 minutes, 7 speed.

10. If storing in 60ml syringe, remove syringe plunger, add stopper to syringe, and using the spindle attachment push gel into each syringe.

11. Package and label with barcode label and affix "beyond use date" and "external use only" labels

12. Tape labels

13. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxiliary sticker

Beyond use date sticker

Add prepack label if compound lot made to stock for multiple Rx

Add Rx auxiliary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Off-white/opaque colored cream, uniform consistency, no odor

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag or Samix mixing jar

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist