

Formula Name KETOPROFEN GEL INACTIVE 10% GEL

Initials AP **Quantity** 60 **MLS** **Schedule** 0 **Days to Exp** 14 **Date of Formula** 2/2/2021
Cost \$5.38 **Cost Per Unit** \$0.09 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
KETOPROFEN USP	PWD	6.0000	GMS	☐	\$2.22		SOLUBILITY: WATER(I), ALCOHOL(FS)
ETHYL ALCOHOL 95% USP	LIQU	6.0000	MLS	☐	\$0.41		MISCIBLE WITH WATER 1. THIS ALCOHOL IS NOT DENATURED. 2.***DO NOT USE DENATURED ALCOHOL IN ANY COMPOUNDED PRODUCT*** SOLUBILITY:N/A
LIPMAX LECITHIN & ISOPROPYL PALMITATE	SOLN	13.2000	MLS	☐	\$0.49		
POLOX 20% GEL	GEL	60.0000	GMS	☒	\$2.27		NONE

Therapy**Description**

Solid light yellow, oily, highly viscous organogel

Classification(s) Non-Preserved Aqueous

Disease State(s)

Instructions**A. Calculations/Special Considerations:**

None

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding
2. Measure out ingredients
3. Heat Ethyl Alcohol to 40°C in an appropriate size beaker and spin with stir bar
4. Triturate Ketoprofen in mortar.
5. Add Ketoprofen powder to heated Ethyl Alcohol and wait until dissolved
6. Add Lipmax and mix until evenly dispersed
7. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe
8. In the same syringe QS with polox, add plunger loosely, invert and tap mixture to the bottom, once tip is clear, fully insert plunger and expel extra air
9. Attach a new syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (typically 20 transfers)
10. Package and label with barcode label and affix "external use only" label
11. Tape labels
12. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:

Solid light yellow, oily, highly viscous organogel

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

PCCA 9142

Peacock, G. G., Sauvageot, J., & Addo, R. T. (2013) Evaluation of the stability of ketoprofen in pluronic lecithin organogel and the determination of an appropriate beyond-use date. International Journal of Pharmaceutical Compounding, 18(4), 348-350