

Formula Name KETOPROFEN/LIDOCAINE/CYCLOBENZAPRINE HCL ANHYDROUS GEL 20/5/2% GEL

Initials	AP	Quantity	100	MLS	Schedule	0	Days to Exp	180	Date of Formula	11/8/2021
Cost	\$63.13	Cost Per Unit	\$0.63	AWP	\$0.00	AWP Per Unit	\$0.00	Last Review	1/6/2016	

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
KETOPROFEN USP	PWD	20.0000	GMS	☐	\$7.40		SOLUBILITY: WATER(I), ALCOHOL(FS)
LIDOCAINE USP	PWD	5.0000	GMS	☐	\$1.28		SOLUBILITY: WATER (I) ALCOHOL (VS) 1. LIDOCAINE BASE IS BEST TO USE ON TOPICAL FORMULATIONS. HAS AN INCREASED LIPOPHILIC ACTIVITY OVER THE HCL.
CYCLOBENZAPRINE HCL USP	PWD	2.0000	GMS	☐	\$23.20		SOLUBILITY: WATER(FS), ALCOHOL(FS)
ETHYL ALCOHOL 95% USP	LIQU	10.0000	MLS	☐	\$0.00		MISCIBLE WITH WATER 1. THIS ALCOHOL IS NOT DENATURED. 2.**DO NOT USE DENATURED ALCOHOL IN ANY COMPOUNDED PRODUCT** SOLUBILITY:N/A
VERSAPRO (ANHYDROUS GEL BASE)	GEL	100.0000	GMS	☒	\$31.20		NONE
PROPYLENE GLYCOL USP	LIQU	3.5000	MLS	☐	\$0.05		SOLUBILITY:N/A

Therapy**Description**

Off-white/opaque colored cream, uniform consistency, no odor

Classification(s) Nonaqueous**Disease State(s)****Instructions**

A. Calculations/Special Considerations:

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log

2. Measure out ingredients

3. Triturate Ketoprofen, Lidocaine, and Cyclobenzaprine HCL in mortar.

4. Wet step 3 with Ethyl Alcohol and Propylene Glycol (for 240 mL or less: use syringe to syringe method; for 240+ mL: use mixer on lowest speed)

5. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe, add plunger, and expel air

6. In new syringe fill with Anhydrous base add plunger loosely, invert and tap Anhydrous base to the bottom, once tip is clear of Anhydrous base, fully insert plunger and expel extra air

7. Attach syringe from step 5 to Anhydrous base syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (complete 40 transfers)

8. Package and label with barcode label and affix "external use only" label

9. Tape labels

10. Have compound certified pharmacist perform final verification

D. Labeling:

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Label with lot number and expiration barcode and ingredient(s) label.
External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's
Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Off-white/opaque colored cream, uniform consistency, no odor

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist