

Formula Name KETOPROFEN/LIDOCAINE/CYCLOBENZAPRINE HCL GEL INACTIVE 20/5/2% GEL

Initials **Quantity** 100 **MLS** **Schedule** 0 **Days to Exp** 30 **Date of Formula** 8/24/2021
Cost \$31.88 **Cost Per Unit** \$0.32 **AWP** **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
LIPODERM BASE	CRM	0.0000	GMS	☒	\$0.00		NONE
KETOPROFEN USP	PWD	20.0000	GMS	☐	\$7.40		SOLUBILITY: WATER(I), ALCOHOL(FS)
LIDOCAINE USP	PWD	5.0000	GMS	☐	\$1.28		SOLUBILITY: WATER (I) ALCOHOL (VS) 1. LIDOCAINE BASE IS BEST TO USE ON TOPICAL FORMULATIONS. HAS AN INCREASED LIPOPHILIC ACTIVITY OVER THE HCL.
CYCLOBENZAPRINE HCL USP	PWD	2.0000	GMS	☐	\$23.20		SOLUBILITY: WATER(FS), ALCOHOL(FS)
ETHYL ALCOHOL 95% USP	LIQU	10.0000	MLS	☐	\$0.00		MISCIBLE WITH WATER 1. THIS ALCOHOL IS NOT DENATURED. 2.***DO NOT USE DENATURED ALCOHOL IN ANY COMPOUNDED PRODUCT*** SOLUBILITY:N/A
DISTILLED WATER	LIQU	3.5000	MLS	☐	\$0.00		NONE

Therapy**Description**

Off-white/opaque colored cream, uniform consistency, no odor

Classification(s)

Disease State(s)

Instructions

A. Calculations/Special Considerations:

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log

2. Measure out ingredients

3. Triturate Ketoprofen, Lidocaine, and Cyclobenzaprine HCL in mortar.

4. Wet step 3 with Ethyl Alcohol and Distilled Water (for 240 mL or less: use syringe to syringe method; for 240+ mL: use mixer on lowest speed)

5. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe, add plunger, and expel air

6. In new syringe fill with lipoderm and expel air and extra lipoderm so the sum of both syringes equals the final volume

7. Attach syringe from step 5 to lipoderm syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (typically 20 transfers)

8. Package and label with barcode label and affix "external use only" label

9. Tape labels

10. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rxs

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Off-white/opaque colored cream, uniform consistency, no odor

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 30 days

I. Formula Source & BUD Testing:

USP 795

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist