

Formula Name LEVETIRACETAM SUPP INACTIVE 1000MG SUPP

Initials ETJ **Quantity** 100 EACH **Schedule** 0 **Days to Exp** 90 **Date of Formula** 1/14/2016
Cost \$210.40 **Cost Per Unit** \$2.10 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
SPG SUPPOSI-BASE	BASE	106.0000	GMS	☐	\$2.82		SOLUBILITY:N/A
SILICA GEL (PPTD-MICRONIZED)	PWD	2.0000	GMS	☐	\$0.58		SOLUBILITY:N/A
LEVETIRACETAM USP	PWD	100.0000	GMS	☐	\$207.00		SOLUBILITY: 1:0.4 WATER(VS), 1:5 ALCOHOL(FS)

Therapy**Description**

Off-white/cream colored semi-solid, smooth to touch bullet shaped

Classification(s)**Disease State(s)****Instructions****Equipment Needed:**

Beakers, weigh boats, scales, suppository mold, Pam spray, paper towels, microwave/hot plate, spin bars, labels and plastic containers

Labelling & Storage:

Label with lot number and expiration barcode and ingredient(s) label.

Package in amber unit dose blisters with foil label.

Special Considerations:

None

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Melt SPG Base on the hot plate or in the microwave until fully liquified (~1 1/2 minutes then in 15 second intervals, if necessary)
4. Triturate Levetiracetam powder in mortar
5. Add Levetiracetam powder and Silica gel to beaker on hot plate at 120 degrees F and disperse in liquified base using stir bar. Let spin until dissolved (~15min)
6. Pour mixture into mold and, using the spatula, spread the mixture until each mold is full. Fill the center holes initially then move outwards. Once finished with one half push any excess to the other half
7. Fill remaining half of mold
8. Allow to harden slightly and then refrigerate for at least 120 minutes
9. Remove from fridge and let sit for at least 120 minutes, push in each suppository until it clicks, then remove clamps
10. Package and label suppositories in amber suppository unit dose blisters and foil label

Formula Source:

PCCA 11803

Testing:

PCCA 11803

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist