

Formula Name LIDOCAINE/HYDROCORTISONE SUPP 2/1% SUPP

Initials AP **Quantity** 100 EACH **Schedule** 0 **Days to Exp** 180 **Date of Formula** 2/10/2021
Cost \$7.83 **Cost Per Unit** \$0.08 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
SPG SUPPOSI-BASE	BASE	182.0000	GMS	☐	\$3.90		SOLUBILITY:N/A
SILICA GEL (PPTD-MICRONIZED)	PWD	2.0000	GMS	☐	\$0.34		SOLUBILITY:N/A
LIDOCAINE USP	PWD	3.7500	GMS	☐	\$0.96		SOLUBILITY: WATER (I) ALCOHOL (VS) 1. LIDOCAINE BASE IS BEST TO USE ON TOPICAL FORMULATIONS. HAS AN INCREASED LIPOPHILIC ACTIVITY OVER THE HCL.
HYDROCORTISONE USP MICONIZED (CORTISOL)	PWD	1.8800	GMS	☐	\$2.63		SOLUBILITY: WATER (I) ALCOHOL 1:40 PROTECT FROM LIGHT WHITE OR ALMOST WHITE ODORLESS CRYSTALLINE POWDER CORTICOSTEROID

Therapy**Description**

Off-white/cream colored semi-solid, smooth
to touch bullet shaped

Classification(s) Nonaqueous

Disease State(s)

Instructions

A. Calculations/Special Considerations:
None

B. Equipment Required:

Beakers, weigh boats, scales, suppository mold, mortar & pestle, paper towels, microwave/hot plate, spin bars, labels and plastic containers

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Melt SPG Base on the hot plate or in the microwave until liquified (~45 seconds then in 15 second intervals, if necessary)
4. Add Lidocaine, Hydrocortisone, and Silica gel to mortar and triturate to a fine powder
5. Add powder to beaker on hot plate (set at 40°C) and disperse in liquified base using stir bar. Let spin until dissolved (~15min)
6. Pour mixture into mold and, using the spatula, spread the mixture until each mold is full
7. Allow to harden at room temperature until solid (at least 2 hours)
8. Remove bolts, push down each suppository until it clicks, place mold on side and separate top mold by applying downward pressure to the lower mold
9. Package suppositories in amber suppository unit dose blisters
10. Have compound certified pharmacist perform final verification
11. Label blisters with prepack label foil sticker

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label

Add prepack label if compound lot made to stock for multiple Rx

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:

Off-white/cream colored semi-solid, smooth to touch bullet shaped

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Number of suppositories made is within 5% of intended batch size

Weigh 10% of batch (weighing each suppository individually) and confirm individual masses are within acceptable range.

Theoretical weight of suppository 1.896g

Acceptable weight range of sampled suppositories 1.801g - 1.991g

Weight #1: _____

Weight #2: _____

Weight #3: _____

Weight #4: _____

Weight #5: _____

Weight #6: _____

Weight #7: _____

Weight #8: _____

Weight #9: _____

Weight #10: _____

Pharmacist signature: _____

G. Packaging Container:

Package in UV light protective amber unit dose blisters with foil label.

H. Storage Requirements

Store refrigerated (36° - 46 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Formulation - PCCA 11432

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3

INGREDIENT WEIGHT #4

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist