

Formula Name LIDOCAINE/MICONAZOLE/ZINC OXIDE/ANTACID CREAM 3:3:3:1 UNG

Initials **Quantity** 200 **GMS** **Schedule** 0 **Days to Exp** 35 **Date of Formula** 5/25/2023
Cost \$5.64 **Cost Per Unit** \$0.03 **AWP** **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
LIDOCAINE 2% JELLY	GEL	60.0000	GMS	☐	\$0.00		NONE
MICONAZOLE 2%	CRM	60.0000	GMS	☐	\$4.84		
ZINC OXIDE 20% OINT USP	OINT	60.0000	GMS	☐	\$0.68		NONE
ANTACID LIQUID REGULAR STRENGTH	SUSP	20.0000	MLS	☐	\$0.13		SOLUBILITY:1:0.45 WATER(VS) , ALCOHOL(SL)

Therapy**Description**

Off-white opaque viscous semi-solid

Classification(s)**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

B. Equipment Required:

Wax paper, mixer, plastic spatula, metal spatula, ointment jar, graduated cylinder

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients using wax paper so that balances are not damaged
3. Combine ingredients using geometric dilution on wax paper or using Kitchen Aid mixer
4. Scrape wax paper/bowl with spatula and package in ointment jar (tap to lower ointment as needed)
5. Package and label with barcode label and affix "external use only" label
6. Tape labels
7. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

 (place label here) **E. Description of Finished Product:**

Off-white opaque viscous semi-solid

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Weight of final product is within 5% of intended batch size

Final Gross Weight: g - Weight of Ointment Jar: g = Final Net Weight: gPharmacist signature: **G. Packaging Container:**

Package in white ointment jar

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 35 days

I. Formula Source & BUD Testing

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3