

**Formula Name METOPROLOL ANHYDROUS SUSP 25MG/5ML SUSP**

**Initials** AP **Quantity** 30 **MLS** **Schedule** 0 **Days to Exp** 90 **Date of Formula** 11/26/2021  
**Cost** \$0.03 **Cost Per Unit** \$0.00 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

**Chemical Ingredients**

| Chemical Name                         | Form | Qty    | Units | QS                                  | Est Cost | Ha | Note                                 |
|---------------------------------------|------|--------|-------|-------------------------------------|----------|----|--------------------------------------|
| METOPROLOL 50MG TABLETS               | TAB  | 3.0000 | TABS  | <input type="checkbox"/>            | \$0.00   |    | NONE                                 |
| PROPYLENE GLYCOL USP                  | LIQU | 2.0000 | MLS   | <input type="checkbox"/>            | \$0.03   |    | SOLUBILITY:N/A                       |
| ANHYDROUS SUSPENDING AGENT (UNISPEND) | SUSP | 0.0000 | MLS   | <input checked="" type="checkbox"/> | \$0.00   |    | SOLUBILITY:N/A<br>SWEET ORAL VEHICLE |

**Therapy****Description**

Pink opaque suspension with fine suspended particles

**Classification(s)**Nonaqueous Oral**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

**B. Equipment Required:**

Counting tray, Mortar &amp; pestle, graduated cylinder, 60mL Luer Lock syringe, Stir bar, stir plate, Beaker, amber bottle, tape

**C. Method of Preparation:**

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. In a mortar, triturate Metoprolol tartrate tablets into a fine powder using the pestle
4. Wet powder with propylene glycol
5. Add about 15% of final volume of Anhydrous suspension to mortar and briefly mix
6. Pour suspension into a beaker (if small batch may use syringe to syringe method add suspension to 60mL syringe with black stopper at base)
7. Rinse mortar and add remaining amount of Anhydrous suspension required to Q.S.
8. Add stir bar to beaker, place on stir plate and spin until uniform (If using syringe to syringe method, mix suspension until uniform)
9. Have compound certified pharmacist perform final verification
10. Transfer contents to amber bottle
11. Label with RX label and "Shake well" "Refrigerate" auxilliary label
12. Tape Labels

**D. Labeling:**

Label with lot number and expiration barcode and ingredient(s) label.

Store in refrigerator auxillary sticker

Shake Well auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

---

(place label here)

---

**E. Description of Finished Product:**

Pink opaque suspension with fine suspended particles

**F. Quality Control Procedures:**

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist

Pharmacist signature: \_\_\_\_\_

G. Packaging Container:

Bulk store in UV protective amber bottle

Dispense in UV protective amber bottle with volume graduations

H. Storage Requirements:

Store refrigerated (36° - 46 °F)

Maximum BUD - 90 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

---

|                                                                                                                                                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------|
| Appropriate References Should Be<br>Consulted. Safety And Efficacy Of<br>These Formulas Are The<br>Responsibility Of The Compounding<br>Pharmacist |
|----------------------------------------------------------------------------------------------------------------------------------------------------|