

Formula Name MICAFUNGIN 100MG IN 100ML OF 0.9% SODIUM CHLORIDE INJ Sol (MiniBag) 1MG/ML INJ

Initials **Quantity** 100 **MLS** **Schedule** 0 **Days to Exp** 30 **Date of Formula** 7/19/2023
Cost \$0.00 **Cost Per Unit** \$0.00 **AWP** **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
MICAFUNGIN 100MG INJ VIAL	PWD	1.0000	VIAL	☐	\$0.00		
SODIUM CHLORIDE 0.9% 100ML INJ SOL (MiniBag)	INJ	100.0000	MLS	☐	\$0.00		

Therapy**Description**

Clear colorless solution free of particulate matter

Classification(s)**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

B. Equipment Required:

Alcohol Pad, Sterile gloves

C. Method of Preparation:

1. Perform hand-hygiene procedure by wetting hands and forearms and washing using soap and warm water for at least 30 seconds. Dry hands with a lint-free towel.
2. Don sterile gloves and use waterless alcohol-based surgical hand scrub.
3. Clean and disinfect the isolator hood according to proper policy and procedures.
4. Spray down all compounding items (needles, syringes, alcohol pads, stock bottles, etc) with 70% isopropyl alcohol (IPA), dry them and place them in the isolator hood
5. Place a second set of gloves over the gloves in the isolator hood.
6. Disinfect injection ports and stoppers with 70% IPA.
7. Connect 1 vial of Micafungin to mini-bag of sodium chloride 0.9% sol.
8. Label the bag with the appropriate pharmacy label and appropriate auxiliary label.

D. Labeling:

Record Manufacturer, Lot Number, NDC, and Expiration date for Micafungin

Record Manufacturer, Lot Number, NDC, and Expiration date for stock 0.9% Sodium Chloride Bag

Label with lot number and expiration label

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxiliary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Clear colorless solution free of particulate matter

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Container-closure integrity is verified to be free of any defect

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

Dispense in IV admixture bag

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

H. Storage Requirements

Store at room temperature (68° - 77 °F)

Maximum BUD - 30 days after connection before reconstitution

I. Formula Source & BUD Testing:

USP 797