

Formula Name MILRINONE 200MG IN 500ML OF 5% DEXTROSE INJ Sol (BAG) 0.4MG/ML INJ

Initials **Quantity** 500 **MLS** **Schedule** 0 **Days to Exp** 14 **Date of Formula** 7/20/2023
Cost \$0.00 **Cost Per Unit** \$0.00 **AWP** **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
MILRINONE 50MG/50ML INJ VIAL	INJ	20.0000	MLS	☐	\$0.00		
DEXTROSE 5% SOLUTION 500ML	SOLN	500.0000	MLS	☐	\$0.00		

Therapy**Description**

Clear colorless solution free of particulate matter

Classification(s)**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

B. Equipment Required:

60ml syringe (4), transfer needles (4), Alcohol Pad, Port protector sticker,

C. Method of Preparation:

1. Perform hand-hygiene procedure by wetting hands and forearms and washing using soap and warm water for at least 30 seconds. Dry hands with a lint-free towel.
2. Don sterile gloves and use waterless alcohol-based surgical hand scrub.
3. Clean and disinfect the isolator hood according to proper policy and procedures.
4. Spray down all compounding items (needles, syringes, alcohol pads, stock bottles, etc) with 70% IPA, dry them and place them in the isolator hood.
5. Place a second set of gloves over the gloves in the isolator hood.
6. Disinfect injection ports and stoppers with 70% IPA.
7. Using 60ml syringes, remove 200ml of fluid from 5% dextrose bag.
8. Pull up 200ml from milrinone vials.
9. Inject 200ml of milrinone into the D5W bag.
10. Cover entry port with port protector sticker.
11. Label the bag with the appropriate pharmacy label.

D. Labeling:

Record Manufacturer, Lot Number, NDC, and Expiration date for Milrinone

Record Manufacturer, Lot Number, NDC, and Expiration date for stock 5% Dextrose Bag

Label with lot number and expiration label

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

 (place label here) **E. Description of Finished Product:**

Clear colorless solution free of particulate matter

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Container-closure integrity is verified to be free of any defect

Volume of final product is within 5% of intended batch size

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Pharmacist signature: _____

G. Packaging Container:

Dispense in IV admixture bag covered by a UV protective foil bag

H. Storage Requirements

Store at room temperature (68° - 77 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

USP 797

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist