

Formula Name ABH DOUBLE GEL INACTIVE 2/50/2MG/ML GEL

Initials AP **Quantity** 60 **MLS** **Schedule** 4 **Days to Exp** 14 **Date of Formula** 2/2/2021
Cost \$21.97 **Cost Per Unit** \$0.37 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
DIPHENHYDRAMINE HCL USP	PWD	3.0000	GMS	☐	\$1.34		SOLUBILITY: WATER(S), ALCOHOL(S)
HALOPERIDOL USP	PWD	0.1200	GMS	☐	\$0.65		SOLUBILITY: 1:10,000 WATER (VSS), 1:60 ALCOHOL(SP)
LORAZEPAM USP	PWD	0.1200	GMS	☐	\$5.40		NONE
ETHOXY DIGLYCOL	LIQU	5.0000	MLS	☐	\$0.53		SOLUBILITY:N/A SOLVENT
DISTILLED WATER	LIQU	1.0000	MLS	☐	\$0.00		NONE
LIPODERM BASE	CRM	60.0000	GMS	☒	\$14.04		NONE

Therapy**Description**

Off-white/opaque colored cream, uniform consistency, no odor

Classification(s) Non-Preserved Aqueous

Disease State(s)

Instructions

A. Calculations/Special Considerations:
None

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding and control usage logs
2. Measure out ingredients
3. Triturate Diphenhydramine, Haloperidol, and Lorazepam in mortar.
4. Wet Step 3 with Ethoxy Diglycol and Distilled Water (for 240 mL or less: use syringe to syringe method; for 240+ mL: use mixer on lowest speed)
5. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe
6. In new syringe fill with lipoderm, add plunger loosely, invert and tap lipoderm to the bottom, once tip is clear of lipoderm, fully insert plunger and expel extra air
7. Attach syringe from step 5 to lipoderm syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (typically 20 transfers)
8. Package and label with barcode label and affix "store in refrigerator" and "external use only" labels
9. Tape labels
10. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

Store in refrigerator auxillary sticker

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:

Off-white/opaque colored cream, uniform consistency, no odor

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store refrigerated (36° - 46 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

PCCA 9428

J. Testing:

USP 795