

Formula Name AMLODIPINE SUPP 2.5MG SUPP

Initials AP **Quantity** 100 EACH **Schedule** 0 **Days to Exp** 180 **Date of Formula** 2/10/2021
Cost \$5.89 **Cost Per Unit** \$0.06 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
SPG SUPPOSI-BASE	BASE	173.2000	GMS	☐	\$3.71		SOLUBILITY:N/A
SILICA GEL (PPTD-MICRONIZED)	PWD	2.0000	GMS	☐	\$0.34		SOLUBILITY:N/A
AMLODIPINE 2.5MG TAB	EACH	100.0000	TABS	☐	\$1.84		

Therapy**Description**

Off-white/cream colored semi-solid, smooth to touch bullet shaped

Classification(s)Nonaqueous**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

B. Equipment Required:

Beakers, weigh boats, scales, suppository mold, mortar & pestle, paper towels, microwave/hot plate, spin bars, labels and plastic containers

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Melt SPG Base on the hot plate or in the microwave until liquified (~45 seconds then in 15 second intervals, if necessary)
4. Triturate Amlodipine tablets in mortar & pestle, add Silica gel and triturate as well
5. Add Amlodipine and Silica gel mixture to beaker on hot plate (set at 40°C) and disperse in liquified base using stir bar. Let spin until dissolved (~15min)
6. Pour mixture into mold and, using the spatchula, spread the mixture until each mold is full
7. Allow to harden at room temperature until solid (at least 2 hours)
8. Remove bolts, push down each suppository until it clicks, place mold on side and separate top mold by applying downward pressure to the lower mold
9. Individually weigh at least 10% of suppositories to ensure the weight is within 10% of theoretical weight
10. Package suppositories in amber suppository unit dose blisters
11. Have compound certified pharmacist perform final verification
12. Label blisters with prepack label foil sticker

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Off-white/cream colored semi-solid, smooth to touch bullet shaped

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Number of suppositories made is within 5% of intended batch size

Weigh 10% of batch (weighing each suppository individually) and confirm individual masses are within acceptable range.

Theoretical weight of suppository 1.850g

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Acceptable weight range of sampled suppositories 1.665g - 2.035g

Weight #1: _____

Weight #2: _____

Weight #3: _____

Weight #4: _____

Weight #5: _____

Weight #6: _____

Weight #7: _____

Weight #8: _____

Weight #9: _____

Weight #10: _____

Pharmacist signature: _____

G. Packaging Container:

Package in UV light protective amber unit dose blisters with foil label.

H. Storage Requirements

Store refrigerated (36° - 46 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist