

Formula Name AMPICILLIN/SULBACTAM 1.5G IN 50ML OF 0.9% SODIUM CHLORIDE INJ Sol (BAG) 30MG/ML INJ 30MG/ML INJ

Initials AP **Quantity** 50 **MLS** **Schedule** 0 **Days to Exp** 2 **Date of Formula** 7/19/2023

Cost \$0.00 **Cost Per Unit** \$0.00 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
SODIUM CHLORIDE 0.9% 50ML INJ SOL	INJ	50.0000	MLS	☐	\$0.00		
AMPICILLIN / SULBACTAM 1.5G VIAL	PWD	1.0000	VIAL	☐	\$0.00		
STERILE WATER FOR INJECTION	SOLN	3.2000	MLS	☐	\$0.00		NONE

Therapy

Description Clear colorless to pale yellow solution free of particulate matter

Classification(s) CATEGORY 1 CSP

Disease State(s)

Instructions

A. Calculations/Special Considerations:

None

B. Equipment Required:

5mL Luer Lock Syringe, 22G Transfer Needle, Alcohol Pad, Port Seal

C. Method of Preparation:

1. Perform hand-hygiene procedure by wetting hands and forearms and washing using soap and warm water for at least 30 seconds. Dry hands with a lint-free towel.
2. Don sterile gloves and use waterless alcohol-based surgical hand scrub.
3. Clean and disinfect the isolator hood according to proper policy and procedures.
4. Spray down all compounding items (needles, syringes, alcohol pads, stock bottles, etc) with 70% isopropyl alcohol (IPA), dry them and place them in the isolator hood
5. Place a second set of gloves over the gloves in the isolator hood.
6. Disinfect injection ports and stoppers with 70% IPA.
7. Reconstitute 1.5G vial of ampicillin/sulbactam by SLOWLY adding 3.2ml of sterile water sol.
-ENSURE ALL POWDER IS WETTED BY GENTLY ROTATING THE VIAL
-ALLOW MIXED PRODUCT TO STAND TO DISSIPATE ANY FOAM
8. Transfer entire contents of the 1.5G vial into one 50ml sodium chloride 0.9% IV bag.
9. Remove air from the bag and cover the injection port.
10. Label the bag with the appropriate pharmacy label and refrigerator auxiliary label.

D. Labeling:

Record Manufacturer, Lot Number, NDC, and Expiration date for ampicillin/sulbactam

Record Manufacturer, Lot Number, NDC, and Expiration date for stock 0.9% Sodium Chloride Bag and Sterile Water vial

Label with lot number and expiration label

Store in refrigerator auxiliary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxiliary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Clear colorless to pale yellow solution free of particulate matter

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Container-closure integrity is verified to be free of any defect

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

Dispense in IV admixture bag covered by a UV protective foil bag with cold packs

H. Storage Requirements

Store refrigerated (36° - 46 °F)

Maximum BUD - 48 hours

I. Formula Source & BUD Testing:

USP 797

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist