

Formula Name ANTACID+/LIDOCAINE/DIPHENHYDRAMINE/NYSTATIN SUSP INACTIVE 1:1:1:1 SUSP

Initials ETJ **Quantity** 4000 **MLS** **Schedule** 0 **Days to Exp** 14 **Date of Formula** 1/7/2016
Cost \$87.35 **Cost Per Unit** \$0.02 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
PROPYLENE GLYCOL USP	LIQU	50.0000	MLS	☐	\$0.68		SOLUBILITY:N/A
LIDOCAINE USP	PWD	20.0000	GMS	☐	\$5.60		SOLUBILITY: WATER (I) ALCOHOL (VS) 1. LIDOCAINE BASE IS BEST TO USE ON TOPICAL FORMULATIONS. HAS AN INCREASED LIPOPHILIC ACTIVITY OVER THE HCL.
PRESERVED WATER (PARABENS)	LIQU	700.0000	MLS	☐	\$0.00		SEE FORMULA FOR PRESERVED WATER (PARABEN)
DIPHENHYDRAMINE HCL USP	PWD	2.5000	GMS	☐	\$0.28		SOLUBILITY: WATER(S), ALCOHOL(S)
NYSTATIN ORAL 100,000 U/ML SUSPENSION	SUSP	946.0000	MLS	☐	\$0.00		NONE
SIMPLE SYRUP	LIQU	946.0000	MLS	☐	\$24.10		NONE
XANTHAN GUM 1% SUSPENSION	SUSP	250.0000	MLS	☐	\$0.00		NONE
FLAVOR CHERRY CONCENTRATE *COLORLESS*	SOLN	60.0000	MLS	☐	\$8.09		NONE
FOOD COLOR RED LIQUID	SOLN	40.0000	GTTS	☐	\$48.61		NONE
ANTACID LIQUID REGULAR STRENGTH	SUSP	0.0000	MLS	☒	\$0.00		SOLUBILITY:1:0.45 WATER(VS) , ALCOHOL(SL)

Therapy**Description**Light salmon colored uniform suspension,
medium viscosity, bubblegum and mint scent**Classification(s)**Nonpreserved Aqueous**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

20 gtts = 1mL

B. Equipment Required:

Weigh boats; beakers; graduated cylinder; homogenizer (prn); funnel; scales; stir rod/bar; tape

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. To measure lidocaine cut off bottle tops with box cutter. Use small spatula to scrape into an appropriate beaker
4. Heat Propylene Glycol in microwave (~10 + 5 seconds; NOT 15 seconds)
5. Dissolve Lidocaine in Propylene Glycol using stir rod/bar (NOTE: Do NOT add this to the water solution yet)
6. Dissolve Diphenhydramine in Water using stir rod/bar
7. Add Nystatin, Simple syrup, Xanthan gum, Cherry Concentrate (Colorless), and Food Coloring to Water
8. Homogenize until evenly dispersed
9. SLOWLY add Lidocaine mixture being sure to homogenize with each addition
10. QS with Antacid+
11. Have compound certified pharmacist perform final verification
12. Pour into container
13. Homogenize / Shake well until evenly dispersed
14. Label with RX label and "Shake well" "Refrigerate" auxiliary label
15. Tape labels

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

D. Labelling:

Label with lot number and expiration barcode and ingredient(s) label.

Store in refrigerator auxillary sticker

Shake Well auxillary sticker

E. Description of Finished Product:

Light salmon colored uniform suspension, medium viscosity, bubblegum and mint scent

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

G. Packaging Container:

Bulk store in UV protective amber bottle

Dispense in UV protective amber bottle with volume graduations

H. Storage Requirements

Store refrigerated (36° - 46 °F)

Maximum BUD - 14 days

I. Formula Source:

J. Testing:

USP 795

Revised by: Austin Philipp

12/07/2020

Verified by:

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist