

**Formula Name BACLOFEN ANHYDROUS SUSP INACTIVE 5MG/ML SUSP**

**Initials** AP **Quantity** 1000 **MLS** **Schedule** 0 **Days to Exp** 90 **Date of Formula** 11/18/2021  
**Cost** \$110.80 **Cost Per Unit** \$0.11 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

**Chemical Ingredients**

| Chemical Name                         | Form | Qty       | Units | QS                                  | Est Cost | Ha | Note                                                                                        |
|---------------------------------------|------|-----------|-------|-------------------------------------|----------|----|---------------------------------------------------------------------------------------------|
| BACLOFEN USP                          | PWD  | 5.0000    | GMS   | <input type="checkbox"/>            | \$14.24  |    | SOLUBILITY: WATER (SS)<br>ALCOHOL (SS)<br>1. EXCELLENT CHEMICAL<br>FOR DIABETIC NEUROPATHY. |
| PROPYLENE GLYCOL USP                  | LIQU | 20.0000   | MLS   | <input type="checkbox"/>            | \$0.56   |    | SOLUBILITY:N/A                                                                              |
| ANHYDROUS SUSPENDING AGENT (UNISPEND) | SUSP | 1000.0000 | MLS   | <input checked="" type="checkbox"/> | \$96.00  |    | SOLUBILITY:N/A<br>SWEET ORAL VEHICLE                                                        |

**Therapy****Description**

White opaque suspension with fine suspended particles

**Classification(s)**

Nonaqueous Oral

**Disease State(s)****Instructions**

A. Calculations/Special Considerations:  
None

B. Equipment Required:  
Weigh boats; beakers; scales; mortar & pestle; homogenizer; class A graduated cylinder; funnel; tape

**C. Method of Preparation:**

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. In a mortar, triturate Baclofen into a fine powder using the pestle
4. Wet powder with propylene glycol
5. Add ~10mL of Anhydrous suspension to mortar and briefly mix
6. Pour mixture into beaker
7. Rinse mortar and pestle with ~10% of Anhydrous suspension and pour into beaker, repeat until all baclofen completely transferred from mortar and pestle
8. Add up to 90% of final volume to beaker, if large particles persist, use homogenizer to further mix
9. QS to final volume with Anhydrous suspension in class A graduated cylinder
10. Pour into amber bottle
11. Label with RX label and "Shake well" "Refrigerate" auxilliary label
12. Have compound certified pharmacist perform final verification
13. Tape Labels

**D. Labeling:**

Label with lot number and expiration barcode and ingredient(s) label.  
Store in refrigerator auxillary sticker  
Shake Well auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx  
Add Rx auxillary label if compound lot made to order for a unique Rx

\_\_\_\_\_ (place label here) \_\_\_\_\_

**E. Description of Finished Product:**

White opaque suspension with fine suspended particles

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: \_\_\_\_\_

G. Packaging Container:

Bulk store in UV protective amber bottle

Dispense in UV protective amber bottle with volume graduations

H. Storage Requirements:

Store refrigerated (36° - 46 °F)

Maximum BUD - 90 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1