

Formula Name BACLOFEN SF SUSP INACTIVE 20MG/5ML SUSP

Initials AP **Quantity** 1000 **MLS** **Schedule** 0 **Days to Exp** 14 **Date of Formula** 2/10/2021
Cost \$21.07 **Cost Per Unit** \$0.02 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
BACLOFEN USP	PWD	4.0000	GMS	☐	\$8.90		SOLUBILITY: WATER (SS) ALCOHOL (SS) 1. EXCELLENT CHEMICAL FOR DIABETIC NEUROPATHY.
FLAVOR TUTTI FRUTTI, ARTIFICIAL	SOLN	15.0000	MLS	☐	\$4.75		SOLUBILITY:N/A
SUSPENDIT LIQUID	SUSP	0.0000	MLS	☒	\$0.00		SOLUBILITY:N/A SWEET ORAL VEHICLE
FOOD COLOR YELLOW LIQUID	SOLN	2.0000	GTTS	☐	\$7.43		

Therapy**Description**

Thick orange opaque suspension with small suspended particles

Classification(s) Non-Preserved Aqueous

Disease State(s)

Instructions**A. Calculations/Special Considerations:**

None

B. Equipment Required:

Weigh boats; beakers; scales; mortar & pestle; class A graduated cylinder; funnel; tape

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. In a mortar, triturate Baclofen into a fine powder using the pestle
4. Add Suspendit to Step 3 slowly using 10mL's at a time while mixing thoroughly, then transfer to a beaker equipped with a spinbar. Use an amount of Suspendit that is approximately 10% of the final volume. Use a rubber spatula to thoroughly scrape the sides of the mortar during this process.
5. Rinse the Step 4 mortar with Suspendit. Use an amount of Suspendit that is approximately 10% of the final volume. Combine the rinsing into the Step 2 beaker, using the rubber spatula to thoroughly scrape the sides of the mortar during this process. Do this step a total of two times to completely rinse the mortar.
6. Add Tutti Frutti and Food Coloring to mixture, then add Suspendit to mixture to achieve ~90% of final volume, mix well.
7. In a Class A rated graduated cylinder, bring Step 6 to the final volume with Suspendit and mix well.
8. Pour into container
9. Label with a compound label, individual bar scan labels, and "Shake Well" sticker
10. Have compound certified pharmacist perform final verification
11. Shake well until evenly dispersed
12. Tape Labels

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

Store in refrigerator auxillary sticker

Shake Well auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:

Thick orange opaque suspension with small suspended particles

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

Bulk store in UV protective amber bottle

Dispense in UV protective amber bottle with volume graduations

H. Storage Requirements:

Store refrigerated (36° - 46 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

PCCA Formula 11199

He, Y., Brunstrom-Hernandez, J. E., Thio, L. L., Lackey, S., Gaebler-Spira, D., Kuroda, M. M., ... Jusko, W. J. (2014). Population Pharmacokinetics of Oral Baclofen in Pediatric Patients with Cerebral Palsy. *The Journal of Pediatrics*, 164(5), 1181–1188.e8. doi: 10.1016/j.jpeds.2014.01.029

Vadlamudi, N. B., Hitch, M. C., Dimmitt, R. A., & Thame, K. A. (2013). Baclofen for the Treatment of Pediatric GERD. *Journal of Pediatric Gastroenterology and Nutrition*, 57(6), 808-812. doi: 10.1097/MPG.0b013e3182a2747b