

Formula Name BANHURT SUSP (PREDNISONE/TETRACYCLINE/DIPHENHYDRAMINE/NYSTATIN)
5/240/25mg/500k/30mL SUSP

Initials AP **Quantity** 3000 **MLS** **Schedule** 0 **Days to Exp** 35 **Date of Formula** 1/24/2022
Cost \$12.55 **Cost Per Unit** \$0.00 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
PREDNISONE 50MG	TAB	10.0000	TABS	☐	\$0.00		NONE
TETRACYCLINE HCL	PWD	24.0000	GMS	☐	\$0.00		SOLUBILITY: 1:10 WATER(FS), 1:100 ALCOHOL(SS)
GLYCERIN USP (NATURAL)	LIQU	20.0000	MLS	☐	\$2.13		SOLUBILITY:N/A
DIPHENHYDRAMINE HCL 12.5MG/5ML ORAL SOLUTION	SOLN	1000.0000	MLS	☐	\$10.42		NONE
ORA-BLEND	SUSP	0.0000	MLS	☒	\$0.00		SOLUBILITY:N/A SWEET ORAL VEHICLE
NYSTATIN ORAL 100,000 U/ML SUSPENSION	SUSP	500.0000	MLS	☐	\$0.00		NONE
PRESERVED WATER (PARABENS)	LIQU	600.0000	MLS	☐	\$0.00		SEE FORMULA FOR PRESERVED WATER (PARABEN)

Therapy

Description

Pinkish-red opaque suspension, medium viscosity

Classification(s) Preserved Aqueous

Disease State(s)

Instructions

A. Calculations/Special Considerations:
None

B. Equipment Required:

Weigh boats; beakers; graduated cylinder; mortar/pestle; stir bar, stir plate, homogenizer (prn); funnel; tape

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Triturate Prednisone tablets in mortar with pestle to a fine powder
4. Add Tetracycline HCl powder to prednisone using geometric dilution
5. Add Glycerin to form a smooth transferrable consistency
6. Add Diphenhydramine to a large beaker on a stir plate and spin.
7. Add Prednisone and Tetracycline mixture from step 5 to beaker
8. Rinse glassware to remove any active powder residue with Ora-Blend as needed.
9. Add water
10. Add Nystatin suspension
11. Add Ora-Blend until ~90% of final volume is achieved
12. Homogenize until evenly dispersed
13. QS to final volume with Ora-Blend
14. Have compound certified pharmacist perform final verification
15. Transfer to amber bottle
16. Shake well until evenly dispersed
17. Label with RX label and "Shake well" "Refrigerate" auxilliary label
18. Tape labels

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

Store in refrigerator auxillary sticker

Shake Well auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Pinkish-red opaque suspension, medium viscosity

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

Bulk store in UV protective amber bottle

Dispense in UV protective amber bottle with volume graduations

H. Storage Requirements:

Store refrigerated (36° - 46 °F)

Maximum BUD - 35 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1