

Formula Name BHR GEL INACTIVE 25/1/10MG/ML GEL

Initials AP **Quantity** 60 **MLS** **Schedule** 0 **Days to Exp** 14 **Date of Formula** 2/2/2021
Cost \$7.22 **Cost Per Unit** \$0.12 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
DIPHENHYDRAMINE HCL USP	PWD	1.5000	GMS	☐	\$0.67		SOLUBILITY: WATER(S), ALCOHOL(S)
HALOPERIDOL USP	PWD	0.0600	GMS	☐	\$0.33		SOLUBILITY: 1:10,000 WATER (VSS), 1:60 ALCOHOL(SP)
METOCLOPRAMIDE HCL USP	PWD	0.7090	GMS	☐	\$2.98		SOLUBILITY: 1:0.7 WATER(VS), 1:3 ALCOHOL(FS)
ETHOXY DIGLYCOL	LIQU	5.0000	MLS	☐	\$0.53		SOLUBILITY:N/A SOLVENT
POLOX 20% GEL	GEL	60.0000	GMS	☒	\$2.27		NONE
LIPMAX LECITHIN & ISOPROPYL PALMITATE	SOLN	12.0000	MLS	☐	\$0.44		

Therapy**Description**

Solid yellow, oily, highly viscous organogel

Classification(s)Non-Preserved Aqueous**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

Metoclopramide 1mg = Metoclopramide HCl 1.182 mg

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Triturate Diphenhydramine, Haloperidol, and Metoclopramide in mortar.
4. Wet step 3 with Ethoxy Diglycol (for 240 mL or less: use syringe to syringe method; for 240+ mL: use mixer on lowest speed)
5. Add Lipmax and mix until evenly dispersed
6. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe
7. In the same syringe QS with polox, add plunger loosely, invert and tap lipoderm to the bottom, once tip is clear of lipoderm, fully insert plunger and expel extra air
8. Attach a new syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (typically 20 transfers)
9. Package and label with barcode label and affix "external use only" label
10. Tape labels
11. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here)_____

E. Description of Finished Product:

Solid yellow, oily, highly viscous organogel

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

PCCA 3226

PCCA 2248