

Formula Name MORPHINE SULF./LIDOCAINE HCL IN HYDROGEL 0.5/5% GEL

Initials AP **Quantity** 60 **GMS** **Schedule** 2 **Days to Exp** 35 **Date of Formula** 1/22/2024
Cost \$2.90 **Cost Per Unit** \$0.05 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
HYDROGEL	GEL	0.0000	GMS	☒	\$0.00		
PRESERVED WATER (PARABENS)	LIQU	5.0000	MLS	☐	\$0.00		SEE FORMULA FOR PRESERVED WATER (PARABEN)
MORPHINE SULFATE (PENTAHYDRATE) USP	PWD	0.3000	GMS	☐	\$2.06		SOLUBILITY: 1:16 WATER(S), 1:570 ALCOHOL(SL)
LIDOCAINE HCL USP	PWD	3.0000	GMS	☐	\$0.84		SOLUBILITY: WATER(VS), ALCOHOL(VS)

Therapy**Description**

White opaque gel, medium viscosity, uniform consistency with suspended air bubbles

Classification(s)

Preserved Aqueous

Disease State(s)**Instructions**

A. Calculations/Special Considerations:
None

B. Equipment Required:
Beakers; weigh boats; mortar& pestle, rubber spatula.
For batches 240 mL or less: 60 mL syringes, luer lock adapters.
For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding and control usage logs
2. Measure out ingredients (Note: cut the back end off of the Hydrogel tube and squeeze it out through the new opening. If necessary, use a spatula to scrape any remaining gel out of the tube.)
3. Dissolve Lidocaine HCl and Morphine in Preserved Water
4. Add the Lidocaine HCl and Morphine mixture from step 4 to a 60ml syringe
5. Fill another 60ml syringe with the hydrogel. ***QS the Hydrogel so both syringes equal the final volume. Ex 5ml in active syringe and 55ml in hydrogel syringe = 60ml QS
6. Mix using the syringe to syringe method, completing 40 transfers
7. Package and label with barcode and affix "external use only" & "refrigerate" label
8. Tape labels
9. Have compound certified pharmacist perform final verification

D. Labelling:

Label with lot number and expiration barcode and ingredient(s) label.
External use only auxillary sticker
Refrigerate sticker

E. Description of Finished Product:

White opaque gel, medium viscosity, uniform consistency with suspended air bubbles

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist
Volume of final product is within 10% of intended batch size

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag
May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

H. Storage Requirements

Store at refrigerated temperature (36°–46 °F)

Maximum BUD - 35 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist