

**Formula Name MORPHINE SULFATE SUPP INACTIVE 10MG SUPP**

**Initials** ETJ **Quantity** 100 EACH **Schedule** 2 **Days to Exp** 90 **Date of Formula** 1/26/2016  
**Cost** \$7.26 **Cost Per Unit** \$0.07 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

**Chemical Ingredients**

| Chemical Name                       | Form | Qty      | Units | QS                       | Est Cost | Ha | Note                                            |
|-------------------------------------|------|----------|-------|--------------------------|----------|----|-------------------------------------------------|
| SPG SUPPOSI-BASE                    | BASE | 183.0000 | GMS   | <input type="checkbox"/> | \$3.92   |    | SOLUBILITY:N/A                                  |
| SILICA GEL (PPTD-MICRONIZED)        | PWD  | 2.0000   | GMS   | <input type="checkbox"/> | \$0.34   |    | SOLUBILITY:N/A                                  |
| MORPHINE SULFATE (PENTAHYDRATE) USP | PWD  | 1.0000   | GMS   | <input type="checkbox"/> | \$3.00   |    | SOLUBILITY: 1:16 WATER(S),<br>1:570 ALCOHOL(SL) |

**Therapy****Description**

Off-white/cream colored semi-solid, smooth to touch bullet shaped

**Classification(s)****Disease State(s)****Instructions****Equipment Needed:**

Beakers, weigh boats, scales, suppository mold, Pam spray, paper towels, microwave/hot plate, spin bars, labels and plastic containers

**Labelling & Storage:**

Label with lot number and expiration barcode and ingredient(s) label.

Package in amber unit dose blisters with foil label.

**Special Considerations:**

None

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Melt SPG Base on the hot plate or in the microwave until fully liquified (~1 1/2 minutes then in 15 second intervals, if necessary)
4. Add Morphine and Silica gel to beaker on hot plate and disperse in liquified base using stir bar. Let spin until dissolved (~15min)
5. Spray the suppository mold generously with Pam; turn upside down to tap out any extra Pam. Set mold on hot plate until warm to touch
6. Pour mixture into mold and, using the spatchula, spread the mixture until each mold is full
7. Allow to harden slightly and then refrigerate for at least 90 minutes
8. Remove from fridge, push in each suppository until it clicks, then remove clamps
9. Package and label suppositories in amber suppository unit dose blisters and foil label

**Formula Source:**

PCCA 9171

**Testing:**

PCCA 9171

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist