

**Formula Name PROMETHAZINE HCl 25MG/ML GEL INACTIVE 25MG/ML GEL**

**Initials** AP **Quantity** 120 **MLS** **Schedule** 0 **Days to Exp** 90 **Date of Formula** 2/2/2021  
**Cost** \$36.19 **Cost Per Unit** \$0.30 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

**Chemical Ingredients**

| Chemical Name                 | Form | Qty      | Units | QS                       | Est Cost | Ha | Note                  |
|-------------------------------|------|----------|-------|--------------------------|----------|----|-----------------------|
| PROMETHAZINE HCL USP          | PWD  | 2.5000   | GMS   | <input type="checkbox"/> | \$2.56   |    | SOLUBILITY: WATER(VS) |
| PROPYLENE GLYCOL USP          | LIQU | 3.0000   | MLS   | <input type="checkbox"/> | \$0.03   |    | SOLUBILITY:N/A        |
| VERSAPRO (ANHYDROUS GEL BASE) | GEL  | 120.0000 | GMS   | <input type="checkbox"/> | \$33.60  |    | NONE                  |

**Therapy****Description**

Off-white/opaque colored cream, uniform consistency, no odor

**Classification(s)**

Nonaqueous

**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

**B. Equipment Required:**

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

**C. Method of Preparation:**

1. Calculate required amounts and fill out compounding log

2. Measure out ingredients

3. Add Promethazine HCl to mortar & pestle, triturate, and wet with Propylene Glycol

4. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe

5. In new syringe fill with PermE8, add plunger loosely, invert and tap PermE8 to the bottom, once tip is clear of PermE8, fully insert plunger and expel extra air

6. Attach syringe from step 5 to PermE8 syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (typically 20 transfers)

7. Package and label with barcode label and affix "store in refrigerator" and "external use only" labels

8. Tape labels

9. Have compound certified pharmacist perform final verification

**D. Labeling:**

Label with lot number and expiration barcode and ingredient(s) label.

Store in refrigerator auxillary sticker

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

\_\_\_\_\_ (place label here) \_\_\_\_\_

**E. Description of Finished Product:**

Off-white/opaque colored cream, uniform consistency, no odor

**F. Quality Control Procedures**

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist

Pharmacist signature: \_\_\_\_\_

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store refrigerated (36° - 46 °F)

Maximum BUD - 90 days

I. Formula Source & BUD Testing:

USP 795

---

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist