

**Formula Name DIAZEPAM SUPP SHELL (INACTIVE) 15MG SUPP**

**Initials** AP    **Quantity** 60 EACH    **Schedule** 4    **Days to Exp** 180    **Date of Formula** 2/10/2021  
**Cost** \$4.99    **Cost Per Unit** \$0.08    **AWP** \$0.00    **AWP Per Unit** \$0.00    **Last Review** 1/6/2016

**Chemical Ingredients**

| Chemical Name               | Form | Qty      | Units | QS                       | Est Cost | Ha | Note           |
|-----------------------------|------|----------|-------|--------------------------|----------|----|----------------|
| SPG SUPPOSI-BASE            | BASE | 112.0020 | GMS   | <input type="checkbox"/> | \$3.15   |    | SOLUBILITY:N/A |
| SILICA GEL (PTD-MICRONIZED) | PWD  | 1.2000   | GMS   | <input type="checkbox"/> | \$0.40   |    | SOLUBILITY:N/A |
| DIAZEPAM USP                | PWD  | 0.9000   | GMS   | <input type="checkbox"/> | \$1.44   |    | NONE           |

**Therapy****Description**

Off-white/cream colored semi-solid, smooth to touch bullet shaped suppository encased in white shell

**Classification(s)**Nonaqueous**Disease State(s)****Instructions**

A. Calculations/Special Considerations:  
None

**B. Equipment Required:**

Beakers, weigh boats/paper, spatulas and spoons, weight scale, mortar and pestle, microwave/hot plate, 60mL luer lock syringe & syringe cap, spin bar, spin bar catcher, suppository shells, shell stand, suppository backlight, suppository sealer, labels, ziplock bag

**C. Method of Preparation:**

1. Calculate required amounts and fill out compounding log and control logs
2. Measure out ingredients
3. Melt SPG Base on the hot plate or in the microwave until liquified (~45 seconds then in 15 second intervals, if necessary)
4. Add Diazepam and Silica gel to mortar and triturate to a fine powder
5. Add powder to beaker on hot plate (set at 40°C) and disperse in liquified base using stir bar. Let spin until dissolved (~15min)
6. Set empty shells on stand and light backlight
7. Pour mixture into stoppered 60mL syringe, insert plunger, and fill each shell to the fill line - shake syringe every 5 suppositories to ensure homogenous mixture
8. Allow filled shells to harden at room temperature until solid (at least 2 hours)
9. Have compound certified pharmacist perform final verification
10. Enclose each shell with sealer, ensuring the entire width of the shell is sufficiently enclosed
11. Individually weigh at least 10% of suppositories to ensure the weight is within 10% of theoretical weight
12. Label suppositories horizontally along the sealed area of the shell and store in a plastic bag

**D. Labeling:**

Label with lot number and expiration barcode and ingredient(s) label

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxiliary label if compound lot made to order for a unique Rx

\_\_\_\_\_ (place label here) \_\_\_\_\_

**E. Description of Finished Product:**

Off-white/cream colored semi-solid, smooth to touch bullet shaped suppository encased in white shell

**F. Quality Control Procedures**

Physical appearance of compound verified by compounding certified pharmacist

Number of suppositories made is within 5% of intended batch size

Weigh 10% of batch (weighing each suppository individually) and confirm individual masses are within acceptable range.

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist

Theoretical weight of suppository in sealed shell 2.208g  
Acceptable weight range of sampled suppositories 1.987g - 2.429g

Weight #1: \_\_\_\_\_

Weight #2: \_\_\_\_\_

Weight #3: \_\_\_\_\_

Weight #4: \_\_\_\_\_

Weight #5: \_\_\_\_\_

Weight #6: \_\_\_\_\_

Weight #7: \_\_\_\_\_

Weight #8: \_\_\_\_\_

Weight #9: \_\_\_\_\_

Weight #10: \_\_\_\_\_

Pharmacist signature: \_\_\_\_\_

G. Packaging Container:

Package suppository shell in plastic bag with label.

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Formulation - PCCA 9919

BUD- USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3

Appropriate References Should Be  
Consulted. Safety And Efficacy Of  
These Formulas Are The  
Responsibility Of The Compounding  
Pharmacist