

Formula Name DICLOFENAC SOD/ GABAPENTIN / LIDOCAINE HCL ANHYDROUS GEL 5/6/5% GEL

Initials	AP	Quantity	60	MLS	Schedule	0	Days to Exp	180	Date of Formula	7/30/2025
Cost	\$14.44	Cost Per Unit	\$0.24	AWP	\$0.00	AWP Per Unit	\$0.00	Last Review	1/6/2016	

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
VERSAPRO (ANHYDROUS GEL BASE)	GEL	60.0000	GMS	☒	\$11.04		NONE
DICLOFENAC SODIUM	PWD	3.0000	GMS	☐	\$2.16		SOLUBILITY: WATER (SS) ALCOHOL (S) WHITE TO SLIGHTLY YELLOWISH HYGROSCOPIC POWDER PROTECT FROM LIGHT NONSTEROIDAL ANTI- INFLAMMATORY POSSIBLY EFFECTIVE IN OSTEOPOROSIS (PREVENTS BONE RESORPTION IN
GABAPENTIN USP	PWD	3.6000	GMS	☐	\$0.00		
LIDOCAINE HCL USP	PWD	3.0000	GMS	☐	\$0.84		SOLUBILITY: WATER(VS), ALCOHOL(VS)
PROPYLENE GLYCOL USP	LIQU	4.0000	MLS	☐	\$0.11		SOLUBILITY:N/A
STERILE WATER FOR IRRIGATION	LIQU	5.0000	MLS	☐	\$0.28		1000ML PLASTIC BOTTLE

Therapy**Description**

Off-white/opaque colored cream, uniform consistency, no odor

Classification(s) Nonaqueous**Disease State(s)****Instructions**

A. Calculations/Special Considerations:

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log

2. Measure out ingredients

3. Triturate Diclofenac in mortar

4. Wet step 3 with Propylene Glycol

5. Triturate Gabapentin and Lidocaine HCl in mortar

6. Wet step 5 with Sterile Water for irrigation

7. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour active mixtures into syringe, add plunger, and expel air

8. In new syringe fill with Anhydrous base add plunger loosely, invert and tap Anhydrous base to the bottom, once tip is clear of Anhydrous base, fully insert plunger and expel extra air.

9. QS with anhydrous base so the active syringe and base syringe equal the final volume of the compound

10. Attach syringe from step 7 to Anhydrous base syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (complete 40 transfers)

11. Package and label with barcode label and affix "external use only" label

12. Tape labels

13. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxillary sticker

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Add prepack label if compound lot made to stock for multiple Rx's
Add Rx auxiliary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Off-white/opaque colored cream, uniform consistency, no odor

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Formulation - PCCA 13204

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3

INGREDIENT WEIGHT #4

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist