

Formula Name FIRST OMEPRAZOLE SUSPENSION INACTIVE 2MG/ML SOLN

Initials **Quantity** 150 **MLS** **Schedule** 0 **Days to Exp** 30 **Date of Formula** 2/1/2020
Cost \$26.43 **Cost Per Unit** \$0.18 **AWP** **AWP Per Unit** \$0.48 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
OMEPRAZOLE POWDER	PWD	0.3000	GMS	☐	\$26.43		
FIRST-PPI Suspension	SOLN	150.0000	MLS	☐	\$0.00		

Therapy**Description**

Light pink uniform suspension with fine suspended particles

Classification(s) **Disease State(s)** **Instructions**

Equipment Needed:
None

Labelling & Storage:

Label with patient specific prescription label
Shake well and refrigerate sticker
expiration date sticker (30 days)

Special Considerations:
None

1. Hold the neck of the bottle containing the powder and tap the bottom edges on a hard surface to loosen the powder.
2. Remove the cap from the bottle.
3. Tap the top of the induction seal liner to loosen any powder which may have adhered to the liner.
4. Carefully and slowly peel back the inner foil seal liner from the bottle.
5. Using the enclosed tool, scrape the powder from the seal into the bottle. Now using the smaller end of the tool, with firm strength loosen any powder from the inside lower edges of the bottle.
6. Shake the suspension bottle for a few seconds.
6. Open the diluent bottle and transfer about half the contents into the powder bottle.
7. Replace the cap and shake the powder bottle vigorously for approximately 60 seconds.
8. Add the remaining suspension into the powder bottle, allowing to drain for 10 seconds.
9. Replace the cap and shake the bottle vigorously for approximately 30 seconds.
10. Instruct the patient to shake well before each use and to use an oral dosing device that measures the appropriate volume of the oral solution in milliliters.

Formula Source:

CutisPharma
www.cutispharma.com

Testing:

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist