

Formula Name HALOPERIDOL RAPID DISSOLVE TABLET 2MG TAB

Initials **Quantity** 96 EACH **Schedule** 0 **Days to Exp** 180 **Date of Formula** 6/15/2024
Cost \$11.48 **Cost Per Unit** \$0.12 **AWP** **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
HALOPERIDOL USP	PWD	0.1920	GMS	☐	\$1.11		SOLUBILITY: 1:10,000 WATER (VSS), 1:60 ALCOHOL(SP)
MANGO FLAVOR	PWD	0.4000	GMS	☐	\$0.20		
STEVIA 90%	PWD	0.2000	GMS	☐	\$0.12		
BITTERNESS REDUCING AGENT (NF01)	PWD	0.1000	GMS	☐	\$0.11		
Rapid Dissolve Tablet BASE	PWD	71.1080	GMS	☐	\$9.95		

Therapy**Description**

Off-white/cream tablet

Classification(s)**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

RDT - Tray is calibrated for 750mg

B. Equipment Required:

weigh paper/boats, metal spatula, rubber spatula, scale, mortar & pestle, 50-mesh sieve, RDT mold & tamper, oven, labels and unit dose sheets, Inversina tumbler

C. Method of Preparation:

1. Calculate required amounts and fill out compounding logs
2. Measure out ingredients, pre-heat oven to 110° Celsius
3. Do not triturate Medi-RDT base, triturate all other ingredients.
4. Pass all ingredients through 50-mesh sieve
5. Add powder to Inversina tumbler, and mix for 3 minutes, if completing multiple tray batches, separate into batches of 72g/tray
6. Pour half of powder mixture on to mold and spread over cavities
7. Tap cavity to settle mixture, add additional powder to fill cavities
8. Apply the top compression plate to compact the powder, pressing down with moderate pressure
9. Add additional powder, spread over cavities, repeat steps 7,8,9 two additional times.
10. Place mold in oven (at 110° Celsius) and bake for 15 minutes
11. Remove mold from the oven, place wax paper over mold and apply compression plate, put do not apply pressure
12. Turn mold upside down and allow mold to cool for 2 minutes
13. Tap bottom cavity plate with rubber mallet to remove tablets
14. Allow tablets to cool on wax paper for 15-30 minutes
15. Weigh 20% of the tablets and record their weight
16. Have compound certified pharmacist perform final verification
17. Package in amber blister packs, apply adhesive foil tamper labels and place in sleeve.
18. Label blisters with prepack label foil sticker

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxiliary label if compound lot made to order for a unique Rx

_____(place label here)_____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:
Off-white/cream tablet

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Number of tablets made is within 5% of intended batch size

Weigh 20% of batch (weighing each tablet individually) and confirm individual masses are within acceptable range.

Theoretical weight of tablet 0.750g

Acceptable weight range of sampled suppositories 0.675g - 0.825g

Weight #1: _____ Weight #11: _____

Weight #2: _____ Weight #12: _____

Weight #3: _____ Weight #13: _____

Weight #4: _____ Weight #14: _____

Weight #5: _____ Weight #15: _____

Weight #6: _____ Weight #16: _____

Weight #7: _____ Weight #17: _____

Weight #8: _____ Weight #18: _____

Weight #9: _____ Weight #19: _____

Weight #10: _____ Weight #20: _____

Pharmacist signature: _____

G. Packaging Container:

Package in UV light protective amber unit dose blisters with foil label.

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

Medisca F-005 729v4

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3

INGREDIENT WEIGHT #4

INGREDIENT WEIGHT #5

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist