

Formula Name HALOPERIDOL SOLUTAB 1MG TAB

Initials **Quantity** 50 EACH **Schedule** 0 **Days to Exp** 180 **Date of Formula** 8/14/2024
Cost \$0.47 **Cost Per Unit** \$0.01 **AWP** **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
HALOPERIDOL USP	PWD	0.0550	GMS	☐	\$0.32		SOLUBILITY: 1:10,000 WATER (VSS), 1:60 ALCOHOL(SP)
LACTOSE MONOHYDRATE	PWD	2.8600	GMS	☐	\$0.05		
SUCROSE POWDERED	PWD	0.7150	GMS	☐	\$0.04		NONE
TABLET TRITURATE EXCIPIENT	SOLN	1.2000	MLS	☐	\$0.06		

Therapy**Description**

Off-white/cream round tablet

Classification(s)**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

Tablet Triturate Plate is 60mg size (0.053mL volume)

Plate is calibrated to 0.066g blank weight per tablet

Formula has 10% added weight to account for powder loss; total dry batch weight of 3.63g

B. Equipment Required:

weigh paper/boats, metal spatula, rubber spatula, scale, mortar & pestle, tablet triturate plate, baking sheet, labels and unit dose sheets

C. Method of Preparation:

1. Calculate required amounts and fill out compounding logs

2. Measure out ingredients

3. Using geometric dilution, mix active drug, sucrose, and lactose together with trituration in a mortar and pestle

4. Moisten powder mixture with the tablet triturate excipient with 50% of excipient volume.

5. Mix well with rubber spatula using firm, pressing motion and mix until uniform

6. Mix the remaining excipient until the base holds together without crumbling and adheres to the spatula with a playdough-like consistency

7. Place tablet triturate plate on a glass pill tile and begin filling the die cavities

8. When mold appears full, slide the mold towards you and lift off the pill tile. Turn over plate to fill any remaining cavities not completely full.

9. Immediately press the mold over the pegs of the mold base with firm, even pressure to eject the tablets.

****Do not allow tablets to dry in the mold****

10. Eject tablets onto a paper lined baking sheet to allow to air dry for 12 hours. Cover with aluminum foil to protect from light

11. Weigh 10% of the tablets and record their weight

12. Have compound certified pharmacist perform final verification

13. Package in prescription vial or in Unit dose

14. Label blisters with prepack label foil sticker

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxiliary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Off-white/cream round tablet

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Number of tablets made is within 5% of intended batch size

Weigh 10% of batch (weighing each tablet individually) and confirm individual masses are within acceptable range.

Theoretical weight of tablet 0.066g

Adjusted tablet weight for loss 0.060g

Acceptable weight range of sampled suppositories 0.054g - 0.066g

Weight #1: _____ Weight #11: _____

Weight #2: _____ Weight #12: _____

Weight #3: _____ Weight #13: _____

Weight #4: _____ Weight #14: _____

Weight #5: _____ Weight #15: _____

Weight #6: _____ Weight #16: _____

Weight #7: _____ Weight #17: _____

Weight #8: _____ Weight #18: _____

Weight #9: _____ Weight #19: _____

Weight #10: _____ Weight #20: _____

Pharmacist signature: _____

G. Packaging Container:

Package in UV light protective amber vial or unit dose blisters with foil label

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 180 days

I. Formula Source & BUD Testing:

PCCA 10677

BUD - USP 795

INGREDIENT WEIGHT #1

INGREDIENT WEIGHT #2

INGREDIENT WEIGHT #3

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist