

Formula Name IBUPROFEN SUSP INACTIVE 400MG/5ML SUSP

Initials AP **Quantity** 480 **MLS** **Schedule** 0 **Days to Exp** 14 **Date of Formula** 2/10/2021
Cost \$7.27 **Cost Per Unit** \$0.02 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
DISTILLED WATER	LIQU	100.0000	MLS	☐	\$0.01		NONE
CITRIC ACID ANHYDROUS USP	PWD	0.4800	GMS	☐	\$0.03		SOLUBILITY: 1:0.5 WATER(VS) 1:2 ALCOHOL(FS)
POTASSIUM SORBATE NF	PWD	0.8000	GMS	☐	\$0.41		SOLUBILITY: 1:4.5 WATER(FS), 1:35 ALCOHOL(SP)
XANTHAN GUM NF	PWD	2.4000	GMS	☐	\$0.19		SOLUBILITY:N/A
SORBITOL 70% USP SOLUTION	SOLN	200.0000	MLS	☐	\$1.48		SOLUBILITY:1:0.45 WATER(VS) , ALCOHOL(SL)
IBUPROFEN USP	PWD	38.4000	GMS	☐	\$2.69		SOLUBILITY: WATER(VSS), ALCOHOL(VS)
POLYSORBATE 80 NF	LIQU	2.2000	MLS	☐	\$0.44		SOLUBILITY: WATER(VS), ALCOHOL(S)
GLYCERIN USP (SYNTHETIC)	LIQU	48.0000	MLS	☐	\$0.72		SOLUBILITY:N/A
FLAVOR CHERRY CONCENTRATE	SOLN	16.0000	MLS	☐	\$1.30		SOLUBILITY:N/A
METHOCEL 2% SOL	SOLN	0.0000	MLS	☒	\$0.00		

Therapy**Description**

Thick, opaque, red in color suspension with heavy granulated particulate matter

Classification(s) Non-Preserved Aqueous

Disease State(s)

Instructions

A. Calculations/Special Considerations:
None

B. Equipment Required:
Weigh boats; beakers; scales; stir rod/bar; hot/stir plate; homogenizer; graduated cylinder; glass mortar and pestle; funnel; tape

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. Heat Distilled Water to 122 degrees F
4. Add Citric Acid and Potassium Sorbate to Distilled Water and mix well using homogenizer
5. SLOWLY Add Xanthan Gum to Step 4 and stir until dispersion is complete using homogenizer
6. In a separate beaker, warm the Sorbitol
7. In a mixer blend the Ibuprofen and the Polysorbate 80 and triturate in the Glycerin until a frosting like consistency is achieved
8. Mix Sorbitol and Step 7 in mixer
9. Add Step 8 to the beaker containing step 5 and homogenize
10. Add the Flavor to Step 9 and mix well
11. Bring Step 10 to final volume with Methocel 2% and mix thoroughly until a uniform suspension is obtained
12. Have compound certified pharmacist perform final verification
13. Pour into container
14. Label with RX label and "Shake well" "Refrigerate" auxilliary label
15. Tape labels

D. Labeling:
Label with lot number and expiration barcode and ingredient(s) label.
Store in refrigerator auxillary sticker
Shake Well auxillary sticker

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Add prepack label if compound lot made to stock for multiple Rx's
Add Rx auxiliary label if compound lot made to order for a unique Rx

_____ (place label here) _____

E. Description of Finished Product:

Thick, opaque, red in color suspension with heavy granulated particulate matter

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

Bulk store in UV protective amber bottle

Dispense in UV protective amber bottle with volume graduations

H. Storage Requirements:

Store refrigerated (36° - 46 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

PCCA 8801

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist