

Formula Name ISOPROPYL ALCOHOL ANTISEPTIC TOPICAL SOLUTION INACTIVE 60% SOLN

Initials AP **Quantity** 4000 **MLS** **Schedule** 0 **Days to Exp** 30 **Date of Formula** 2/10/2021
Cost \$3.19 **Cost Per Unit** \$0.00 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
DISTILLED WATER	LIQU	0.0000	MLS	☒	\$0.00		NONE
ISOPROPYL ALCOHOL 70%	LIQU	3428.4000	MLS	☐	\$0.00		NONE
HYDROGEN PEROXIDE TOPICAL SOLUTION 3%	LIQU	166.8000	MLS	☐	\$0.00		
GLYCERIN USP (NATURAL)	LIQU	30.0000	MLS	☐	\$3.19		SOLUBILITY:N/A

Therapy**Description**

Clear solution

Classification(s)Non-Preserved Aqueous**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

None

B. Equipment Required:

graduated cylinders; erlenmeyer flask; stir plate; stir bar, funnel; tape

C. Method of Preparation:

1. Calculate required amounts and fill out compounding log
2. Measure out ingredients
3. In a calibrated erlenmeyer flask equipped with a spin bar, add Isopropyl Alcohol and Hydrogen Peroxide topical solution
4. Transfer glycerin stepwise and quantitatively into the erlenmeyer flask and mix well after each addition
5. Rinse the glycerin container several times with water and add contents to
6. Bring to a final volume with distilled water and mix well
7. Have compound certified pharmacist perform final verification
8. Package and label

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

(place label here)

E. Description of Finished Product:

Clear solution

F. Quality Control Procedures:

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

Store in UV protective amber bottle

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

Maximum BUD - 30 days

I. Formula Source & BUD Testing:
PCCA 13716

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist