

Formula Name KETAMINE GEL INACTIVE 1% GEL

Initials AP **Quantity** 60 **MLS** **Schedule** 3 **Days to Exp** 14 **Date of Formula** 2/2/2021
Cost \$4.20 **Cost Per Unit** \$0.07 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
KETAMINE HCL USP	PWD	0.6900	GMS	☐	\$1.46		SOLUBILITY: WATER (FS) ALCOHOL (S) MAXIMUM SOLUBILITY IS 200MG/ML WHITE CRYSTALLINE POWDER WITH A CHARACTERISTIC COLOR SOLUTIONS MAY BE CLEAR TO SLIGHTLY GREEN USED IN ANESTHESIA MAY CAUSE AMNESIA
DISTILLED WATER	LIQU	3.0000	MLS	☐	\$0.00		NONE
POLOX 20% GEL	GEL	60.0000	GMS	☒	\$2.27		NONE
LIPMAX LECITHIN & ISOPROPYL PALMITATE	SOLN	13.0000	MLS	☐	\$0.48		

Therapy**Description**

Solid light yellow, oily, highly viscous organogel

Classification(s) Non-Preserved Aqueous

Disease State(s)

Instructions**A. Calculations/Special Considerations:**

Ketamine 1mg = Ketamine HCl 1.15 mg

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, rubber spatula.

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment

C. Method of Preparation:

1. Calculate required amounts and fill out compounding and control usage logs
2. Measure out ingredients
3. Triturate Ketamine in mortar.
4. Wet step 3 with distilled water (for 240 mL or less: use syringe to syringe method; for 240+ mL: use mixer on lowest speed)
5. Add Lipmax and mix until evenly dispersed
6. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe
7. In the same syringe QS with polox, add plunger loosely, invert and tap mixture to the bottom, once tip is clear, fully insert plunger and expel extra air
8. Attach a new syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (typically 20 transfers)
9. Package and label with barcode label and affix "external use only" label
10. Tape labels
11. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxillary sticker

Add prepack label if compound lot made to stock for multiple Rx's

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:

Solid light yellow, oily, highly viscous organogel

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at controlled room temperature (68°–77 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

PCCA 3900