

CS215: Data Analysis and Interpretation

www.cse.iitb.ac.in/~sharat

(Many slides obtained from Prof. Ajit and Prof. Sunita and gratefully acknowledged. Fair use of internet resources as available to professors under copyright act)

1

Agenda

- Last time
 - Expectation of a function of a random variable
 - Leibnitz rule
 - Pareto principle
 - Course logistics
- Today
 - Linearity of expectations
 - Joint distributions

2

Expected Value May Be Technically Undefined

$$P(X = x) = k/x^2 \text{ for } x \geq 1, x \in \mathbb{Z}^+$$

$$\sum_{x=1}^{\infty} P(X = x) = \sum_{x=1}^{\infty} k/x^2 = 1 \text{ if } k = 6/\pi^2$$

$$E(X) = \sum_{x=1}^{\infty} x P(X = x) = k \sum_{x=1}^{\infty} 1/x = \infty$$

3

3

Pareto principle


- The Pareto **rule of thumb**
 - 20% of pea pods contained 80% of the peas
 - 1992 [United Nations Development Program](#) Report
 - Richest 20% of the world's population receives 82.7% of the world's income
- A small fraction of causes often produces a large fraction of effects
 - Many things in life are not distributed evenly

4

4

Pareto distribution

- Alpha = 1.16 exhibits the Pareto principle
- The type I Pareto distribution is characterized by a scale parameter x_m and a shape parameter α , which is known as the *tail index*.
- The expectation is technically undefined in some cases



$$f_X(x) = \begin{cases} \frac{\alpha x_m^\alpha}{x^{\alpha+1}} & x \geq x_m, \\ 0 & x < x_m. \end{cases}$$

$$\bar{F}(x) = \Pr(X > x) = \begin{cases} \left(\frac{x_m}{x}\right)^\alpha & x \geq x_m, \\ 1 & x < x_m, \end{cases}$$

$$E(X) = \begin{cases} \infty & \alpha \leq 1, \\ \frac{\alpha x_m}{\alpha - 1} & \alpha > 1. \end{cases}$$

5

5

Linearity of Expectations

- A function f is linear if $f(ax) = af(x)$ and $f(x+y) = f(x) + f(y)$
- If $f(x) = af(x) + b$, then the function is affine, but commonly termed as linear
- Expectation is not a function but an operator — it takes a function as input — and it is linear in the same sense.

$$E(a g(X) + b) = \int_{-\infty}^{+\infty} (a g(x) + b) f_X(x) dx$$

Why is this useful?

$$= \int_{-\infty}^{+\infty} a g(x) f_X(x) dx + \int_{-\infty}^{+\infty} b f_X(x) dx$$

$$= a E(g(X)) + b$$

6

6

Mean squared prediction error

- You want to predict the value of a random variable X
 - Suppose we predict it as c
 - What is the error? Maybe $(X - c)$?
- Define the error to be the mean squared error
- Equality holds exactly when $c = \mu$
 - The expected value is the predictor with least mean squared error.

Recall "sample mean"

$$\begin{aligned}
 E((X - c)^2) &= E((X - \mu + \mu - c)^2) \\
 &= E((X - \mu)^2 + (\mu - c)^2 + 2(X - \mu)(\mu - c)) \\
 &\geq E((X - \mu)^2)
 \end{aligned}$$

7

7

Absolute value

- What minimizes the quantity $J(c)$
 - The median
 - The median need not be unique

$$J(c) = \int_{-\infty}^{+\infty} |x - c| f_X(x) dx$$

8

8

Variance

- Variance tells you how much the values of X deviates from the mean, on the average.

$$\text{Var}(X) = E[(X - \mu)^2] = \int_{-\infty}^{\infty} (x - \mu)^2 f_X(x) dx$$

- The positive square root of the variance is the standard deviation

- Low-variance densities concentrate around a point
- High-variance densities are spread out

$$\text{Var}(X) = E[(X - \mu)^2] = \sum_i (x_i - \mu)^2 P(X = x_i)$$

9

9

Variance may be technically undefined

- For some distributions, the defining integral has no finite value, so the variance is undefined.
 - Example: the Pareto distribution with $\alpha < 2$.
- In some cases, the mean is defined but the variance is not
- In some cases, both are undefined.

10

10

Hand Calculation of Variance

$$\begin{aligned}\text{Var}(X) &= E[(X - \mu)^2] = E[X^2 + \mu^2 - 2X\mu] = E[X^2] + \mu^2 - 2E[X]\mu = E[X^2] + \mu^2 - 2\mu^2 \\ &= E[X^2] - \mu^2 = E[X^2] - (E[X])^2\end{aligned}$$

- The middle step uses $E[X] = \mu$
 - This form — mean of the square minus square of the mean — is usually the easier one to compute with.
- Variance when you throw a fair dice
 - Earlier: expected value was $7/2$
 - Option 1: $[(1 - 7/2)^2 + (2 - 7/2)^2 + \dots + (6 - 7/2)^2] (1/6)$
 - Option 2: $E[X^2] = 1/6(1 + 4 + 9 + 16 + 25 + 36) = 91/6$, $\text{Var} = 91/6 - 49/4$

Recall “sample variance”

11

11

Variance

- Shifting a random variable does not change its variance
- Scaling it scales the variance by the square.

$$\begin{aligned}\text{Var}(aX + b) &= E[(aX + b - E(aX + b))^2] = E[(aX + b - (a\mu + b))^2] \\ &= E[a^2(X - \mu)^2] = a^2E[(X - \mu)^2] = a^2\text{Var}(X)\end{aligned}$$

12

12

Jointly distributed random variables

- In machine learning one often needs to model relationships between two or more random variables.
 - CPI at IITB and the salary offered during placements
 - Hours spent on Instagram each day and hours of sleep at night.
 - Literacy level and crime rate
 - Price of a product, peer feedback and probability that a user clicks “Buy Now”
- If we observe multiple paired instances of X and Y , then some combinations of outcomes are more likely than others
 - Random variables rarely live alone. In real data, quantities interact, move together, or constrain one another.
 - Joint distributions captures in a mathematical way these relationships

13

13

Joint Distribution function

- Recall: A random variable is synonymous with a cumulative distribution function, or simply a distribution function
 - We define the cumulative distribution function (cdf) $F_X(x) = \Pr\{X \leq x\}$
- The relationship between two random variables is defined by the joint cumulative probability distribution function

$$G(x, y) = \Pr\{X \leq x, Y \leq y\}$$
- All probability questions about X and Y can be answered in terms of G , for example

$$F_X(x) = \Pr\{X \leq x\} = \Pr\{X \leq x, Y < \infty\} = G(x, \infty)$$

14

Joint pmfs

For discrete random variables X and Y , the **joint probability mass function** is

$$p_{X,Y}(x_i, y_j) = P(X = x_i, Y = y_j)$$

The **pmf** of X follows by summing out the other

$$\begin{aligned} P\{X = x_i\} &= P\left(\bigcup_j \{X = x_i, Y = y_j\}\right) \\ &= \sum_j P\{X = x_i, Y = y_j\} \\ &= \sum_j p(x_i, y_j) \end{aligned}$$

This is called marginalizing

15

15

Example

- 15% of families are childless, 20% have one child, 35% have two and 30% have three (and that's it)
 - Male and female children are equally likely and independent.
 - Let B (G) denote random variables for boys (girls)
 - What is the joint pmf $p(B, G)$
- No children? $P(B = 0, G = 0) = 0.15$
- One girl, no boy? $P(B = 0, G = 1) = 0.2 \times 0.5 = 0.1$

16

16

Example

- 15% of families are childless, 20% have one child, 35% have two and 30% have three. What is the joint pmf $p(B, G)$
- Two boys and one girl? $P(B = 2, G = 1) = 0.3 \times (1/8) \times 3 = 0.1125$
 - All 8 combinations of three children are equally likely; three of them have two boys and one girl.

$i \backslash j$	0	1	2	3	Row Sum $= P\{B = i\}$
0	.15	.10	.0875	.0375	.3750
1	.10	.175	.1125	0	.3875
2	.0875	.1125	0	0	.2000
3	.0375	0	0	0	.0375
Column Sum $= P\{G = j\}$	.3750	.3875	.2000	.0375	

17