

# ECH Parallel Stratification, $B(x)\hat{x}$

---

Open Additional files:

Get dispersion routines by evaluating Plasma\_Dispersion.np

Get plotting and printing routines by evaluating PlotPack.nb

Set Parameters by opening a Parameter Window

Note: Slab profile models defined in initialization cells at the bottom of this notebook.

Plot Real and Imaginary parts of  $n_z$  from 4th order cold plasma dispersion relation (Fast and Slow roots)

N.B. Here  $n_z$  is perpendicular to B. But in nzColdDisFS  $n_z$  is parallel to B. So think of it as exchanging  $n_z \longleftrightarrow n_x$ .

$\alpha = 1.5$ ,  $n_x = 0.25$


```

In[834]:= nParallelFS[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzColdDisFS[freq, ne, b, nz, etaList]];

ntFS = Table[Flatten[{x, nParallelFS[x]}], {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ ]];

nF = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[3]]}];
g1 = PPComplexListPlot[nF, "x (m)", "nx"];
g2 = PPComplexListPlot[nS, "x (m)", "nx"];
Show[{g1, g2}, PlotRange → {0., 10.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```



dataSet=Parallel Stratification 28HHz

xProfileMin=0.5

xProfileMax=1.5

nXmin= $1.46 \times 10^{-19}$

nXmax= $1.46 \times 10^{-19}$

BXmin=0.5

BXmax=1.5

freq=28000.

nz=0.5

etaList={0., 1., 0., 0., 0.}

xmin=0.5

xmax=1.5

Plot Real and Imaginary parts of  $n_z^2$  from 4nd order  
cold plasma dispersion relation (Fast and Slow roots)


N.B. Here  $n_z$  is perpendicular to B. But in nzColdDisFS  $n_z$  is parallel to B. So think of it as exchanging  $n_z \leftrightarrow n_x$ .

$\alpha = 1.5$ ,  $n_x = 0$ .

```
In[842]:= nParallel2FS[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzSqColdDisFS[freq, ne, b, nz, etaList]];

ntFS = Table[Flatten[{x, nParallel2FS[x]}], {x, xmin, xmax,
  (xmax - xmin)/(nPoints - 1)}];

nF = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[3]]}];
g1 = PPComplexListPlot[nF, "x (m)", "nx"];
g2 = PPComplexListPlot[nS, "x (m)", "nx"];
Show[{g1, g2}, PlotRange -> {0., 10.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];
```



dataSet=Parallel Stratification 28HHz

xProfileMin=0.5

xProfileMax=1.5

nXmin= $1.46 \times 10^{19}$

nXmax= $1.46 \times 10^{19}$

BXmin=0.5

BXmax=1.5

freq=28000.

nz=0.5

etaList={0., 1., 0., 0., 0.}

xmin=0.5

xmax=1.5

Plot Real and Imaginary parts of  $n_z$  from 4nd order

cold plasma dispersion relation (Plus and Minus roots)

N.B. Here  $n_z$  is perpendicular to B. But in nzColdDisPM  $n_z$  is parallel to B. So think of it as exchanging  $n_z \longleftrightarrow n_x$ .

$$\alpha = 1.5, n_x = 0.$$


```

In[858]:= nParallelPM[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzColdDisPM[freq, ne, b, nz, etaList]];

ntFS = Table[Flatten[{x, nParallelPM[x]}], {x, xmin, xmax,  $\frac{x_{\text{max}} - x_{\text{min}}}{n_{\text{Points}} - 1}$ }}];

nF = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[3]]}];
g1 = PPComplexListPlot[nF, "x (m)", "nx"];
g2 = PPComplexListPlot[nS, "x (m)", "nx"];
Show[{g1, g2}, PlotRange → {0., 10.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```



dataSet=Parallel Stratification 28HHz

xProfileMin=0.5

xProfileMax=1.5

nXmin= $1.46 \times 10^{-19}$

nXmax= $1.46 \times 10^{-19}$

BXmin=0.5

BXmax=1.5

freq=28000.

nz=0.5

etaList={0., 1., 0., 0., 0.}

xmin=0.5

xmax=1.5

Plot Real and Imaginary parts of  $n_z^2$  from 4nd order cold plasma dispersion relation (Plus and Minus roots)


N.B. Here  $n_z$  is perpendicular to B. But in nzColdDisPM  $n_z$  is parallel to B. So think of it as exchanging  $n_z \leftrightarrow n_x$ .

$\alpha = 1.5$ ,  $n_x = 0$ .

```
In[866]:= nParallel2PM[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzSqColdDisPM[freq, ne, b, nz, etaList]];

ntFS = Table[Flatten[{x, nParallel2FS[x]}], {x, xmin, xmax,
   $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ }}];

nF = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[ntFS][[3]]}];
g1 = PPComplexListPlot[nF, "x (m)", "nx"];
g2 = PPComplexListPlot[nS, "x (m)", "nx"];
Show[{g1, g2}, PlotRange -> {0., 10.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];
```



dataSet=Parallel Stratification 28HHz

xProfileMin=0.5

xProfileMax=1.5

nXmin= $1.46 \times 10^{19}$

nXmax= $1.46 \times 10^{19}$

BXmin=0.5

BXmax=1.5

freq=28 000.

nz=0.5

etaList={0., 1., 0., 0., 0.}

xmin=0.5

xmax=1.5

Plot Real and Imaginary parts of  $n_z$  from 4nd order  
cold plasma dispersion relation (Fast and Slow roots)

$\alpha = 0.1$ ,  $n_x = 0.5$ .


```

In[882]:= nParallel2FS[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzSqColdDisFS[freq, ne, b, nz, etaList]];

nt2FS = Table[Flatten[{x, nParallel2FS[x]}], {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ ]];

nF = Transpose[{Transpose[nt2FS][[1]], Transpose[nt2FS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[nt2FS][[3]]}];
g1 = PPComplexListPlot[nF, "x (m)", "nz2"];
g2 = PPComplexListPlot[nS, "x (m)", "nz2"];
Show[{g1, g2}, PlotRange → {-2., 2.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```



```
dataSet=Parallel Stratification 28HHz
```

```
xProfileMin=0.5
```

```
xProfileMax=1.5
```

```
nXmin= $1. \times 10^{18}$ 
```

```
nXmax= $1. \times 10^{18}$ 
```

```
BXmin=0.5
```

```
BXmax=1.5
```

```
freq=28000.
```

```
nz=0.5
```

```
etaList={0., 1., 0., 0., 0.}
```

```
xmin=0.5
```

```
xmax=1.5
```



Plot Real and Imaginary parts of  $n_z$  from 4nd order  
cold plasma dispersion relation (Fast and Slow roots)

$$\alpha = 0.1, \quad n_x = 0.5$$


```

In[890]:= nParallelFS[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzColdDisFS[freq, ne, b, nz, etaList]];

nt2FS = Table[Flatten[{x, nParallelFS[x]}], {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ ]];

nF = Transpose[{Transpose[nt2FS][[1]], Transpose[nt2FS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[nt2FS][[3]]}];
g1 = PPComplexListPlot[nF, "x (m)", "nz"];
g2 = PPComplexListPlot[nS, "x (m)", "nz"];
Show[{g1, g2}, PlotRange → {-2., 2.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```



```

dataSet=Parallel Stratification 28HHz
xProfileMin=0.5
xProfileMax=1.5
nXmin=1. × 1018
nXmax=1. × 1018
BXmin=0.5
BXmax=1.5
freq=28 000.
nz=0.5
etaList={0., 1., 0., 0., 0.}
xmin=0.5
xmax=1.5

```

Plot Real and Imaginary parts of  $n_z^2$  from 4th order  
cold plasma dispersion relation (Plus and Minus roots)

$$\alpha = 0.1, \quad n_x = 0.5$$


```

In[898]:= nPerp2PM[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  nzSqColdDisPM[freq, ne, b, nz, etaList]]

nt2PM = Table[Flatten[{x, nPerp2PM[x]}], {x, xmin, xmax,  $\frac{x_{\text{max}} - x_{\text{min}}}{n_{\text{Points}} - 1}$ }]];

nP = Transpose[{Transpose[nt2PM][[1]], Transpose[nt2PM][[2]]}];
nM = Transpose[{Transpose[nt2PM][[1]], Transpose[nt2PM][[3]]}];
g1 = PPComplexListPlot[nP, "x (m)", "nx2"];
g2 = PPComplexListPlot[nM, "x (m)", "nx2"];
Show[{g1, g2}, PlotRange → {-2., 2.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```




```

dataSet=Parallel Stratification 28HHz
xProfileMin=0.5
xProfileMax=1.5
nXmin=1. × 1018
nXmax=1. × 1018
BXmin=0.5
BXmax=1.5
freq=28000.
nz=0.5
etaList={0., 1., 0., 0., 0.}
xmin=0.5
xmax=0.5


```

## Plot Profiles


```
In[515]:= Plot[nprof[x], {x, xmin, xmax}]
```



```
In[516]:= Plot[bprof[x], {x, xmin, xmax}]
```



```
In[517]:= Plot[tprof[x], {x, xmin, xmax}]
```



```
In[518]:=  $\alpha$ Hcut[BXmax, freq, nz, 1]
```

Out[518]= 1.00015

## Initialization

### Magnetic field,Density and Temperature Profiles

```
In[519]:= bprof[x_] := If[Abs[(BXmax - BXmin) / BXmax] > 10-6,
    BXmin + (x - xProfileMin) / (xProfileMax - xProfileMin) (BXmax - BXmin), BXmin];
```

```
In[520]:= nprof[x_] := If[Abs[(nXmax - nXmin) / nXmax] > 10-6,
    nXmin + (x - xProfileMin) / (xProfileMax - xProfileMin) (nXmax - nXmin), nXmin];
```

```
In[521]:= tprof[x_] := If[Abs[(TXmax - TXmin) / TXmax] > 10-6,
    TXmin + (x - xProfileMin) / (xProfileMax - xProfileMin) (TXmax - TXmin), TXmin];
```

```
In[522]:=  $\alpha$ Hcut[B_, freq_, nz_, sgn_] := (1 - nz2) × (1 + sgn * 2.79926 B / freq)
```