

# DIII-D ECH with linear Slab Profiles

---

## Open Additional files:

Get dispersion routines by evaluating Disper.nb

Get plotting and printing routines by evaluating PlotPack.nb

Set Parameters by opening a Parameter Window


Note: Slab profile models defined in initialization cells at the bottom of this notebook.

---

## First Do Cold Plasma

Plot Real and Imaginary parts of  $n_x$  from 2nd order warm plasma dispersion relation (i.e.  $E_{||} \equiv 0$ )

```
nPerpCold[x_] := Module[{ne, b, x0}, x0 = x;  
  ne = nprof[x0];  
  b = bprof[x0];  
  ColdDis0[freq, ne, b, nz, etaList]]  
nt = Table[{x, nPerpCold[x]}, {x, xmin, xmax,  $\frac{x_{\text{max}} - x_{\text{min}}}{n_{\text{Points}} - 1}$ }}];  
ComplexListPlot[nt, "x (m)", "nx"]  
paramPrint[{dataSet, xProfileMin, xProfileMax,  
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];
```



```
dataSet=DIII-D slab
```

```
xProfileMin=-0.68
```

```
xProfileMax=0.68
```

```
nXmin= $2.5 \times 10^{19}$ 
```

```
nXmax= $3.5 \times 10^{19}$ 
```

```
BXmin=1.9
```

```
BXmax=2.1
```

```
freq=55990
```

```
nz=0.1
```

```
etaList={1., 0., 0., 0., 0.}
```

```
xmin=-0.68
```


```
xmax=0.68
```

## Plot Real and Imaginary parts of $kx$ from 2nd order cold plasma dispersion relation (i.e. $E_{\parallel} \equiv 0$ )

```

nPerpCold[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  ColdDis0[freq, ne, b, nz, etaList]]
nt = Table[{x, k0 nPerpCold[x]}, {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ }}];
ComplexListPlot[nt, "x (m)", "kx (m-1)"]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```



```

dataSet=DIII-D slab
xProfileMin=-0.68
xProfileMax=0.68
nXmin= $2.5 \times 10^{19}$ 
nXmax= $3.5 \times 10^{19}$ 
BXmin=1.9
BXmax=2.1
freq=56990
nz=0.1
etaList={1., 0., 0., 0., 0.}
xmin=-0.68
xmax=0.68


```

## Plot Real and Imaginary parts of nx from 4nd order cold plasma dispersion relation (i.e. fast and slow)

```

nPerp2FS[x_] := Module[{ne, b, x0}, x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  ColdDis2FS[freq, ne, b, nz, etaList]]
nt2FS = Table[Flatten[{x, nPerp2FS[x]}], {x, xmin, xmax,  $\frac{x_{\text{max}} - x_{\text{min}}}{n_{\text{Points}} - 1}$ }}];
nF = Transpose[{Transpose[nt2FS][[1]], Transpose[nt2FS][[2]]}];
nS = Transpose[{Transpose[nt2FS][[1]], Transpose[nt2FS][[3]]}];
g1 = ComplexListPlot[nF, "x (m)", "nx"];
g2 = ComplexListPlot[nS, "x (m)", "nx"];
Show[{g1, g2}, PlotRange → {-2., 2.}]
paramPrint[{dataSet, xProfileMin, xProfileMax,
  nXmin, nXmax, BXmin, BXmax, freq, nz, etaList, xmin, xmax}];

```



```

dataSet=DIII-D slab
xProfileMin=-0.68
xProfileMax=0.68
nXmin= $2.5 \times 10^{19}$ 
nXmax= $3.5 \times 10^{19}$ 
BXmin=1.9
BXmax=2.1
freq=55990
nz=0.1
etaList={1., 0., 0., 0., 0.}
xmin=-0.68
xmax=0.68

```

## Now Warm Plasma Stuff

Plot Real and Imaginary parts of  $n_x$  from 6th order warm plasma dispersion relation (expanded to 2nd order in  $k_{\perp} \rho$ )

```

nPerpWarm6[x_] := Module[{ne, te, b, x0, TL},
  x0 = x;
  ne = nprof[x0];
  b = bprof[x0];
  TL = tprof[x0] * TLlist;
  WarmDis6[freq, ne, b, nz, etaList, TL]]

nxwarm = Table[Flatten[{x, nPerpWarm6[x]}], {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ }}];
roots = rootSort[nxwarm];
rootsRe = Table[Flatten[{roots[[i]][[1]],
  Table[Re[roots[[i]][[j]]], {j, 2, Length[roots[[i]]}]}], {i, Length[roots]}}];

rootsIm = Table[Flatten[{roots[[i]][[1]],
  Table[Im[roots[[i]][[j]]], {j, 2, Length[roots[[i]]}]}], {i, Length[roots]}}];

g6 = ComplexVectorListPlot[roots, "x (m)", "nx"];
paramPrint[{dataSet, xProfileMin, xProfileMax, nXmin, nXmax,
  BXmin, BXmax, freq, nz, etaList, TLlist, modelList, xmin, xmax}];
Show[g6, PlotRange → All]
Show[g6, PlotRange → {-1.5, 1.5}]
ComplexVectorListPlot[rootsRe, "x", "Re[kx]", PlotRange → {-2., 2.}]
ComplexVectorListPlot[rootsIm, "x", "Re[kx]"]




dataSet=DIII-D slab
xProfileMin=-0.68
xProfileMax=0.68
nXmin= $2.5 \times 10^{19}$ 
nXmax= $3.5 \times 10^{19}$ 
BXmin=1.9
BXmax=2.1
freq=55 990
nz=0.1
etaList={1., 0., 0., 0., 0.}
TLlist={1., 1., 0., 0., 0., 0.}


```

```
modelList={1, 1, 0, 0, 0, 0}
```

```
xmin=-0.68
```

```
xmax=0.68
```





## Now Try with Hot Plasma Dispersion using root finder

```
nPerpHot[x_, nxGuess_] := Module[{ne, b, x0, TL, nx0},
  x0 = x;
  nx0 = nxGuess;
  ne = nprof[x0];
  b = bprof[x0];
  TL = tprof[x0] * TLlist;
  rootRule = FindRoot[DisFuncGeneral[freq, ne, b, nz, nx, etaList, TL,
    nminList, nmaxList, modelList], {nx, nx0}, MaxIterations -> 30];
  nx /. rootRule]
```


First try root finding on warm plasma dispersion rel (model = 1)

```
modelList = Table[1, {i, 1, 6}];
```


```

nxhot[iRoot_] := Module[{iRoot0, nxWarm, rootsWarm, nxH, x0, ne, b, t, TL},
  nxWarm = Table[Flatten[{x, nPerpWarm6[x]}], {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ }}];
  iRoot0 = iRoot;
  rootsWarm = rootSort[nxWarm];
  nxH = Table[0., {i, 1, nPoints}];
  Do[
    ( $x0 = xmin + (i - 1) \frac{x_{\max} - x_{\min}}{nPoints - 1}$ ;
    nxGuess = rootsWarm[[i]][[iRoot0 + 1]];
    (*Print["x0 = ", x0, " nxGuess= ", nxGuess];*)
    nxH[[i]] = {x0, nPerpHot[x0, nxGuess]};), {i, 1, nPoints}];
  nxH];
g7 = ComplexVectorListPlot[nxhot[1], "x (m)", "nx"];
g8 = ComplexVectorListPlot[nxhot[2], "x (m)", "nx"];
g9 = ComplexVectorListPlot[nxhot[3], "x (m)", "nx"];
g10 = ComplexVectorListPlot[nxhot[4], "x (m)", "nx"];
g11 = ComplexVectorListPlot[nxhot[5], "x (m)", "nx"];
g12 = ComplexVectorListPlot[nxhot[6], "x (m)", "nx"];
Show[{g7, g8, g10, g11}, PlotRange -> All]
Show[{g7, g8, g10, g11}, PlotRange -> {- .1, .1}]
paramPrint[{dataSet, ne0, nsep, B, freq, nz, etaList,
  TList, modelList, rmaj, rmin, rsep, sol, alphan, alphaT}];

```







```

dataSet=DIII-D slab
ne0=ne0
nsep=nsep
B=B
freq=55990
nz=0.1
etaList={1., 0., 0., 0., 0.}
TList={1., 1., 0., 0., 0., 0.}
modelList={1, 1, 1, 1, 1, 1}
rmaj=rmaj
rmin=rmin
rsep=rsep
sol=sol
alphan=alphan
alphaT=alphaT

```

Now try with hot plasma (model = 2) for all species. Can I find the warm plasma roots with the full dispersion relation?

```

xPoint = 0.;
guesses = nPerpWarm6[xPoint] (* Warm plasma roots at x=0 *)
{0.472427 + 0.0323801 i, 1.10079 + 0.00415331 i, 22.5351 + 0.667857 i,
 -0.472427 - 0.0323801 i, -1.10079 - 0.00415331 i, -22.5351 - 0.667857 i}

■ Change model to 2
modelList = Table[2, {i, 1, 6}];

```

```

{nPerpHot[xPoint, guesses[[1]]],
 nPerpHot[xPoint, guesses[[2]]],
 nPerpHot[xPoint, guesses[[3]]],
 nPerpHot[xPoint, guesses[[4]]],
 nPerpHot[xPoint, guesses[[5]]],
 nPerpHot[xPoint, guesses[[6]]]}
paramPrint[{dataSet, ne0, nsep, B, freq, nz,
  etaList, TList, modelList, rmaj, rmin, rsep, sol, alphan, alphaT}];
{0.590246 + 1.7869 × 10-27 i, 1.18681 + 3.41307 × 10-28 i, 1.18681 + 7.67968 × 10-26 i,
 -0.590246 - 1.7869 × 10-27 i, -1.18681 - 3.41307 × 10-28 i, -1.18681 - 7.67968 × 10-26 i}

dataSet=DIII-D slab
ne0=ne0
nsep=nsep
B=B
freq=55 990
nz=0.1
etaList={1., 0., 0., 0., 0.}
TList={1., 1., 0., 0., 0., 0.}
modelList={2, 2, 2, 2, 2, 2}
rmaj=rmaj
rmin=rmin
rsep=rsep
sol=sol
alphan=alphan
alphaT=alphaT
■ Hot plasma root finder gets all roots. What about at xmin?

xPoint = xmin;
guesses = nPerpWarm6[xPoint] (* Warm plasma roots at x=0 *)
{0.590188, 1.18597, 19.3865, -0.590188, -1.18597, -19.3865}

```

```

{nPerpHot[xPoint, guesses[[1]]],
 nPerpHot[xPoint, guesses[[2]]],
 nPerpHot[xPoint, guesses[[3]]],
 nPerpHot[xPoint, guesses[[4]]],
 nPerpHot[xPoint, guesses[[5]]],
 nPerpHot[xPoint, guesses[[6]]]}
paramPrint[{dataSet, ne0, nsep, B, freq, nz,
  etaList, TList, modellList, rmaj, rmin, rsep, sol, alphan, alphaT}];
{0.590246 + 1.7869 × 10-27 i, 1.18681 + 3.41307 × 10-28 i, 1.18681 + 7.67968 × 10-26 i,
 -0.590246 - 1.7869 × 10-27 i, -1.18681 - 3.41307 × 10-28 i, -1.18681 - 7.67968 × 10-26 i}

dataSet=DIII-D slab
ne0=ne0
nsep=nsep
B=B
freq=55 990
nz=0.1
etaList={1., 0., 0., 0., 0.}
TList={1., 1., 0., 0., 0., 0.}
modellList={2, 2, 0, 0, 0, 0}
rmaj=rmaj
rmin=rmin
rsep=rsep
sol=sol
alphan=alphan
alphaT=alphaT

```


- Hot plasma root finder gets 2 of the roots

## Plot hot plasma vs x

```

g7 = ComplexVectorListPlot[nxhot[1], "x (m)", "nx"];
g8 = ComplexVectorListPlot[nxhot[2], "x (m)", "nx"];
g9 = ComplexVectorListPlot[nxhot[3], "x (m)", "nx"];
g10 = ComplexVectorListPlot[nxhot[4], "x (m)", "nx"];
g11 = ComplexVectorListPlot[nxhot[5], "x (m)", "nx"];
g12 = ComplexVectorListPlot[nxhot[6], "x (m)", "nx"];
Show[{g7, g8, g9, g10, g11, g12}, PlotRange → All, AxesOrigin → {0., 0.}]
Show[{g7, g8, g9, g10, g11, g12}, PlotRange → {-2., 2.}, AxesOrigin → {0., 0.}]
paramPrint[{dataSet, ne0, nsep, B, freq, nz, etaList,
  TList, modellList, rmaj, rmin, rsep, sol, alphan, alphaT}];

```




```

dataSet=DIII-D slab
ne0=ne0
nsep=nsep
B=B
freq=55990
nz=0.1
etaList={1., 0., 0., 0., 0.}
TList={1., 1., 0., 0., 0., 0.}
modelList={2, 2, 0, 0, 0, 0}
rmaj=rmaj
rmin=rmin
rsep=rsep
sol=sol
alphan=alphan
alphaT=alphaT

```

```
Show[{g7, g10}, PlotRange → All, AxesOrigin → {0., 0.}]
Show[{g8, g11}, PlotRange → All, AxesOrigin → {0., 0.}]
Show[{g9, g12}, PlotRange → All, AxesOrigin → {0., 0.}]
```



## Focus on the damping


```

nxhotIm[iRoot_] := Module[{iRoot0, nxWarm, rootsWarm, nxH, x0, ne, b, t, TL},
  nxWarm = Table[Flatten[{x, nPerpWarm6[x]}], {x, xmin, xmax,  $\frac{x_{\max} - x_{\min}}{nPoints - 1}$ ]];
  iRoot0 = iRoot;
  rootsWarm = rootSort[nxWarm];
  nxH = Table[0., {i, 1, nPoints}];
  Do[
    ( $x0 = xmin + (i - 1) \frac{x_{\max} - x_{\min}}{nPoints - 1}$ ;
    nxGuess = rootsWarm[[i]][[iRoot0 + 1]];
    (*Print["x0 = ", x0," nxGuess= ",nxGuess];*)
    nxH[[i]] = {x0, Im[nPerpHot[x0, nxGuess]]};), {i, 1, nPoints}];
  nxH];

```


### X - mode

```
ListPlot[nxhotIm[1], PlotRange -> All]
```



## O - mode


```
ListPlot[nxhotIm[2], PlotRange -> All]
```




---

## Plot Profiles


```
Plot[nprof[x], {x, xmin, xmax}]
```



Plot[bprof[x], {x, xmin, xmax}]



Plot[tprof[x], {x, xmin, xmax}]



## Initialization

### Magnetic field, Density and Temperature Profiles

```
bprof[x_] := If[Abs[(BXmax - BXmin) / BXmax] > 10-6,
  BXmin + (x - xProfileMin) / (xProfileMax - xProfileMin) (BXmax - BXmin), BXmin];
```

```
nprof[x_] := If[Abs[(nXmax - nXmin) / nXmax] > 10-6,
  nXmin + (x - xProfileMin) / (xProfileMax - xProfileMin) (nXmax - nXmin), nXmin];
```

```
tprof[x_] := If[Abs[(TXmax - TXmin) / TXmax] > 10-6,
  TXmin + (x - xProfileMin) / (xProfileMax - xProfileMin) (TXmax - TXmin), TXmin];
```