
Sapphire 60445 bacterial contaminant

Peter M. Letcher
The University of Alabama
14 Sept 2015

Summary:


- Samples received from Days 1, 3, and 4 of infection
- Day 1: abundance of healthy algal cells, no sign of infection
- Day 3: Abundant infection, bacteria both inside and outside infected cells; bacterial mostly rod-shaped, some hook-shaped, division by binary fission
- Day 4: host cells fully degraded, often cell wall broken down; a protist component also present, but in low numbers.

Day 1: healthy algal cells: notice chloroplasts and nucleus





Day 3: infected algal cells: bacterial cysts inside infected cells





Day 3: infected algal cells: cysts inside, and rodshaped bacteria outside






Day 3: infected algal cells: the "fish hook"; I think it's the same as the rod-shaped.






Day 4: algal remnants: many algal cells empty; in many cases cell wall degraded





Day 4: Protist flagellated cells also present (notice tubular cristae in mitochondrion)




Sapphire 60445 bacterial contaminant, part 2

Peter M. Letcher
The University of Alabama
150918

Additional images

- Slides 3 and 4 are long bacteria
- Slide 5 are images of binary fission
- Slide 6 is the crescent-shaped bacterium with an appendage
- Slide 7 may be a bacterium crossing the host cell wall and plasma membrane. We may get better images of this at Day 2


Long bacterium external to (L) and inside (R) host cell







~5 μm length

Serial sections of same bacterium inside host, so I think what we are seeing is a long bacterium coiled inside





Cell division via binary fission



Hook-shaped or crescent-shaped bacterium (and look-alike Caulobacter)





https://microbewiki.kenyon.edu/index.php/Caulobacter_crescentus

Maybe the bacterium crossing the host cell wall (arrow)

