

de Rham Cohomology

and a crash course on differential geometry

Noah Chrein

Department of Mathematics
University of Rochester

November 30, 2016

- Introduction to Smooth Manifolds, John M. Lee
- "de Rham Theorem", Marco Perez
(<http://www1.mat.uniroma1.it/people/piazza/deRham-thm.pdf>)

de Rham Cohomology

and a crash course on differential geometry

- 1 Smooth Manifolds
 - Smooth Manifolds
 - Visualization
 - Smooth Maps
- 2 Tangent space
 - Definition
 - Push Forward
 - Tangent Bundle and Sections
 - Cotangent Space
 - Pullback
 - Differential Map
- 3 Tensors
 - Tensors
 - Tensor Product
 - Tensor space and bundle
 - Alternating Tensors
 - Wedge Product
- 4 Differential Forms
 - Differential Forms
 - Exterior Derivative
 - Closed and Exact Forms
- 5 de Rham Cohomology
 - The Chain Complex
 - Invariance of Homology
 - Proof of Homotopy Equivalence
 - Calculating some de Rham groups
 - Mayer-Vietoris Sequence
 - Smooth Singlar Cohomology
 - de Rham Theorem

Smooth Manifolds

Definition

- An n -dimensional manifold M is a "nice" locally euclidean space. That is, there is a collection of open sets $\{U_i\}$ and homeomorphisms $\{\phi_i : U_i \rightarrow \mathbb{R}^n\}$ that covers all of M

Smooth Manifolds

Definition

- An n -dimensional manifold M is a "nice" locally euclidean space. That is, there is a collection of open sets $\{U_i\}$ and homeomorphisms $\{\phi_i : U_i \rightarrow \mathbb{R}^n\}$ that covers all of M
- These local homeomorphisms, called coordinate charts, allow us to talk about "local coordinates" around each point in M


Smooth Manifolds

Definition

- An n -dimensional manifold M is a "nice" locally euclidean space. That is, there is a collection of open sets $\{U_i\}$ and homeomorphisms $\{\phi_i : U_i \rightarrow \mathbb{R}^n\}$ that covers all of M
- These local homeomorphisms, called coordinate charts, allow us to talk about "local coordinates" around each point in M
- A smooth manifold is a manifold whose transition maps, $\phi_a \circ \phi_b^{-1}$, are smooth maps (as defined for maps between \mathbb{R}^n)

Smooth Manifolds

Visualization



Smooth Manifolds

Smooth Maps

- A smooth map is a continuous map $F : M \rightarrow N$ such that the composition $\phi^{-1} \circ F \circ \psi$ is a smooth map (as defined for maps from R^m to R^n)

Smooth Manifolds

Smooth Maps

- A smooth map is a continuous map $F : M \rightarrow N$ such that the composition $\phi^{-1} \circ F \circ \psi$ is a smooth map (as defined for maps from R^m to R^n)
- Smoothness is a local property, that is, smoothness need only apply in some neighborhood around each point to be smooth globally

Smooth Manifolds

Smooth Maps

- A smooth map is a continuous map $F : M \rightarrow N$ such that the composition $\phi^{-1} \circ F \circ \psi$ is a smooth map (as defined for maps from \mathbb{R}^m to \mathbb{R}^n)
- Smoothness is a local property, that is, smoothness need only apply in some neighborhood around each point to be smooth globally
- Define $C^\infty(M, N)$ to be the set of smooth maps from M to N . $C^\infty(M) := C^\infty(M, \mathbb{R})$

Tangent Space

Definition

- let $p \in M$ A derivation at p is a linear map $X : C^\infty(M) \rightarrow R$ such that $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$

Tangent Space

Definition

- let $p \in M$ A derivation at p is a linear map $X : C^\infty(M) \rightarrow R$ such that $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$
- Define the vector space $T_p M$ as the set of derivations at p

Tangent Space

Definition

- let $p \in M$ A derivation at p is a linear map $X : C^\infty(M) \rightarrow \mathbb{R}$ such that $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$
- Define the vector space $T_p M$ as the set of derivations at p
- If $M = \mathbb{R}^n$ and $v \in \mathbb{R}^n$ define $D_v \in T_p \mathbb{R}^n$ by
$$D_v f_p = \frac{\partial}{\partial t} f(p + tv)_{t=0}$$

Tangent Space

Definition

- let $p \in M$ A derivation at p is a linear map $X : C^\infty(M) \rightarrow R$ such that $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$
- Define the vector space $T_p M$ as the set of derivations at p
- If $M = R^n$ and $v \in R^n$ define $D_v \in T_p R^n$ by
$$D_v f_p = \frac{\partial}{\partial t} f(p + tv)_{t=0}$$
- All derivations of R^n are of this form. if $v = v^i e_i$ then
$$D_v f(p) = v^i \frac{\partial f}{\partial x^i}(a),$$
 so the derivations $\frac{\partial}{\partial x^i}$ form a basis for $T_p R^n$

Tangent Space

Definition

- let $p \in M$ A derivation at p is a linear map $X : C^\infty(M) \rightarrow \mathbb{R}$ such that $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$
- Define the vector space $T_p M$ as the set of derivations at p
- If $M = \mathbb{R}^n$ and $v \in \mathbb{R}^n$ define $D_v \in T_p \mathbb{R}^n$ by
$$D_v f_p = \frac{\partial}{\partial t} f(p + tv)_{t=0}$$
- All derivations of \mathbb{R}^n are of this form. if $v = v^i e_i$ then
$$D_v f(p) = v^i \frac{\partial f}{\partial x^i}(a),$$
 so the derivations $\frac{\partial}{\partial x^i}$ form a basis for $T_p \mathbb{R}^n$
- Likewise if $\{x^i\}$ are local coordinates for a point $p \in M$ then the derivations $\frac{\partial}{\partial x^i}$ form a basis for $T_p M$

Tangent Space

Definition

- let $p \in M$ A derivation at p is a linear map $X : C^\infty(M) \rightarrow \mathbb{R}$ such that $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$
- Define the vector space $T_p M$ as the set of derivations at p
- If $M = \mathbb{R}^n$ and $v \in \mathbb{R}^n$ define $D_v \in T_p \mathbb{R}^n$ by
$$D_v f_p = \frac{\partial}{\partial t} f(p + tv)_{t=0}$$
- All derivations of \mathbb{R}^n are of this form. if $v = v^i e_i$ then
$$D_v f(p) = v^i \frac{\partial f}{\partial x^i}(a),$$
 so the derivations $\frac{\partial}{\partial x^i}$ form a basis for $T_p \mathbb{R}^n$
- Likewise if $\{x^i\}$ are local coordinates for a point $p \in M$ then the derivations $\frac{\partial}{\partial x^i}$ form a basis for $T_p M$
- Thus one can think of the tangent space as the directional derivatives on a smooth manifold.

Tangent Space

Push Forward

- For $F : M \rightarrow N$ define the pushforward $F_* : T_p M \rightarrow T_{F(p)} N$ by $(F_* X)(f) = X(f \circ F)$

Tangent Space

Push Forward

- For $F : M \rightarrow N$ define the pushforward $F_* : T_p M \rightarrow T_{F(p)} N$ by $(F_* X)(f) = X(f \circ F)$
- $(G \circ F)_* = G_* \circ F_*$ and $Id_* = Id_{(T_p M)}$, so $T_p : Man^\infty \rightarrow Vect_R$ is a covariant functor.

Tangent Space

Push Forward

- For $F : M \rightarrow N$ define the pushforward $F_* : T_p M \rightarrow T_{F(p)} N$ by $(F_* X)(f) = X(f \circ F)$
- $(G \circ F)_* = G_* \circ F_*$ and $Id_* = Id_{(T_p M)}$, so $T_p : Man^\infty \rightarrow Vect_R$ is a covariant functor.
- If one looks at a coordinate neighborhood of p , one will find that the pushforward is none other than the jacobian of the map $\phi F \phi^{-1}$.

Tangent Space

Tangent Bundle and Sections

- Define the Tangent Bundle $TM = \coprod_{p \in M} T_p M$ and define the projection $\pi : TM \rightarrow M$ by $\pi(p, X) = p$

Tangent Space

Tangent Bundle and Sections

- Define the Tangent Bundle $TM = \coprod_{p \in M} T_p M$ and define the projection $\pi : TM \rightarrow M$ by $\pi(p, X) = p$
- There is a unique smooth structure on TM such that the projection and pushforwards are smooth, making TM into a smooth $2n$ -dimensional manifold

Tangent Space

Tangent Bundle and Sections

- Define the Tangent Bundle $TM = \coprod_{p \in M} T_p M$ and define the projection $\pi : TM \rightarrow M$ by $\pi(p, X) = p$
- There is a unique smooth structure on TM such that the projection and pushforwards are smooth, making TM into a smooth $2n$ -dimensional manifold
- A smooth section, or a smooth vector field, is a map $\sigma : M \rightarrow TM$ such that $\pi \circ \sigma = Id_M$

Tangent Space

Tangent Bundle and Sections

- Define the Tangent Bundle $TM = \coprod_{p \in M} T_p M$ and define the projection $\pi : TM \rightarrow M$ by $\pi(p, X) = p$
- There is a unique smooth structure on TM such that the projection and pushforwards are smooth, making TM into a smooth $2n$ -dimensional manifold
- A smooth section, or a smooth vector field, is a map $\sigma : M \rightarrow TM$ such that $\pi \circ \sigma = Id_M$
- Define $\mathfrak{T}(M)$ the $C^\infty(M)$ - module of sections of TM .
($f \cdot \sigma$ is a section of TM , and $\sigma(p), \tau(p) \in T_p M$ so $\sigma + \tau$ is well defined.)

Tangent Space

Tangent Bundle and Sections

- Define the Tangent Bundle $TM = \coprod_{p \in M} T_p M$ and define the projection $\pi : TM \rightarrow M$ by $\pi(p, X) = p$
- There is a unique smooth structure on TM such that the projection and pushforwards are smooth, making TM into a smooth $2n$ -dimensional manifold
- A smooth section, or a smooth vector field, is a map $\sigma : M \rightarrow TM$ such that $\pi \circ \sigma = Id_M$
- Define $\mathfrak{T}(M)$ the $C^\infty(M)$ - module of sections of TM .
($f \cdot \sigma$ is a section of TM , and $\sigma(p), \tau(p) \in T_p M$ so $\sigma + \tau$ is well defined.)
- We can pushforward sections on M to N by applying F_* , however, these maps aren't always sections on N

- Recall the definition of the dual space of a vector space V^*

Tangent Space

Cotangent Space

- Recall the definition of the dual space of a vector space V^*
- define for each point p in M , the cotangent space
 $T_p^*M = (T_pM)^*$.

Tangent Space

Cotangent Space

- Recall the definition of the dual space of a vector space V^*
- define for each point p in M , the cotangent space $T_p^*M = (T_pM)^*$.
- Analogous to the tangent bundle and the space of sections, define cotangent bundle and it's sections $T^*M = \coprod_{p \in M} T_p^*M$ and $\mathfrak{T}^*(M)$

Tangent Space

Cotangent Space

- Recall the definition of the dual space of a vector space V^*
- define for each point p in M , the cotangent space $T_p^*M = (T_pM)^*$.
- Analogous to the tangent bundle and the space of sections, define cotangent bundle and its sections $T^*M = \coprod_{p \in M} T_p^*M$ and $\mathfrak{T}^*(M)$
- For local coordinates $\{x_p^i\}$, to the basis $\frac{\partial}{\partial x^i}$ of T_pM , there is a dual basis $\{\epsilon^i\}$ for T_p^*M .

Tangent Space

Pullback

- for a smooth map $F : M \rightarrow N$ define the pullback map $F^* : T_{F(p)}^* N \rightarrow T_p^* M$ by $(F^* \epsilon)_p(X) = \epsilon_{F(p)}(F_* X)$

Tangent Space

Pullback

- for a smooth map $F : M \rightarrow N$ define the pullback map $F^* : T_{F(p)}^* N \rightarrow T_p^* M$ by $(F^* \epsilon)_p(X) = \epsilon_{F(p)}(F_* X)$
- Unlike the pushforward, the pullback of a section of N will be a section of M

Tangent Space

Pullback

- for a smooth map $F : M \rightarrow N$ define the pullback map $F^* : T_{F(p)}^* N \rightarrow T_p^* M$ by $(F^* \epsilon)_p(X) = \epsilon_{F(p)}(F_* X)$
- Unlike the pushforward, the pullback of a section of N will be a section of M
- note that $(F \circ G)^* = G^* \circ F^*$ and that $Id^* = Id_{T_p^* M}$ thus $T^* : Man^\infty \rightarrow Vect_R$ is a contravariant functor

Tangent Space

Differential Map

- Define the map $d : C^\infty(M) \rightarrow \mathfrak{T}^*(M)$ by $[df]_p(X) = X_p(f)$

Tangent Space

Differential Map

- Define the map $d : C^\infty(M) \rightarrow \mathcal{T}^*(M)$ by $[df]_p(X) = X_p(f)$
- Locally, we can write the differential of a function in terms of a basis for T_p^*M
$$df_p = \frac{\partial f}{\partial x^i}(p) \epsilon^i$$

Tangent Space

Differential Map

- Define the map $d : C^\infty(M) \rightarrow \mathfrak{T}^*(M)$ by $[df]_p(X) = X_p(f)$
- Locally, we can write the differential of a function in terms of a basis for T_p^*M
$$df_p = \frac{\partial f}{\partial x^i}(p) \epsilon^i$$
- Thus we have $\{dx^i\}$ as a basis for T_p^*M

Tangent Space

Differential Map

- Define the map $d : C^\infty(M) \rightarrow \mathfrak{T}^*(M)$ by $[df]_p(X) = X_p(f)$
- Locally, we can write the differential of a function in terms of a basis for T_p^*M
$$df_p = \frac{\partial f}{\partial x^i}(p)\epsilon^i$$
- Thus we have $\{dx^i\}$ as a basis for T_p^*M
- A generalization of this differential map to tensor spaces is the main object of study for de Rham Cohomology.

Tensors

- A k -covariant Tensor on a vector space V is a linear map
 $T : V^k \rightarrow \mathbb{R}$

Tensors

- A k -covariant Tensor on a vector space V is a linear map
 $T : V^k \rightarrow \mathbb{R}$
- denote the vector space of such tensors $T^k(V)$. Addition and scalar multiplication are defined pointwise.

Tensors

- A k -covariant Tensor on a vector space V is a linear map
 $T : V^k \rightarrow \mathbb{R}$
- denote the vector space of such tensors $T^k(V)$. Addition and scalar multiplication are defined pointwise.
- note that $T^1(V) = V^*$

Tensors

Tensor Product

- if $T \in T^k(V)$, $S \in T^{k'}(V)$ define $T \otimes S \in T^{k+k'}(V)$ by
$$T \otimes S(X_1, \dots, X_{k+k'}) = T(X_1, \dots, X_k) \cdot S(X_{k+1}, \dots, X_{k+k'})$$

Tensors

Tensor Product

- if $T \in T^k(V)$, $S \in T^{k'}(V)$ define $T \otimes S \in T^{k+k'}(V)$ by
$$T \otimes S(X_1, \dots, X_{k+k'}) = T(X_1, \dots, X_k) \cdot S(X_{k+1}, \dots, X_{k+k'})$$
- If $\{e_i\}$ is a basis for V and $\{\epsilon_j\}$ is the dual basis for V^* , then $\epsilon_{j_1} \otimes \dots \otimes \epsilon_{j_k}$ is a basis for $T^k(V)$. Thus
$$T^k(V) \cong \underbrace{V^* \otimes \dots \otimes V^*}_{k\text{-times}}$$

Tensors

Tensor space and bundle

- let M be a smooth manifold, at every point $p \in M$ define
 $T_p^k M = T^k(T_p M)$, and the tensor bundle
 $T^k M = \coprod_{p \in M} T_p^k M$

Tensors

Tensor space and bundle

- let M be a smooth manifold, at every point $p \in M$ define
 $T_p^k M = T^k(T_p M)$, and the tensor bundle
 $T^k M = \coprod_{p \in M} T_p^k M$
- define $\mathfrak{T}^k(M) = \{\text{Sections of } T^k M\}$

Tensors

Tensor space and bundle

- let M be a smooth manifold, at every point $p \in M$ define
 $T_p^k M = T^k(T_p M)$, and the tensor bundle
 $T^k M = \coprod_{p \in M} T_p^k M$
- define $\mathfrak{T}^k(M) = \{\text{Sections of } T^k M\}$
- Note: $T^1 M = T^* M$ and $\mathfrak{T}^1(M) = \mathfrak{T}^*(M)$.
define $T^0(M) = R$

Tensors

Tensor space and bundle

- let M be a smooth manifold, at every point $p \in M$ define
 $T_p^k M = T^k(T_p M)$, and the tensor bundle
 $T^k M = \coprod_{p \in M} T_p^k M$
- define $\mathfrak{T}^k(M) = \{\text{Sections of } T^k M\}$
- Note: $T^1 M = T^* M$ and $\mathfrak{T}^1(M) = \mathfrak{T}^*(M)$.
define $T^0(M) = R$
- for a smooth map $F : M \rightarrow N$ define the pullback map
 $F^* : \mathfrak{T}^k(N) \rightarrow \mathfrak{T}^k(M)$ by
 $(F^* \sigma)_p(X_1, \dots, X_k) = \sigma_{F(p)}(F_* X_1, \dots, F_* X_k)$

Tensors

Alternating Tensors

- The ideas behind a signed k -dimensional volume are captured by Alternating Tensors.

Tensors

Alternating Tensors

- The ideas behind a signed k -dimensional volume are captured by Alternating Tensors.
- An alternating tensor $T \in T^k(V)$ is one that $\forall i \neq j$
$$T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = -T(X_1, \dots, X_j, \dots, X_i, \dots, X_k).$$

Tensors

Alternating Tensors

- The ideas behind a signed k -dimensional volume are captured by Alternating Tensors.
- An alternating tensor $T \in T^k(V)$ is one that $\forall i \neq j$
 $T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = -T(X_1, \dots, X_j, \dots, X_i, \dots, X_k)$.
- The alternating Tensors form a subspace $\Lambda^k(V) \subset T^k(V)$

Tensors

Alternating Tensors

- The ideas behind a signed k -dimensional volume are captured by Alternating Tensors.
- An alternating tensor $T \in T^k(V)$ is one that $\forall i \neq j$
 $T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = -T(X_1, \dots, X_j, \dots, X_i, \dots, X_k)$.
- The alternating Tensors form a subspace $\Lambda^k(V) \subset T^k(V)$
- For the basis $\{\epsilon^i\}$ of V^* , and an ordered k -tuple I , define the elementary alternating k -tensor ϵ^I , by

$$\epsilon^I(X_1, \dots, X_k) = \det \begin{pmatrix} \epsilon^{i_1}(X_1) & \dots & \epsilon^{i_k}(X_1) \\ \vdots & & \vdots \\ \epsilon^{i_1}(X_k) & \dots & \epsilon^{i_k}(X_k) \end{pmatrix}$$

Tensors

Alternating Tensors

- The ideas behind a signed k -dimensional volume are captured by Alternating Tensors.
- An alternating tensor $T \in T^k(V)$ is one that $\forall i \neq j$
 $T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = -T(X_1, \dots, X_j, \dots, X_i, \dots, X_k)$.
- The alternating Tensors form a subspace $\Lambda^k(V) \subset T^k(V)$
- For the basis $\{\epsilon^i\}$ of V^* , and an ordered k -tuple I , define the elementary alternating k -tensor ϵ^I , by

$$\epsilon^I(X_1, \dots, X_k) = \det \begin{pmatrix} \epsilon^{i_1}(X_1) & \dots & \epsilon^{i_k}(X_1) \\ \vdots & & \vdots \\ \epsilon^{i_1}(X_k) & \dots & \epsilon^{i_k}(X_k) \end{pmatrix}$$

- then the set $\{\epsilon^I \mid I \text{ is increasing}\}$ is a basis for $\Lambda^k(V)$

Tensors

Alternating Tensors

- The ideas behind a signed k -dimensional volume are captured by Alternating Tensors.
- An alternating tensor $T \in T^k(V)$ is one that $\forall i \neq j$
 $T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = -T(X_1, \dots, X_j, \dots, X_i, \dots, X_k)$.
- The alternating Tensors form a subspace $\Lambda^k(V) \subset T^k(V)$
- For the basis $\{\epsilon^i\}$ of V^* , and an ordered k -tuple I , define the elementary alternating k -tensor ϵ^I , by

$$\epsilon^I(X_1, \dots, X_k) = \det \begin{pmatrix} \epsilon^{i_1}(X_1) & \dots & \epsilon^{i_k}(X_1) \\ \vdots & & \vdots \\ \epsilon^{i_1}(X_k) & \dots & \epsilon^{i_k}(X_k) \end{pmatrix}$$

- then the set $\{\epsilon^I \mid I \text{ is increasing}\}$ is a basis for $\Lambda^k(V)$
- some simple consequences of this is that $\Lambda^k(V) = 0$ if $k > \dim(V)$ and that $\Lambda^1(V) = T^1(V)$

Tensors

Wedge Product

- define $Alt : T^k(V) \rightarrow \Lambda^k(V)$ by
$$Alt T(X_1, \dots, X_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (sgn \sigma) T(X_{\sigma 1}, \dots, X_{\sigma k})$$

Tensors

Wedge Product

- define $Alt : T^k(V) \rightarrow \Lambda^k(V)$ by
$$Alt T(X_1, \dots, X_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (sgn \sigma) T(X_{\sigma 1}, \dots, X_{\sigma k})$$
- for two alternating tensors $\omega \in \Lambda^k(V), \eta \in \Lambda^l(V)$ define the wedge product $\omega \wedge \eta = \frac{(k+l)!}{k!l!} Alt(\omega \otimes \eta)$

Tensors

Wedge Product

- define $Alt : T^k(V) \rightarrow \Lambda^k(V)$ by
$$Alt T(X_1, \dots, X_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (sgn \sigma) T(X_{\sigma 1}, \dots, X_{\sigma k})$$
- for two alternating tensors $\omega \in \Lambda^k(V), \eta \in \Lambda^l(V)$ define the wedge product $\omega \wedge \eta = \frac{(k+l)!}{k!l!} Alt(\omega \otimes \eta)$
- $\epsilon^I \wedge \epsilon^J = \epsilon^{IJ}$

Tensors

Wedge Product

- define $Alt : T^k(V) \rightarrow \Lambda^k(V)$ by
$$Alt T(X_1, \dots, X_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (sgn \sigma) T(X_{\sigma 1}, \dots, X_{\sigma k})$$
- for two alternating tensors $\omega \in \Lambda^k(V), \eta \in \Lambda^l(V)$ define the wedge product $\omega \wedge \eta = \frac{(k+l)!}{k!l!} Alt(\omega \otimes \eta)$
- $\epsilon^I \wedge \epsilon^J = \epsilon^{IJ}$
- The key properties of the wedge product are bilinearity, associativity, anticommutativity ($\omega \wedge \eta = (-1)^{kl} \eta \wedge \omega$) and the two following formula:
$$\epsilon^{i_1} \wedge \dots \wedge \epsilon^{i_k} = \epsilon^I$$
$$\omega^1 \wedge \dots \wedge \omega^k(X_1, \dots, X_k) = \det \omega^i(X_j)$$

Differential Forms

- Define the space of alternating k -tensors at p ,
 $\Lambda_p^k(M) = \Lambda^k(T_p M)$ and the bundle $\Lambda^k(M) = \coprod_{p \in M} \Lambda_p^k(M)$.

Differential Forms

- Define the space of alternating k -tensors at p ,
 $\Lambda_p^k(M) = \Lambda^k(T_p M)$ and the bundle $\Lambda^k(M) = \coprod_{p \in M} \Lambda_p^k(M)$.
- define the space of " k -forms" as the sections of $\Lambda^k(M)$, that is
 $\Omega^k(M) = \{\omega \text{ section of } \Lambda^k(M)\}$

Differential Forms

- Define the space of alternating k-tensors at p,
 $\Lambda_p^k(M) = \Lambda^k(T_p M)$ and the bundle $\Lambda^k(M) = \coprod_{p \in M} \Lambda_p^k(M)$.
- define the space of "k-forms" as the sections of $\Lambda^k(M)$, that is
 $\Omega^k(M) = \{\omega \text{ section of } \Lambda^k(M)\}$
- in a neighborhood around a point p, $\{dx^i\}$ is a basis for $T_p^* M$
so a section $\omega \in \Omega^k(M)$ can be written locally as
$$\omega = \omega_I \cdot dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

(ω_I are smooth functions)

Differential Forms

- Define the space of alternating k -tensors at p ,
 $\Lambda_p^k(M) = \Lambda^k(T_p M)$ and the bundle $\Lambda^k(M) = \coprod_{p \in M} \Lambda_p^k(M)$.
- define the space of "k-forms" as the sections of $\Lambda^k(M)$, that is
 $\Omega^k(M) = \{\omega \text{ section of } \Lambda^k(M)\}$
- in a neighborhood around a point p , $\{dx^i\}$ is a basis for T_p^*M
so a section $\omega \in \Omega^k(M)$ can be written locally as
$$\omega = \omega_I \cdot dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

(ω_I are smooth functions)
- Note: $\Omega^0(M) = C^\infty(M)$, $\Omega^1(M) = \mathfrak{T}^*(M)$ and $\Omega^k(M) = 0$ if
 $k > \dim(M)$

Differential Forms

Exterior Derivative

- We can define a unique linear map $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ with the following properties:

Differential Forms

Exterior Derivative

- We can define a unique linear map $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ with the following properties:
- 1) if f is smooth $df(X) = X(f)$
- 2) if $\omega \in \Omega^k(M), \eta \in \Omega^l(M)$ then
$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^k \omega \wedge d\eta$$
- 3) $d(d(\omega)) = 0$.

Differential Forms

Exterior Derivative

- We can define a unique linear map $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ with the following properties:
 - 1) if f is smooth $df(X) = X(f)$
 - 2) if $\omega \in \Omega^k(M), \eta \in \Omega^l(M)$ then
$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^k \omega \wedge d\eta$$
 - 3) $d(d(\omega)) = 0$.
- It is non-trivial to show that this map exists and is unique, the proof can be found on page 215 of Lee.

Differential Forms

Exterior Derivative

- We can define a unique linear map $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ with the following properties:
 - 1) if f is smooth $df(X) = X(f)$
 - 2) if $\omega \in \Omega^k(M), \eta \in \Omega^l(M)$ then
$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^k \omega \wedge d\eta$$
 - 3) $d(d(\omega)) = 0$.
- It is non-trivial to show that this map exists and is unique, the proof can be found on page 215 of Lee.
- if $\{x^i\}$ are local coordinates around p , there is an explicit formulation for the map d .
$$d\left(\sum'_I \omega_I dx^{i_1} \wedge \dots \wedge dx^{i_k}\right) = \sum'_I \left[\sum_i \frac{\partial \omega_I}{\partial x^i} dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \right]$$

Differential Forms

Closed and Exact Forms

- if $d\omega = 0$, we call ω closed, and if there is an η such that $d\eta = \omega$ we say it is exact. If ω is exact, then it is closed, but the converse is not true. Measuring the extent of this failure is the point of de Rham Cohomology.

Differential Forms

Closed and Exact Forms

- if $d\omega = 0$, we call ω closed, and if there is an η such that $d\eta = \omega$ we say it is exact. If ω is exact, then it is closed, but the converse is not true. Measuring the extent of this failure is the point of de Rham Cohomology.
- Why is this important? As the notation $d\omega$ suggests, we are able to integrate these forms. Evaluating the integrals of these forms is implicitly linked to the differential of said forms by means of stokes theorem

$$\int_M d\omega = \int_{\partial M} \omega$$

Differential Forms

Closed and Exact Forms

- if $d\omega = 0$, we call ω closed, and if there is an η such that $d\eta = \omega$ we say it is exact. If ω is exact, then it is closed, but the converse is not true. Measuring the extent of this failure is the point of de Rham Cohomology.
- Why is this important? As the notation $d\omega$ suggests, we are able to integrate these forms. Evaluating the integrals of these forms is implicitly linked to the differential of said forms by means of stokes theorem
$$\int_M d\omega = \int_{\partial M} \omega$$
- For example, if $\gamma : I \rightarrow M$ is a smooth curve, and ω a 1-form, then we can evaluate $\int_\gamma \omega$ by means of the fundamental theorem of calculus when ω is exact.

de Rham Cohomology

The Cochain Complex

- the vector spaces $\Omega^k(M)$ along with the homomorphism $d_k : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ define a cochain complex.

de Rham Cohomology

The Cochain Complex

- the vector spaces $\Omega^k(M)$ along with the homomorphism $d_k : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ define a cochain complex.
- define the de Rham cohomology group $H_{dR}^k(M) = \text{Ker}(d_{k+1}) / \text{Im}(d_k)$

de Rham Cohomology

The Cochain Complex

- the vector spaces $\Omega^k(M)$ along with the homomorphism $d_k : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ define a cochain complex.
- define the de Rham cohomology group
$$H_{dR}^k(M) = \text{Ker}(d_{k+1}) / \text{Im}(d_k)$$
- If $F : M \rightarrow N$ smooth, then $F^* : \Omega^k(N) \rightarrow \Omega^k(M)$, if we can show that $F^*d = dF^*$ then we can induce a map
$$H(F) : H_{dR}^k(N) \rightarrow H_{dR}^k(M)$$

de Rham Cohomology

The Cochain Complex

- the vector spaces $\Omega^k(M)$ along with the homomorphism $d_k : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ define a cochain complex.
- define the de Rham cohomology group
$$H_{dR}^k(M) = \text{Ker}(d_{k+1}) / \text{Im}(d_k)$$
- If $F : M \rightarrow N$ smooth, then $F^* : \Omega^k(N) \rightarrow \Omega^k(M)$, if we can show that $F^*d = dF^*$ then we can induce a map
$$H(F) : H_{dR}^k(N) \rightarrow H_{dR}^k(M)$$
- But this is easy, since
$$\begin{aligned} G^*d(fd x^{i_1} \wedge \dots \wedge dx^{i_k}) &= G^*(df \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}) = \\ d(f \circ G) \wedge d(x^{i_1} \circ G) \wedge \dots \wedge d(x^{i_k} \circ G) &= \\ d((f \circ G)d(x^{i_1} \circ G) \wedge \dots \wedge d(x^{i_k} \circ G)) &= dG^*(fd x^{i_1} \wedge \dots \wedge dx^{i_k}) \end{aligned}$$

de Rham Cohomology

Invariance of cohomology

- it's simple to see that $H_{dR}^k : \text{Man}^\infty \rightarrow \mathfrak{Ab}$ is a contravariant functor, so the homology groups are invariant under diffeomorphism.

de Rham Cohomology

Invariance of cohomology

- it's simple to see that $H_{dR}^k : \text{Man}^\infty \rightarrow \mathfrak{Ab}$ is a contravariant functor, so the homology groups are invariant under diffeomorphism.
- What's more is that they are actually invariant under homotopy equivalence, yes the *continuous* homotopy equivalence. This means that de Rham cohomology is actually a topological invariant

de Rham Cohomology

Invariance of cohomology

- it's simple to see that $H_{dR}^k : \text{Man}^\infty \rightarrow \mathfrak{Ab}$ is a contravariant functor, so the homology groups are invariant under diffeomorphism.
- What's more is that they are actually invariant under homotopy equivalence, yes the *continuous* homotopy equivalence. This means that de Rham cohomology is actually a topological invariant
- This is the first clue toward the central theorem of de Rham, that the singular cohomology of a smooth manifold is isomorphic to it's de Rham cohomology.

de Rham Cohomology

Proof of Homotopy Equivalence

- We will prove homotopy equivalence by constructing a chain homotopy

de Rham Cohomology

Proof of Homotopy Equivalence

- We will prove homotopy equivalence by constructing a chain homotopy
- for $X \in T_p M$ define $X \lrcorner : \Lambda_p^k M \rightarrow \Lambda_p^{k-1} M$, the "contraction by X " by $X \lrcorner \omega(Y_1, \dots, Y_{k-1}) = \omega(X, Y_1, \dots, Y_{k-1})$
- Let $i_t : M \rightarrow M \times I$ be the injection $i_t(p) = (p, t)$

de Rham Cohomology

Proof of Homotopy Equivalence

- We will prove homotopy equivalence by constructing a chain homotopy
- for $X \in T_p M$ define $X \lrcorner : \Lambda_p^k M \rightarrow \Lambda_p^{k-1} M$, the "contraction by X " by $X \lrcorner \omega(Y_1, \dots, Y_{k-1}) = \omega(X, Y_1, \dots, Y_{k-1})$
- Let $i_t : M \rightarrow M \times I$ be the injection $i_t(p) = (p, t)$
- then define $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ by $h(\omega) = \int_0^1 (\frac{\partial}{\partial t} \lrcorner \omega) dt$

de Rham Cohomology

Proof of Homotopy Equivalence

- We will prove homotopy equivalence by constructing a chain homotopy
- for $X \in T_p M$ define $X \lrcorner : \Lambda_p^k M \rightarrow \Lambda_p^{k-1} M$, the "contraction by X " by $X \lrcorner \omega(Y_1, \dots, Y_{k-1}) = \omega(X, Y_1, \dots, Y_{k-1})$
- Let $i_t : M \rightarrow M \times I$ be the injection $i_t(p) = (p, t)$
- then define $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ by $h(\omega) = \int_0^1 (\frac{\partial}{\partial t} \lrcorner \omega) dt$
- we will show that for $\omega \in \Omega^{k-1}(M \times I)$,
$$h(d\omega) + d(h\omega) = i_1^* \omega - i_0^* \omega$$

de Rham Cohomology

Proof of Homotopy Equivalence

- We will prove homotopy equivalence by constructing a chain homotopy
- for $X \in T_p M$ define $X \lrcorner : \Lambda_p^k M \rightarrow \Lambda_p^{k-1} M$, the "contraction by X " by $X \lrcorner \omega(Y_1, \dots, Y_{k-1}) = \omega(X, Y_1, \dots, Y_{k-1})$
- Let $i_t : M \rightarrow M \times I$ be the injection $i_t(p) = (p, t)$
- then define $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ by $h(\omega) = \int_0^1 (\frac{\partial}{\partial t} \lrcorner \omega) dt$
- we will show that for $\omega \in \Omega^{k-1}(M \times I)$,
 $h(d\omega) + d(h\omega) = i_1^* \omega - i_0^* \omega$
- this will then imply that when $H : F \rightarrow G$ is a homotopy, $F = H \circ i_0$, $G = H \circ i_1$ and $\tilde{h} = h \circ H^* : \Omega^k(N) \rightarrow \Omega^{k-1}(M)$

de Rham Cohomology

Proof of Homotopy Equivalence

- We will prove homotopy equivalence by constructing a chain homotopy
- for $X \in T_p M$ define $X \lrcorner : \Lambda_p^k M \rightarrow \Lambda_p^{k-1} M$, the "contraction by X " by $X \lrcorner \omega(Y_1, \dots, Y_{k-1}) = \omega(X, Y_1, \dots, Y_{k-1})$
- Let $i_t : M \rightarrow M \times I$ be the injection $i_t(p) = (p, t)$
- then define $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ by $h(\omega) = \int_0^1 (\frac{\partial}{\partial t} \lrcorner \omega) dt$
- we will show that for $\omega \in \Omega^{k-1}(M \times I)$,
$$h(d\omega) + d(h\omega) = i_1^* \omega - i_0^* \omega$$
- this will then imply that when $H : F \rightarrow G$ is a homotopy, $F = H \circ i_0$, $G = H \circ i_1$ and $\tilde{h} = h \circ H^* : \Omega^k(N) \rightarrow \Omega^{k-1}(M)$
- We will then find that \tilde{h} is the chain homotopy we are looking for, that is:
$$\tilde{h}(d\omega) + d(\tilde{h}\omega) = F^* \omega - G^* \omega$$

de Rham Cohomology

Proof of homotopy equivalence

- We need to show that as $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ is defined:

$$h(d\omega) + d(h\omega) = i_1^*\omega - i_0^*\omega$$

de Rham Cohomology

Proof of homotopy equivalence

- We need to show that as $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ is defined:

$$h(d\omega) + d(h\omega) = i_1^*\omega - i_0^*\omega$$

- explicitly:

$$(h\omega)_q(X_1, \dots, X_{k-1}) = \int_0^1 \left(\frac{\partial}{\partial t} \lrcorner \omega_{(q,t)} \right) (X_1, \dots, X_{k-1}) dt = \int_0^1 \omega_{(q,t)} \left(\frac{\partial}{\partial t}, X_1, \dots, X_{k-1} \right) dt$$

de Rham Cohomology

Proof of homotopy equivalence

- We need to show that as $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ is defined:

$$h(d\omega) + d(h\omega) = i_1^*\omega - i_0^*\omega$$

- explicitly:

$$(h\omega)_q(X_1, \dots, X_{k-1}) = \int_0^1 \left(\frac{\partial}{\partial t} \lrcorner \omega_{(q,t)} \right) (X_1, \dots, X_{k-1}) dt = \int_0^1 \omega_{(q,t)} \left(\frac{\partial}{\partial t}, X_1, \dots, X_{k-1} \right) dt$$

- but we need only show this holds for basis elements, there are two cases of basis elements, $\omega = f(x, t)dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{k-1}}$ and $\omega = f(x, t)dx^{i_1} \wedge \dots \wedge dx^{i_k}$

de Rham Cohomology

Proof of homotopy equivalence

- We need to show that as $h : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ is defined:

$$h(d\omega) + d(h\omega) = i_1^*\omega - i_0^*\omega$$

- explicitly:

$$(h\omega)_q(X_1, \dots, X_{k-1}) = \int_0^1 \left(\frac{\partial}{\partial t} \lrcorner \omega_{(q,t)} \right) (X_1, \dots, X_{k-1}) dt = \int_0^1 \omega_{(q,t)} \left(\frac{\partial}{\partial t}, X_1, \dots, X_{k-1} \right) dt$$

- but we need only show this holds for basis elements, there are two cases of basis elements, $\omega = f(x, t)dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{k-1}}$ and $\omega = f(x, t)dx^{i_1} \wedge \dots \wedge dx^{i_k}$
- I'll do the second case, as it is more enlightening, the first case is rather simple however.

de Rham Cohomology

Proof of homotopy equivalence

since ω does not have any dt term, $\frac{\partial}{\partial t} \lrcorner \omega = 0$ thus $h\omega = 0$ so $d(h\omega) = 0$.

On the other hand

$$h(d\omega) = h\left(\sum_i \frac{\partial f}{\partial t} dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right)$$

de Rham Cohomology

Proof of homotopy equivalence

since ω does not have any dt term, $\frac{\partial}{\partial t} \lrcorner \omega = 0$ thus $h\omega = 0$ so $d(h\omega) = 0$.

On the other hand

$$\begin{aligned} h(d\omega) &= h\left(\sum_i \frac{\partial f}{\partial t} dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right) \\ &= h\left(\frac{\partial f}{\partial t} dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right) \end{aligned}$$

de Rham Cohomology

Proof of homotopy equivalence

since ω does not have any dt term, $\frac{\partial}{\partial t} \lrcorner \omega = 0$ thus $h\omega = 0$ so $d(h\omega) = 0$.

On the other hand

$$h(d\omega) = h(\sum_i \frac{\partial f}{\partial t} dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k})$$

$$= h(\frac{\partial f}{\partial t} dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k})$$

$$= \int_0^1 (\frac{\partial f}{\partial t} dt) dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

de Rham Cohomology

Proof of homotopy equivalence

since ω does not have any dt term, $\frac{\partial}{\partial t} \lrcorner \omega = 0$ thus $h\omega = 0$ so $d(h\omega) = 0$.

On the other hand

$$h(d\omega) = h\left(\sum_i \frac{\partial f}{\partial t} dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right)$$

$$= h\left(\frac{\partial f}{\partial t} dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right)$$

$$= \int_0^1 \left(\frac{\partial f}{\partial t} dt\right) dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

$$= (f(x, 1) - f(x, 0)) dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

de Rham Cohomology

Proof of homotopy equivalence

since ω does not have any dt term, $\frac{\partial}{\partial t} \lrcorner \omega = 0$ thus $h\omega = 0$ so $d(h\omega) = 0$.

On the other hand

$$\begin{aligned} h(d\omega) &= h\left(\sum_i \frac{\partial f}{\partial t} dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right) \\ &= h\left(\frac{\partial f}{\partial t} dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\right) \\ &= \int_0^1 \left(\frac{\partial f}{\partial t} dt\right) dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ &= (f(x, 1) - f(x, 0)) dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ &= i_1^* \omega - i_0^* \omega \end{aligned}$$

de Rham Cohomology

Calculating some de Rham groups

- $H_{dR}^0(M) = R$ when M is connected
- $H_{dR}^k(M) = 0$ when $k > \dim(M)$
- $H^K(\{p\}) = 0$ for $k > 0$.
- $H^K(M) = 0$ for $k > 0$ if M is contractible.

de Rham Cohomology

Calculating some de Rham groups

- $H_{dR}^0(M) = R$ when M is connected
 $H_{dR}^k(M) = 0$ when $k > \dim(M)$
 $H^K(\{p\}) = 0$ for $k > 0$.
 $H^K(M) = 0$ for $k > 0$ if M is contractible.
- if $M = \coprod_i M_i$ then $H_{dR}^k(M) \cong \prod_i H_{dR}^k(M_i)$
This is due to the stronger statement that the pullbacks of inclusion maps ι_i^* induce an isomorphism
 $\Psi : \Omega^k(M) \rightarrow \prod_i \Omega^k(M_i)$ by $\omega \mapsto \{\iota_i^* \omega\}$


de Rham Cohomology

Calculating some de Rham groups

- $H_{dR}^0(M) = R$ when M is connected
 $H_{dR}^k(M) = 0$ when $k > \dim(M)$
 $H^K(\{p\}) = 0$ for $k > 0$.
 $H^K(M) = 0$ for $k > 0$ if M is contractible.
- if $M = \coprod_i M_i$ then $H_{dR}^k(M) \cong \prod_i H_{dR}^k(M_i)$
This is due to the stronger statement that the pullbacks of inclusion maps ι_i^* induce an isomorphism
 $\Psi : \Omega^k(M) \rightarrow \prod_i \Omega^k(M_i)$ by $\omega \mapsto \{\iota_i^* \omega\}$
- For $k \geq 1$, $H_{dR}^k(R^n) = 0$


de Rham Cohomology

Mayer-Vietoris Sequence



de Rham Cohomology

Mayer-Vietoris Sequence




consider the pullbacks i^*, j^*, k^*, l^* the following sequence is exact:

$$0 \rightarrow \Omega^k(X) \xrightarrow{k^* \oplus l^*} \Omega^k(U) \oplus \Omega^k(V) \xrightarrow{i^* - j^*} \Omega(U \cap V) \rightarrow 0$$

de Rham Cohomology

Mayer-Vietoris Sequence



consider the pullbacks i^*, j^*, k^*, l^* the following sequence is exact:

$$0 \rightarrow \Omega^k(X) \xrightarrow{k^* \oplus l^*} \Omega^k(U) \oplus \Omega^k(V) \xrightarrow{i^* - j^*} \Omega(U \cap V) \rightarrow 0$$

Which it turn yields the long exact sequence

$$\begin{aligned} \dots \xrightarrow{\Delta} H_{dR}^k(X) \xrightarrow{k^* \oplus l^*} H_{dR}^k(U) \oplus H_{dR}^k(V) \xrightarrow{i^* - j^*} H_{dR}^k(U \cap V) \xrightarrow{\Delta} \\ \xrightarrow{\Delta} H_{dR}^{k+1}(X) \rightarrow \dots \end{aligned}$$

de Rham Cohomology

Smooth Singular Cohomology

- define the abelian group $\mathfrak{C}_k^\infty(M) = Fr(\{C^\infty(\Delta^k, M)\})$

de Rham Cohomology

Smooth Singular Cohomology

- define the abelian group $\mathfrak{C}_k^\infty(M) = Fr(\{C^\infty(\Delta^k, M)\})$
- for $\sigma \in \mathfrak{C}_k^\infty(M)$ define $\partial_k : \mathfrak{C}_k^\infty(M) \rightarrow \mathfrak{C}_{k-1}^\infty(M)$ by
$$\partial_k(\sigma) = \sum_{i=1}^k (-1)^i \sigma \circ F_{i,k},$$
 where $F_{i,k}$ is the i -th face map.

de Rham Cohomology

Smooth Singular Cohomology

- define the abelian group $\mathfrak{C}_k^\infty(M) = Fr(\{C^\infty(\Delta^k, M)\})$
- for $\sigma \in \mathfrak{C}_k^\infty(M)$ define $\partial_k : \mathfrak{C}_k^\infty(M) \rightarrow \mathfrak{C}_{k-1}^\infty(M)$ by $\partial_k(\sigma) = \sum_{i=1}^k (-1)^i \sigma \circ F_{i,k}$, where $F_{i,k}$ is the i -th face map.
- From this chain complex, define the cochain complex $\mathfrak{C}_\infty^k(M) = Hom(\mathfrak{C}_k^\infty(M), R)$ with coboundary map $\delta^k = Hom(\partial_k, R)$ and retrieve the *smooth* singular cohomology.

de Rham Cohomology

Smooth Singular Cohomology

- define the abelian group $\mathfrak{C}_k^\infty(M) = Fr(\{C^\infty(\Delta^k, M)\})$
- for $\sigma \in \mathfrak{C}_k^\infty(M)$ define $\partial_k : \mathfrak{C}_k^\infty(M) \rightarrow \mathfrak{C}_{k-1}^\infty(M)$ by $\partial_k(\sigma) = \sum_{i=1}^k (-1)^i \sigma \circ F_{i,k}$, where $F_{i,k}$ is the i -th face map.
- From this chain complex, define the cochain complex $\mathfrak{C}_\infty^k(M) = Hom(\mathfrak{C}_k^\infty(M), R)$ with coboundary map $\delta^k = Hom(\partial_k, R)$ and retrieve the *smooth* singular cohomology.
- It can be shown that the normal singular cohomology with coefficients in R is equivalent to the smooth singular cohomology on smooth manifolds.

de Rham Cohomology

de Rham Theorem

- fix a k -form ω on M and a simplex σ . Pullback ω by σ and we have a k -form on Δ^K , define:

$$\Psi_k(\omega)(\sigma) = \int_{\sigma} \omega = \int_{\Delta^k} \sigma^* \omega$$

de Rham Cohomology

de Rham Theorem

- fix a k -form ω on M and a simplex σ . Pullback ω by σ and we have a k -form on Δ^K , define:

$$\Psi_k(\omega)(\sigma) = \int_{\sigma} \omega = \int_{\Delta^k} \sigma^* \omega$$

- this defines a homomorphism $\Psi_k : \Omega^k(M) \rightarrow \mathfrak{C}_{\infty}^k$
which, upon showing these maps commute with the boundary map and differential yields a chain map

$$\Psi : \Omega^*(M) \rightarrow \mathfrak{C}_{\infty}^*$$

de Rham Cohomology

de Rham Theorem

- fix a k -form ω on M and a simplex σ . Pullback ω by σ and we have a k -form on Δ^K , define:
$$\Psi_k(\omega)(\sigma) = \int_{\sigma} \omega = \int_{\Delta^k} \sigma^* \omega$$
- this defines a homomorphism $\Psi_k : \Omega^k(M) \rightarrow \mathfrak{C}_{\infty}^k$
which, upon showing these maps commute with the boundary map and differential yields a chain map
$$\Psi : \Omega^*(M) \rightarrow \mathfrak{C}_{\infty}^*$$

de Rham's Theorem

$H(\Psi) : H_{dR}(M) \rightarrow H_{\infty}(M)$ is an isomorphism for all smooth manifolds