

# **INLAND-SEBKHAS IN DER OSTPROVINZ SAUDI-ARABIENS: RÄUMLICHE VERÄNDERUNGEN DURCH ZUNEHMENDE LANDSCHAFTSDEGRADATION**

**HANS-JÖRG BARTH, REGENSBURG**

## **SUMMARY**

**Inland sabkhas in the Eastern Province of Saudi-Arabia – spatial changes caused by a decreasing vegetation cover.**

Exploitation of natural grazing resources during the last decades leads to a continuous decrease in vegetation cover in the Eastern Province of Saudi Arabia. In an area of unidirectional high energy-winds this development inevitably results in soil erosion and reactivation of fossil dunes. Inland-sabkhas seem to be one of the consequences of ongoing sand erosion. Sedimentological studies proved that inland-sabkhas in the central coastal lowlands of Saudi-Arabia are not residuals of former coastal-sabkhas which originated in the Pleistocene with a higher sea water level than today. Measurements along permanent transect lines showed that inland-sabkhas in degraded areas grow up to 5,4 m in one year in southeasterly directions. The analysis of two Landsat-TM images confirms the expansion of inland-sabkhas within the main wind direction (which is NNW to SSE) as well as sedimentation usually confined to the northern and northwestern sebkha edges. On the other hand inland-sabkhas surrounded by a higher vegetation cover are stabilized. In this case no movement of sabkha edges was observed. For that reason it is concluded that the dynamics of an inland-sabkha are to a high degree influenced by the geo-ecological conditions (ground water, vegetation, sand supply and climatic conditions) of the surrounding area. The origin of inland-sabkhas has to be related to the recent geomorphological history and seems to be a result of a negative sand budget in a subregional wind action system much intensified by overgrazing.

## **ZUSAMMENFASSUNG**

Verstärkter Viehbesatz, der die Tragfähigkeit der Ökosysteme um ein Vielfaches überschreitet, hat bereits seit mehreren Jahrzehnten einen kontinuierlichen Rückgang der Vegetationsdichten in den Ostarabischen Küstentiefländern zur Folge. Dies führt, in Kombination mit einem unimodalen Windsystem und hohen Windgeschwindigkeiten, zur Erosion der äolischen Lockersedimente und zur Reaktivierung ehemals stabiler, fossiler Längsdünenfelder. Die zahlreichen Inland-Sebkhas scheinen ebenfalls das Resultat des verstärkten Materialabtrags zu sein. Sedimentologische Unter-

## LITERATUR

- ALSDIRAWI, F. [Eds]: Range Management in Arid Zones. Proceedings of the Second International Conference on Range Management in the Arabian Gulf, Kuwait 3-6 March 1990, Kuwait, London, New York, 53-70
- ANTON, D. (1984): Aspects of Geomorphological Evolution; Paleosols and Dunes in Saudi Arabia. In: JADO, A. R. & ZÖTL, J. G. [Hrsg.]: Quaternary Period in Saudi Arabia. Springer-Verlag, Wien, New York. Vol. 2, 275-296
- ARAMCO (1984): Oceoanographic, Meterological Services, Survey Unit Dhahran
- BARTH, H.-J. (1998a): Status of Vegetation and an Assessment of the Impact of Overgrazing in an Area North of Jubail, Saudi Arabia. In: OMAR S. A. S., MISAK R., AL-AJMI D. & N. AL-AWADHI [Eds]: Sustainable development in arid zones. Balkema, R'dam, Brookfield, 435-450
- BARTH, H.-J. (1998b): Sebkhas als Ausdruck von Landschaftsdegradation im Zentralen Küstentiefland der Ostprovinz Saudi-Arabiens. Regensburger Geogr. Schriften, Heft 29. Regensburg.
- BREED, C. S., FRYBERGER, S. G., ANDREWS, S., MCCAULEY, C., LENNARTZ, F., GEBEL, D. & HORSTMAN, K. (1979): Regional Studies of Sand Seas, using Landsat (ERTS) Imagery. In: MCKEE, E. D. [Ed.]: A Study of Global Sand Seas. USGS Professional Paper 1052, 305-381
- FRYBERGER, S. G., AL-SARI, A. M., CLISHAM, T. J., RIZVI, S. A. R. & AL-HINAI, K. G. (1984): Wind Sedimentation in the Jafurah Sand Sea, Saudi Arabia, Sedimentology, 31, 413-431
- HOLM, D. A. (1960): Desert Geomorphology in the Arabian Peninsula. Science, 132:1369-1379.
- LANCASTER, N. (1995): Geomorphology of Desert Dunes. Routledge, London, New York
- MAINGUET, M. & F. DUMAY (1998): The concepts of global wind action system and sediment balance, keys for aeolian action and aersertification monitoring. In: OMAR S. A. S., MISAK R., AL-AJMI D. & N. AL-AWADHI [Eds]: Sustainable development in arid zones, p.127-141, Balkema
- MANDAVILLE, J. P. (1990): Flora of Eastern Saudi Arabia. London, New York.
- MORTIMER, C. E. (1996): Chemie. Stuttgart, New York.
- OGUZ, I., KHATTAB, G. AL-HINAI, WASIM, A. & A. Z. ALI (1998): Desert geomorphology. Eolian sand drift rates, and their role in design of sand control measures in the Abqaiq area, Saudi Arabia. In: OMAR S. A. S., MISAK R., AL-AJMI D. & N. AL-AWADHI [Eds]: Sustainable development in arid zones. Balkema, Rotterdam, Brookfield, 213-221
- SARRE, R. D. & CHANCEY, C. C. (1990): Size Segregation During Aeolian Saltation on Sand Dunes. In: Sedimentology, 37, 389-392
- SIDAHMED, A. E. (1995): Towards Sustainable Development of Rangelands/Livestock in Dryland Areas of the Near East and North Africa. In: OMAR, S. A. S., RAZZAQUE, M. A. &

Eingang des Manuskripts: 07.02.1999

Annahme des Manuskripts: 19.11.1999