

NNS Multivariate Dependence vs. multivariance package

Intro

There's a new companion package `multivariance` in R to research demonstrating multivariate dependence,¹

Distance multivariance is a measure of dependence which can be used to detect and quantify dependence.

NNS has had this capability for years!²³

We will show some known cases in 3 dimensions to illustrate the difference in capabilities between the two methods, then use the example found within the `multivariance` package.

Load Packages NNS (>= 4.5)

```
require(devtools); install_github('OVVO-Financial/NNS', ref = "NNS-Beta-Version")
library(NNS)
library(data.table)
library(rgl)
library(multivariance)
```

¹Dependence and Dependence Structures: Estimation and Visualization Using Distance Multivariance. <arXiv:1712.06532>.

²Deriving Nonlinear Correlation Coefficients from Partial Moments <https://ssrn.com/abstract=2148522>

³Beyond Correlation: Using the Elements of Variance for Conditional Means and Probabilities <https://ssrn.com/abstract=2745308>

Case 1: Total Dependence and Positive Correlation

The easiest way to demonstrate this is to show a straight line in 3 dimensions, where the variables would axiomatically be dependent upon one another. All observations would occupy the Co-Upper Partial Moment quadrant (green) or Co-Lower Partial Moment quadrant (red).

```
set.seed(123)

x <- rnorm(1000)


x3 <- cbind(x,x,x)

multicorrelation(x3)

## multicorrelation.2
## 1

NNS.dep.hd(x3, plot = TRUE, independence.overlay = TRUE)

## $actual.observations
## [1] 1000
##
## $independent.null
## [1] 250
##
## $Dependence
## [1] 1
```



Case 2: Total Dependence and Negative Correlation

This case too would be a straight line in 3 dimensions, however, it would span the divergent partial moment quadrants (non-highlighted ones).

```
set.seed(123)

x <- rnorm(1000)


x3 <- cbind(-x,x,x)

multicorrelation(x3)

## multicorrelation.2
##           1

NNS.dep.hd(x3, plot = TRUE, independence.overlay = TRUE)

## $actual.observations
## [1] 0
##
## $independent.null
## [1] 250
##
## $Dependence
## [1] 1
```



Case 3: Independence

3 Normal random variables...

```
set.seed(123)


x3 <- cbind(rnorm(1000),rnorm(1000),rnorm(1000))

multicorrelation(x3)

## multicorrelation.2
##      0.00407423

NNS.dep.hd(x3, plot = TRUE, independence.overlay = TRUE)

## $actual.observations
## [1] 267
##
## $independent.null
## [1] 250
##
## $Dependence
## [1] 0.02266667
```



So far so good!

Case 4: multivariate Example

```
y = rnorm(100)
x = cbind(y,y*2,(y-2)/3,y+1,y*5)

multicorrelation(x)

## multicorrelation.2
## 1
NNS.dep.hd(x, plot = TRUE, independence.overlay = TRUE)

## $actual.observations
## [1] 100
##
## $independent.null
## [1] 6.25
##
## $Dependence
## [1] 1
```

Still good! That's all of the observations within the Co-Partial Moment quadrants per Case 1.

Case 5: multivariate Example Expanded

Let's add some more terms to the matrix, and add more observations to eliminate any small sample concerns.

```
y = rnorm(1000)
x = cbind(y,y*2,(y-2)/3,y+1,y*5,y^2,-y^3,-3*y^4,(y^5-4)/3)

NNS.dep.hd(x, plot = TRUE, independence.overlay = TRUE)

## $actual.observations
## [1] 0
##
## $independent.null
## [1] 3.90625
##
## $Dependence
## [1] 1
multicorrelation(x)
```

```
## multicorrelation.2
## 0.6288725
```

Wait... what happened? All the variables are still very much related according to the pairwise NNS.dep measures.⁴

⁴Nonlinear Correlation and Dependence Using NNS <https://ssrn.com/abstract=3010414>

```
round(NNS.dep(x)$Dependence, 3)
```

```
##          y
## y 1.000 1.000 1.000 1.000 1.000 0.971 0.917 0.920 0.885
##   1.000 1.000 1.000 1.000 1.000 0.971 0.917 0.920 0.885
##   1.000 1.000 1.000 1.000 1.000 0.971 0.917 0.920 0.885
##   1.000 1.000 1.000 1.000 1.000 0.971 0.917 0.920 0.885
##   1.000 1.000 1.000 1.000 1.000 0.971 0.917 0.920 0.885
##   0.971 0.971 0.971 0.971 0.971 1.000 0.987 0.977 0.830
##   0.917 0.917 0.917 0.917 0.917 0.987 1.000 0.988 0.974
##   0.920 0.920 0.920 0.920 0.920 0.977 0.988 1.000 0.988
##   0.885 0.885 0.885 0.885 0.885 0.830 0.974 0.988 1.000
```

There was *not a single observation* in the Co-Partial Moment quadrants versus the expectation of 3.90625 total observations in those quadrants, per Case 2.

Let's increase the number of observations...

```
y = rnorm(10000)
x = cbind(y,y*2,(y-2)/3,y+1,y*5,y^2,-y^3,-3*y^4,(y^5-4)/3)
```

```
NNS.dep.hd(x, plot = TRUE, independence.overlay = TRUE)
```

```
## $actual.observations
## [1] 0
##
## $independent.null
## [1] 39.0625
##
## $Dependence
## [1] 1
```

Again, *not a single observation* in those quadrants. *In the absence of independence, there is dependence.*

Timing

Using the prior example settings of $n = 1,000$, let's compare the timing of the results for each.

```
library(microbenchmark)
y = rnorm(1000)
x = cbind(y,y*2,(y-2)/3,y+1,y*5,y^2,-y^3,-3*y^4,(y^5-4)/3)

microbenchmark(NNS=NNS.dep.hd(x),multivariate=multicorrelation(x),times = 100)
```

```
## Unit: milliseconds
##          expr          min          lq          mean          median          uq
##          NNS    2.267002    2.454802    2.775967    2.502701    2.742451
##  multivariate 256.859000 265.055950 269.075679 268.643450 270.605751
##          max neval
##    13.2757   100
##   346.8588   100
```

Comments

I look forward to further discussions and collaboration with those equally as passionate about these issues, and open to embracing alternative solutions. If you found this presentation interesting or useful, please feel free to reach out via e-mail: ovvo.financial.systems@gmail.com

Thanks for your interest!