


Section 11.3: Integral test

Ex. 1. Consider: $\sum_{k=1}^{\infty} \frac{1}{k^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots$

Then for $f(x) = \frac{1}{x^2}$, $f(k) = Q_k$



$$\sum_{k=1}^{\infty} \frac{1}{k^2} = \boxed{\text{Area of rectangles}} \leq \underbrace{1}_{\text{1st rectangle}} + \int_1^{\infty} \frac{1}{x^2}$$

$$\sum_{k=1}^{\infty} \underbrace{\frac{1}{k^2}}_{Q_k} \leq 1 + \int_1^{\infty} x^{-2} dx = 1 + \left(-x^{-1} \right) \Big|_1^{\infty} = 2.$$

So, any partial sum satisfies

$$S_n = \sum_{k=1}^n \frac{1}{k^2} \leq \sum_{k=1}^{\infty} \frac{1}{k^2} \leq 2$$

\Rightarrow The sequence of partial sums $\{s_n\}$'s bounded:

$$0 < S_n \leq 2$$

But the a_k are nonnegative, so

$$S_{n+1} - S_n = \sum_{k=1}^{n+1} a_k - \sum_{k=1}^n a_k = a_{n+1} > 0$$

$$\Rightarrow S_{n+1} > S_n.$$

The sequence of partial sums $\{s_n\}$ is increasing and bounded \Rightarrow convergent!


$$\Rightarrow \sum_{k=1}^{\infty} \frac{1}{k^2} \text{ is } \underline{\text{convergent}}.$$

(not to 2 though).

Ex. 2 Consider:

$$\sum_{k=1}^{\infty} \frac{1}{k^{1/2}} = \sum_{k=1}^{\infty} \frac{1}{\sqrt{k}}$$

Introduce $f(x) = \frac{1}{\sqrt{x}}$, then $\int_1^{\infty} \frac{1}{\sqrt{x}} dx = +\infty$



$$\sum_{k=1}^{\infty} \frac{1}{k^{1/2}} = \boxed{\text{Area of rectangles}} \geq \int_1^{\infty} \frac{1}{\sqrt{x}} dx =$$

$$= 2\sqrt{x} \Big|_1^{\infty} = \lim_{t \rightarrow \infty} 2\sqrt{x} \Big|_1^t = +\infty$$

$$\Rightarrow \sum_{k=1}^{\infty} \frac{1}{x^{1/2}} = +\infty, \quad \text{divergent.}$$

The integral test for convergence.

Suppose $f(x)$ is continuous, positive, decreasing on $[1, \infty)$, and

$$f(k) = a_k, \quad k \geq 1,$$

then

$$\sum_{k=1}^{\infty} a_k \text{ is convergent} \Leftrightarrow \int_1^{\infty} f(x) dx \text{ is convergent.}$$

Remark: these properties only need to hold for $k \geq k_0$.



