

Section 11.9: Representing functions as power series

Recall:

$$\sum_{k=0}^{\infty} x^k = \frac{1}{1-x} \quad |x| < 1$$

$f(x)$

Ex. 1 Express $\frac{1}{1-x^2}$ as a power series.

$$\frac{1}{1-x^2} = \frac{1}{1-(x^2)} = \sum_{k=0}^{\infty} (x^2)^k = \sum_{k=0}^{\infty} x^{2k}$$

$f(x^2)$

Replaced $x \mapsto x^2$.

Converges when $|x^2| < 1$

$$|x|^2 < 1$$

$$|x| < 1$$

✓

Ex. 2 Express as a power series: $\frac{1}{x+2}$

Know: $\frac{1}{1-x} = f(x)$

$$\frac{1}{x+2} = \frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})} =$$

$$= \frac{1}{2} \cdot \frac{1}{1-(-\frac{x}{2})} = \frac{1}{2} \sum_{k=0}^{\infty} \left(-\frac{x}{2}\right)^k \quad \text{Ⓢ}$$

$f(-\frac{x}{2})$

geometric series, converges when

$$\left|-\frac{x}{2}\right| < 1$$

$$\textcircled{=} \frac{1}{2} \cdot \sum_{k=0}^{\infty} (-1)^k \cdot \frac{x^k}{2^k} = \underbrace{\sum_{k=0}^{\infty} (-1)^k \cdot \frac{x^k}{2^{k+1}}}_{= \frac{1}{x+2}}$$

$$\left| -\frac{x}{2} \right| < 1$$

$$\left| \frac{x}{2} \right| < 1 \quad \cdot 2$$

$$|x| < 2$$

Then, the series is convergent when $|x| < 2$.

Ex. 3 Obtain the power series for

$$\frac{x^3}{x+2}$$

Know: $\frac{1}{x+2} = \sum_{k=0}^{\infty} (-1)^k \cdot \frac{x^k}{2^{k+1}} \quad \times x^3$

Then

$$\begin{aligned} \frac{x^3}{x+2} &= x^3 \sum_{k=0}^{\infty} (-1)^k \frac{x^k}{2^{k+1}} \\ &= \sum_{k=0}^{\infty} (-1)^k \frac{x^{k+3}}{2^{k+1}} \end{aligned}$$

Converges when the series for $\frac{1}{x+2}$ does, for $|x| < 2$.

Differentiation and integration of power series.

Thm. If $\sum_{k=0}^{\infty} c_k(x-a)^k$ has the radius

of convergence $R > 0$, then the function it defines:

$f(x) = \sum_{k=0}^{\infty} c_k(x-a)^k = c_0 + c_1(x-a) + c_2(x-a)^2 + \dots$
is differentiable and continuous, and, moreover, there holds:

1). $f'(x) = \sum_{k=0}^{\infty} k c_k(x-a)^{k-1} = c_1 + 2c_2(x-a) + \dots$

2). $\int f(x) = \sum_{k=0}^{\infty} c_k \frac{(x-a)^{k+1}}{k+1} = c_0(x-a) + c_1 \frac{(x-a)^2}{2} + c_2 \frac{(x-a)^3}{3} + \dots$

In addition, the radii of convergence of the series from 1). and 2). are equal to R . Their centers are still a .

In other words:

The intervals of convergence of the series in 1). and 2). are $(a-R, a+R)$, up to the endpoints.

Thus, differentiation and integration of a convergent power series does not change the radius of convergence.

Ex. 4 Express $\frac{1}{(1-x)^2} = g(x)$ as a power series.

Recall: We know how to represent $\frac{1}{1-x}$:

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k$$

$f(x)$

$$f'(x) = ((1-x)^{-1})' = (-1)(1-x)^{-2} \cdot (-1) = (1-x)^{-2} = g(x).$$

$$g(x) = f'(x) = \left(\sum_{k=0}^{\infty} x^k \right)' \stackrel{\text{Thm}}{=} \sum_{k=0}^{\infty} k \cdot x^{k-1}$$

We know: the radius of convergence is preserved under differentiation, so $\sum_{k=0}^{\infty} k \cdot x^{k-1}$ converges on $(0-1, 0+1)$

This is not saying anything about the convergence at the endpoints, $x = \pm 1$.

Exercise: does this series converge at ± 1 ?

Remark about indexing:

$$\sum_{k=0}^{\infty} k \cdot x^{k-1} = \sum_{k=1}^{\infty} k \cdot x^{k-1}$$

$$= 1 \cdot 1 + 2 \cdot x + 3 \cdot x^2 + 4 \cdot x^3 + \dots$$

By making the change $l = k-1$, ($k = l+1$) the above series is: $\sum_{l=0}^{\infty} (l+1) \cdot x^l$

Ex. 5 Find the power series for $\ln(1+x)$, determine its radius of convergence.

$$f(x) = \ln(1+x), \quad \text{then}$$

$$\begin{aligned} f'(x) &= \frac{1}{1+x} = \frac{1}{1-(-x)} = \sum_{k=0}^{\infty} (-x)^k \\ &= \sum_{k=0}^{\infty} (-1)^k \cdot x^k \end{aligned}$$

convergent when $|x| < 1$

$$f'(x) = \sum_{k=0}^{\infty} (-1)^k x^k \quad \text{when } |x| < 1$$

Integrate both sides:

$$f(x) = \int \frac{1}{1+x} dx = \ln(1+x)$$

$$= \int \sum_{k=0}^{\infty} (-1)^k x^k dx$$

$$= \sum_{k=0}^{\infty} \int (-1)^k x^k dx$$

$$= \sum_{k=0}^{\infty} (-1)^k \int x^k dx$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{x^{k+1}}{k+1} + C$$

→ Convergent for $|x| < 1$ (since integration preserves $R=1$)

Note: the endpoints $x = \pm 1$ need to be checked separately

$$f(x) = \ln(1+x) = \left(\text{let } l = k+1; k=l-1 \right)$$

$$= \sum_{l=1}^{\infty} \underbrace{(-1)^{l+1}}_{(-1)^k = (-1)^{l-1} = (-1)^{l+1} = (-1)^{l-1} \cdot (-1)^2} \frac{x^l}{l} + C$$

Let us determine C .

$$f(x) = \ln(1+x) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^k}{k} + C$$

Set $x=0$ \uparrow

$$\ln(1+0) = 0 = \sum_{k=1}^{\infty} (-1)^{k+1} \cdot \frac{0^k}{k} + C$$

$$0 = 0 + C \Rightarrow C = 0$$

Answer: $\boxed{\ln(1+x) = \sum_{k=0}^{\infty} (-1)^{k+1} \frac{x^k}{k} \quad |x| < 1}$

Example 6: Obtain the power series for $f(x) = \arctan(x)$.

$$f'(x) = \frac{1}{1+x^2} = \frac{1}{1-(-x^2)} = \frac{1}{1-x} \sum_{k=0}^{\infty} (-x^2)^k$$

$$= \sum_{k=0}^{\infty} (-1)^k \cdot x^{2k}$$

Converges when
 $| -x^2 | < 1$
 $|x|^2 < 1$
 $|x| < 1$

Integrate both sides of

$$f'(x) = \sum_{k=0}^{\infty} (-1)^k \cdot x^{2k} \quad |x| < 1$$

$$\begin{aligned} f(x) = \arctan x &= \int \sum_{k=0}^{\infty} (-1)^k x^{2k} dx \\ &= \sum_{k=0}^{\infty} \int (-1)^k x^{2k} dx \\ &= \sum_{k=0}^{\infty} (-1)^k \int x^{2k} dx \\ &= \sum_{k=0}^{\infty} (-1)^k \cdot \frac{x^{2k+1}}{2k+1} + C \end{aligned}$$

Converges for $|x| < 1$ (because integration preserves R).

Note: we are not saying anything about the endpoints of the convergence interval; $x = \pm 1$.

To determine C , set $x=0$

$$f(0) = \underbrace{\arctan(0)}_0 = \underbrace{\sum_{k=0}^{\infty} (-1)^k \frac{0^{2k+1}}{2k+1}}_0 + C$$

$$0 = 0 + C \Rightarrow C = 0$$

$$\arctan(x) = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{2k+1}$$

$$|x| < 1$$

$$\arctan(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

odd function;
 $f(-x) = -f(x)$

odd powers

$$\ln(1+x) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^k}{k}$$

$$|x| < 1$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} - \dots$$

Ex. 7 Expand $\int \frac{1}{1+x^7} dx$ into a power series.

$$\frac{1}{1+x^7} = \frac{1}{1-(-x^7)} \stackrel{\frac{1}{1-x}}{=} \sum_{k=0}^{\infty} (-1)^k \cdot x^{7k}$$

$$\sum_{k=0}^{\infty} \underbrace{(-x^7)^k}_{(-1)^k x^7}$$

converges when
 $|-x^7| < 1$

$$\frac{1}{1+x^7} = \sum_{k=0}^{\infty} (-1)^k x^{7k}, \quad |x| < 1$$

$$\begin{aligned} |x^7| &< 1 \\ |x|^7 &< 1 \\ |x| &< 1. \end{aligned}$$

Integrate both sides of this equality:

$$\int \frac{dx}{1+x^7} = \int \sum_{k=0}^{\infty} (-1)^k x^{7k} dx, \quad |x| < 1$$

$$= \sum_{k=0}^{\infty} \int (-1)^k x^{7k} dx$$

$$= \sum_{k=0}^{\infty} (-1)^k \int x^{7k} dx$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{x^{7k+1}}{7k+1} + C$$

converges when $|x| < 1$, since integration preserves the radius, so $R=1$.

The endpoints of the interval of convergence can change after integration.

Exercise: show that the series for $\int \frac{dx}{1+x^7}$ converges at $x=1$, diverges at $x=-1$.

Answer (to Ex. 7):

$$\int \frac{dx}{1+x^7} = \sum_{k=0}^{\infty} (-1)^k \frac{x^{7k+1}}{7k+1} + C, \quad |x| < 1.$$