

Notes on averaging chipod profiles and comparison to Chameleon data for EQ08 and EQ14

Andy Pickering

April 13, 2017

Contents

1	Overview	2
2	Comparing individual estimates of ϵ	3
3	Comparing time-averaged profiles of ϵ	5
4	γ computed from averaged quantities	8
5	How many profiles need to be averaged to converge?	10
6	Normalized eps vs chi plots	13
7	Summary	14

1 Overview

Comparing average profiles of χ pod estimates (assuming $\gamma = 0.2$) to Chameleon data for the EQ08 and EQ14 experiments. Individual χ pod estimates tend to be biased low, but if we average many profiles together do the average profiles agree?

2 Comparing individual estimates of ϵ




Figure 1: EQ08: Histogram of the ratio of ϵ estimates from χ pod method to the chameleon values, for χ pod method applied to 1m binned profiles. Estimates for each profile were averaged in 10m depth bins.




Figure 2: EQ14: Histogram of the ratio of ϵ estimates from χ pod method to the chameleon values, for χ pod method applied to 1m binned profiles, and applied to just patches. Estimates for each profile were averaged in 10m depth bins.

3 Comparing time-averaged profiles of ϵ




Figure 3: Profiles of ϵ , averaged over 500-cast chunks. Different lines are 1m binned Chameleon ('cham'), χ pod method applied over 1m bins ('bin'), and χ pod method applied to patches ('patch'). *Note χ pod estimates where $\log_{10}\epsilon > -4$ are discarded.* Title of each panel gives profile range used, as well as the number of good profiles actually in that range.




Figure 4: Profiles of ϵ , averaged over 500-cast chunks. Different lines are 1m binned Chameleon ('cham'), χ pod method applied over 1m bins ('bin'), and χ pod method applied to patches ('patch'). *Note χ pod estimates where $\log_{10}\epsilon > -4$ are discarded. Title of each panel gives profile range used, as well as the number of good profiles actually in that range.

4 γ computed from averaged quantities

If we compute gamma from time-averaged N^2, T_z, χ, ϵ do we get $\gamma = 0.2$? Estimates from the averaged data are larger (Figures 5,6) but still slightly less than 0.2 .




Figure 5: Boxplots of $\log_{10}[\gamma]$ for a set of profiles from EQ14. Left is for all 1m avg data. Right is for data from all profiles averaged in 10m bins. Horizontal dashed line indicates $\gamma = 0.2$.




Figure 6: Boxplots of $\log_{10}[\gamma]$ for a set of profiles from EQ14. Left is for all 1m avg data. Right is for data from all profiles averaged in 10m bins. Horizontal dashed line indicates $\gamma = 0.2$.

5 How many profiles need to be averaged to converge?

Next I wanted to see how many profiles we need to average for the χ pod profile to converge to the chameleon profile. Obviously this will depend on the specific profiles used and the characteristics of the turbulence, but here they seem to converge after about 100-150 profiles.




Figure 7: EQ08: Average profiles of ϵ from binned χ_{pod} method and chameleon, for different number of profiles (given by title).




Figure 8: EQ14: Average profiles of ϵ from binned χ_{pod} method and chameleon, for different number of profiles (given by title).

6 Normalized eps vs chi plots

Assuming that

$$\gamma = \frac{N^2 \chi}{2\epsilon < T_z >^2} \quad (1)$$

, plotting $[\epsilon/N\hat{2}]$ vs $[\chi/t_z^2]$ should follow a straight line with slope equal to $1/2\gamma$.




Figure 9: EQ08: 10m binned chameleon $\epsilon/N\hat{2}$ vs χ/t_z^2 . Lines show different values of γ .

7 Summary

- Individual (and 10m binned) χ pod estimates of ϵ are biased low compared to Chameleon values.
- When many profiles are averaged in 10m bins, the average profiles of ϵ agree better.
- γ computed from averaged (across profiles) N^2 , T_z , χ , and ϵ is larger, but not quite 0.2 (mean is around 0.1?)
- Average profiles seem to converge after about 100 profiles?