

Cancer evolution model based on Moran processes

Sergio Peignier¹

INSA Lyon, Biosciences department
BF2I Laboratory (UMR0203)
INSA Btiment Louis Pasteur - 11 avenue Jean Capelle
69621 Villeurbanne Cedex
`sergio.peignier@insa-lyon.fr`

1 Introduction

Neoplasia is a phenomenon involving the excessive and abnormal growth of tissue. Neoplasm are usually classified as "benign", "potentially malignant", or "malignant" (cancer). A single tumor is presumed to be a clonal population of cells, deriving from a unique clone. However this population of cells is often heterogeneous, containing different types of cells. In this context, genetic and epigenetic differences may confer selective advantages to a given individual, enabling it to expand. The dynamics of the populations of mutant clones are tailored by **natural selection** and **drift**, and the fitness of a clone often depends on the interaction with other cells, and its environment. Generally, the dynamics of the **clonal selection** leads to phenomena such as invasion, metastasis and therapeutic resistance. The importance of the evolutionary dynamics of the neoplasm clones, has leaded recently to an increasing interest in research, as reported for instance in [2, 3, 1]. Most of the works in this field, used stochastic process to model the evolutionary dynamics of the clones population. For instance, in [4], the authors use Moran processes to model the evolution of cancer cells.

In this project, we will focus on the genetic instability phenomenon, and its impact on the emergence of resistance to treatments. We refer the reader to [2] (Figure 2a)¹, for a graphical representation (Muller plot) of the frequency of clones in the population, and further explanations.

2 Methodology

In this work, we use a Moran process based model, in order to study the impact of genetic instability in the development of cancer, and its impact on the emergence of resistance to treatments. The major points of the model are detailed hereafter:

¹ https://www.researchgate.net/profile/John_Pepper3/publication/6688835_Cancer_as_an_evolutionary_and_ecological_process/links/00b4952aafdf67a39d000000/Cancer-as-an-evolutionary-and-ecological-process.pdf

- Constant population of size $N = 1000$.
- Two major types of cells coexist: normal cells and cancer cells.
- Among cancer cells, different clones may coexist.
- Each cancer clone c_i is characterized by its fitness f_{c_i} , its frequency n_{c_i} and its mutation rate μ_{c_i} .
- Normal cells have a fitness $f_n = 1$, a frequency $N - \sum_i n_i$, and a mutation rate $\mu_n = 0.001$.

The steps of the process are described hereafter:

- Select an individual with a probability proportional to the fitness f_i of its cell type.
- With a probability μ_i the new individual gives birth to a new cancer clone c_a , with a fitness f_a and a mutation rate μ_a .
- Otherwise, simply increment the frequency of the cell type of the individual that was selected.
- Choose a random individual and kill it.
- The cancer is detected as soon as $\sum_i n_{c_i} > \omega N$, with $\omega \in]0, 1]$.
- Then treatment against a chosen cancer clone c_i can be injected. This treatment reduced the fitness f_{c_i} , by multiplying it by α , during t generations, with $\alpha \in [0, 1]$.
- The experiments ends when $\sum_i n_{c_i} = 0$ (recovery) or $N - \sum_i n_{c_i} = 0$ (death) or $T_{death} = 0$ (natural death).

For the sake of simplicity we consider that $\forall i, \mu_{c_i} = \mu_c$ and $f_{c_i} = f_c$. At the beginning of the simulation, only a small number of cancer cells of type c_0 exist.

3 Questions

3.1 Individual based simulation

- Build a simulator for this model.
- Make a function that computes the diversity in the population, by means of the Shannon diversity index: $H = -\sum_{i=1} p_i \ln p_i$ where $p_i = \frac{n_i}{N}$.
- Set up an experimental protocol that could help you to grasp the impact of the genetic instability in the development of cancer, and the efficiency of the treatment, justify your choice of methodology.
- Execute your experiment(s).
- Analyze your results.

References

1. Beerenwinkel, N., Schwarz, R.F., Gerstung, M., Markowetz, F.: Cancer evolution: mathematical models and computational inference. *Systematic biology* **64**(1), e1–e25 (2014)
2. Merlo, L.M., Pepper, J.W., Reid, B.J., Maley, C.C.: Cancer as an evolutionary and ecological process. *Nature Reviews Cancer* **6**(12), 924 (2006)

3. Michor, F., Iwasa, Y., Nowak, M.A.: Dynamics of cancer progression. *Nature reviews cancer* **4**(3), 197 (2004)
4. Michor, F., Iwasa, Y., Nowak, M.A.: The age incidence of chronic myeloid leukemia can be explained by a one-mutation model. *Proceedings of the National Academy of Sciences* **103**(40), 14931–14934 (2006)