


Solving Mixed Integer Linear Programs with Cutting Planes

Tobias Kohler

May 16, 2024

Mixed Integer Linear Program and Linear Program Relaxation



MILP (standard form)

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & x \in F \\ & := \{x \mid Ax \leq b, x \in \mathbb{Z}_{\geq 0}^{n_1} \times \mathbb{R}_{\geq 0}^{n-n_1}\} \end{aligned}$$

$$c, x \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m$$

Solving Mixed Integer Linear Programs with Cutting Planes

LP Relaxation

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & x \in P \\ & := \{x \mid Ax \leq b, x \in \mathbb{R}_{\geq 0}^n\} \end{aligned}$$


Tobias Kohler

2/26

MILPs


- $n_1 = 0 \Rightarrow$ Linear Program (LP)
- $n_1 = n \Rightarrow$ Integer Linear Program (ILP)
- $x_i \in \{0, 1\} \Rightarrow$ (Mixed) Binary Program ((M)BP)

Observations




- In general, $\text{round}(\hat{x}) \neq x^*$

Observations




- In general, $\text{round}(\hat{x}) \neq x^*$
- The relaxed solution gives a lower bound:
 $c^\top \hat{x} \leq c^\top x^*$.

Observations



- In general, $\text{round}(\hat{x}) \neq x^*$
- The relaxed solution gives a lower bound:
 $c^\top \hat{x} \leq c^\top x^*$.
- \hat{x} can be found at a vertex of P (Simplex Algorithm).


Observations



- In general, $\text{round}(\hat{x}) \neq x^*$
- The relaxed solution gives a lower bound:
 $c^\top \hat{x} \leq c^\top x^*$.
- \hat{x} can be found at a vertex of P (Simplex Algorithm).
- If \hat{x} is already feasible, then $c^\top \hat{x} = c^\top x^*$.


Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem.



Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem.



Valid Inequalities and Cuts

- An inequality $a^\top x \leq r$ is valid for a set F if $a^\top x \leq r$ is satisfied for all $x \in F$.

Valid Inequalities and Cuts

- An inequality $a^\top x \leq r$ is valid for a set F if $a^\top x \leq r$ is satisfied for all $x \in F$.
 - For example: $x \leq 2$ is a valid inequality for $\{x \in \mathbb{Z}_{\geq 0} \mid x \leq 2.718\}$

Valid Inequalities and Cuts

- An inequality $a^\top x \leq r$ is valid for a set F if $a^\top x \leq r$ is satisfied for all $x \in F$.
 - For example: $x \leq 2$ is a valid inequality for $\{x \in \mathbb{Z}_{\geq 0} \mid x \leq 2.718\}$
- A cutting plane (or cut) w.r.t. $\hat{x} \in P \setminus F$ is any valid inequality $a^\top x \leq r$ for F such that:

$$a^\top \hat{x} > r$$

Cutting Planes Algorithm

```
1: LP  $\leftarrow$  Relaxation of the MILP
2: repeat
3:    $\hat{x} \leftarrow$  Optimal solution of the LP
4:   if  $(\hat{x}_1, \dots, \hat{x}_{n_1}) \notin \mathbb{Z}^{n_1}$  then
5:     Add a cut w.r.t.  $\hat{x}$  to the LP
6: until  $(\hat{x}_1, \dots, \hat{x}_{n_1}) \in \mathbb{Z}^{n_1}$ 
7: return  $\hat{x}$ 
```

Cutting Strategy

Question: How to generate “good” and useful cuts?

Cutting Strategy

Question: How to generate “good” and useful cuts?

- Good: Cut away as much as possible (while staying feasible)


Cutting Strategy

Question: How to generate “good” and useful cuts?

- Good: Cut away as much as possible (while staying feasible)
- Useful: Cut away the optimal solution of the relaxation

Convex Hull

- The relaxed solution \hat{x} in $\text{conv}(F)$ also solves the MILP.
- But computing the convex hull is infeasible (exponential).



Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

$$\blacksquare f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

- $f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$
- $\lfloor -a \rfloor = -\lceil a \rceil$ where
 $\lceil a \rceil = \min\{z \in \mathbb{Z} \mid z \geq a\}$

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

- $f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$
- $\lfloor -a \rfloor = -\lceil a \rceil$ where
 $\lceil a \rceil = \min\{z \in \mathbb{Z} \mid z \geq a\}$
- $a \in \mathbb{Z}$ and $a \leq b \Rightarrow a \leq \lfloor b \rfloor$

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

- $f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$
- $\lfloor -a \rfloor = -\lceil a \rceil$ where
 $\lceil a \rceil = \min\{z \in \mathbb{Z} \mid z \geq a\}$
- $a \in \mathbb{Z}$ and $a \leq b \Rightarrow a \leq \lfloor b \rfloor$
- $a \in \mathbb{Z}$ and $a \geq b \Rightarrow a \geq \lceil b \rceil$

Chvátal–Gomory Inequality for Integer Linear Programs

Let $\sum_{j=1}^n a_{ij}x_j \leq b_i$ for an Integer Linear Program ($x \in \mathbb{Z}_{\geq 0}^n$). Then the following inequalities are valid for any $\alpha \geq 0$:

$$\boxed{1} \quad \sum_{j=1}^n \alpha a_{ij}x_j \leq \alpha b_i \qquad \alpha \geq 0$$

Chvátal–Gomory Inequality for Integer Linear Programs

Let $\sum_{j=1}^n a_{ij}x_j \leq b_i$ for an Integer Linear Program ($x \in \mathbb{Z}_{\geq 0}^n$). Then the following inequalities are valid for any $\alpha \geq 0$:

$$1 \quad \sum_{j=1}^n \alpha a_{ij}x_j \leq \alpha b_i \quad \alpha \geq 0$$


$$2 \quad \sum_{j=1}^n \lfloor \alpha a_{ij} \rfloor x_j \leq \alpha b_i \quad x_j \geq 0$$

Chvátal–Gomory Inequality for Integer Linear Programs

Let $\sum_{j=1}^n a_{ij}x_j \leq b_i$ for an Integer Linear Program ($x \in \mathbb{Z}_{\geq 0}^n$). Then the following inequalities are valid for any $\alpha \geq 0$:


- | | | |
|---|--|----------------------|
| 1 | $\sum_{j=1}^n \alpha a_{ij}x_j \leq \alpha b_i$ | $\alpha \geq 0$ |
| 2 | $\sum_{j=1}^n \lfloor \alpha a_{ij} \rfloor x_j \leq \alpha b_i$ | $x_j \geq 0$ |
| 3 | $\sum_{j=1}^n \lfloor \alpha a_{ij} \rfloor x_j \leq \lfloor \alpha b_i \rfloor$ | $x_j \in \mathbb{Z}$ |

ILP vs. MILP



- $\min_{x,y} -y$
s.t. $\frac{3}{2}x + y \leq \frac{5}{4}, (x, y) \in \mathbb{Z}_{\geq 0}^2$
- $(x^*, y^*) = (0, 1)$
- $\lfloor \frac{3}{2} \rfloor \cdot 0 + \lfloor 1 \rfloor \cdot 1 = 1 \leq \lfloor \frac{5}{4} \rfloor \checkmark$

ILP vs. MILP



- $\min_{x,y} -y$
s.t. $\frac{3}{2}x + y \leq \frac{5}{4}, (x, y) \in \mathbb{Z}_{\geq 0} \times \mathbb{R}_{\geq 0}$
- $(x^*, y^*) = (0, \frac{5}{4})$
- $\lfloor \frac{3}{2} \rfloor \cdot 0 + \lfloor 1 \rfloor \cdot \frac{5}{4} = \frac{5}{4} > \lfloor \frac{5}{4} \rfloor$ ✗

Basic Mixed Integer Rounding Inequalities I

Let $x \in \mathbb{Z}_{\geq 0}$, $y \in \mathbb{R}_{\geq 0}$, $b \in \mathbb{R}_{>0} \setminus \mathbb{Z}$. Then

$$x \leq \lfloor b \rfloor \text{ is a valid inequality for } \{x + y \leq b\} \quad (1)$$

and

$$x \geq \lceil b \rceil \text{ is a valid inequality for } \{-x + y \leq -b\} \quad (2)$$

Basic Mixed Integer Rounding Inequalities I

Basic Mixed Integer Rounding Inequalities II

Let $x \in \mathbb{Z}_{\geq 0}$, $y \in \mathbb{R}_{\geq 0}$, $b \in \mathbb{R}_{>0} \setminus \mathbb{Z}$. Then

$$x - \frac{1}{f_b - 1} \leq \lfloor b \rfloor \text{ is a valid inequality for } \{x - y \leq b\} \quad (1)$$

and

$$x + \frac{1}{f_b} \geq \lceil b \rceil \text{ is a valid inequality for } \{-x - y \leq -b\} \quad (2)$$

Basic Mixed Integer Rounding Inequalities II

General Mixed Integer Rounding Inequality

Let $F_{MIR} = \{(x, y) \in \mathbb{Z}_{\geq 0}^2 \times \mathbb{R}_{\geq 0} \mid a_1x_1 + a_2x_2 - y \leq b\}$ where $a \in \mathbb{R}^2$, $b \in \mathbb{R} \setminus \mathbb{Z}$ and assume that $f_1 \leq f_b \leq f_2$. Then the inequality

$$\lfloor a_1 \rfloor x_1 + \left(\lfloor a_2 \rfloor + \frac{f_2 - f_b}{1 - f_b} \right) x_2 - \frac{1}{1 - f_b} y \leq \lfloor b \rfloor$$

is valid for F_{MIR} .

General Mixed Integer Rounding Inequality

Simplex Algorithm

Simplex finds $\hat{x} \in P \times \mathbb{R}_{\geq 0}^{N-n}$ and creates the optimal simplex tableau:

i -th row in the simplex tableau

$$x_{B_i} + \sum_{j \in NB} \bar{a}_{ij} x_j = \bar{b}_i$$

- x_1, \dots, x_{n_1} : Integer problem variables
- x_{n_1+1}, \dots, x_n : Real problem variables
- x_{n+1}, \dots, x_N : (Real) slack variables
- $B = \{B_1, \dots, B_m\}$: Basic variables
- $NB = \{1, \dots, N\} \setminus B$: Nonbasic variables ($\hat{x}_j = 0$ for $j \in NB$)

Gomory Mixed Integer Cut

Let $N_1 = NB \cap \{1, \dots, n_1\}$, $N_2 = NB \cap \{n_1 + 1, \dots, x_N\}$. Consider the i -th row in the optimal simplex tableau

$$x_{B_i} + \sum_{j \in N_1} \bar{a}_{ij} x_j + \sum_{j \in N_2} \bar{a}_{ij} x_j = \bar{b}_i$$

and assume $B_i \leq n_1$ but $\hat{x}_{B_i} = \bar{b}_i \notin \mathbb{Z}$. Then the Gomory Mixed Integer Cut

$$x_{B_i} + \sum_{\substack{j \in N_1 \\ f_{ij} \leq f_i}} \lfloor \bar{a}_{ij} \rfloor x_j + \sum_{\substack{j \in N_1 \\ f_{ij} > f_i}} \left(\lfloor \bar{a}_{ij} \rfloor + \frac{f_{ij} - f_i}{1 - f_i} \right) x_j + \sum_{\substack{j \in N_2 \\ \bar{a}_{ij} < 0}} \left(\frac{\bar{a}_{ij}}{1 - f_i} \right) x_j \leq \lfloor \bar{b}_i \rfloor$$

is a valid inequality for F that is not satisfied by \hat{x} .

Gomory Mixed Integer Cut

Cutting Planes Algorithm

- Let a MILP be given with feasible region $F = \{x \in \mathbb{Z}_{\geq 0}^{n_1} \times \mathbb{R}_{\geq 0}^{n-n_1} \mid Ax \leq b\}$ for some $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$.
- The relaxation is the LP obtained by removing the integer constraints, so its feasible region is the polyhedron $P = \{x \in \mathbb{R}_{\geq 0}^n \mid Ax \leq b\}$.
- Repeat the following two steps until $\hat{x} \in F$:
 - 1 Solve the LP using the Simplex Algorithm and obtain $\hat{x} \in P$
 - 2 If $\hat{x} \notin F$, pick $i \in \{1, \dots, n_1\}$ s.t. $\hat{x}_i \notin \mathbb{Z}$ and add the corresponding CMI-Cut to the LP.

Project Demonstration

Selecting Cutting Planes

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.

Selecting Cutting Planes

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Heuristic to evaluate the efficiency of a cutting plane (e.g. euclidean distance to \hat{x}).

Selecting Cutting Planes

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Heuristic to evaluate the efficiency of a cutting plane (e.g. euclidean distance to \hat{x}).
 - Add multiple cutting planes in each iteration.

Selecting Cutting Planes

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Heuristic to evaluate the efficiency of a cutting plane (e.g. euclidean distance to \hat{x}).
 - Add multiple cutting planes in each iteration.
- Other cutting plane strategies exist.

Branch & Bound

- Like Cutting Planes, we solve the problem relaxation and add constraints until an optimal solution has been found.
- Divide & Conquer

Branching

Cut off the non-integer neighborhood of $x_i^* \notin \mathbb{Z} \Rightarrow$ Two new relaxation problems

Bounding

- Initially, we only know that $-\infty < c^T x_{MILP}^* < \infty$ (not very helpful).
- Improve bounds until Upper Bound – Lower Bound $< \epsilon$:
 - Lower Bound: For any LP-Relaxation, we have $c^T x_{LP}^* \leq c^T x_{MILP}^*$
 - Upper Bound: By definition, $c^T x_{MILP}^* \leq c^T x$ for any feasible $x \in F_{MILP}$

Branch & Cut

- Hello World
- Cutting Planes + Branch & Bound = Branch & Cut

Questions?

References