

Solving Mixed Integer Linear Programs with Cutting Planes


Tobias Kohler

May 16, 2024

Mixed Integer Linear Program

- Optimize linear objective function s.t. linear constraints and some integer constraints.
- Sometimes, decision variables are discrete: Distribution of patients/supplies/vehicles... or binary/logical variables.

Mixed Integer Linear Program and Linear Program Relaxation



MILP (standard form)

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & x \in F \\ & := \{x \mid Ax \leq b, x \in \mathbb{Z}_{\geq 0}^{n_1} \times \mathbb{R}_{\geq 0}^{n-n_1}\} \end{aligned}$$

$$c, x \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m$$

Solving Mixed Integer Linear Programs with Cutting Planes

LP Relaxation

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & x \in P \\ & := \{x \mid Ax \leq b, x \in \mathbb{R}_{\geq 0}^n\} \end{aligned}$$


Tobias Kohler

3/29

Notation


- x_1, \dots, x_{n_1} : Integer variables
- x_{n_1}, \dots, x_n : Real variables
- F, x^* : Feasible region and optimal solution of the MILP
- P, \hat{x} : Feasible region and optimal solution of the LP-relaxation
($F = P \cap \mathbb{Z}_{\geq 0}^{n_1} \times \mathbb{R}_{\geq 0}^{n-n_1}$)
- $n_1 = 0 \Rightarrow$ Linear Program (LP)
- $n_1 = n \Rightarrow$ Integer Linear Program (ILP)
- $x_i \in \{0, 1\} \Rightarrow$ (Mixed) Binary Program ((M)BP)

Observations




- \hat{x} can be found at a vertex of P (Simplex Algorithm).

Observations



- \hat{x} can be found at a vertex of P (Simplex Algorithm).
- In general, $\text{round}(\hat{x}) \neq x^*$


Observations



- \hat{x} can be found at a vertex of P (Simplex Algorithm).
- In general, $\text{round}(\hat{x}) \neq x^*$
- If \hat{x} is already feasible, then $c^\top \hat{x} = c^\top x^*$.


Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem that cut off the relaxed solution.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem that cut off the relaxed solution.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem that cut off the relaxed solution.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem that cut off the relaxed solution.




Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem that cut off the relaxed solution.



Cutting Planes

Solve the problem relaxation. If integer constraints are violated, add additional inequalities to the problem that cut off the relaxed solution.



Valid Inequalities and Cuts

- An inequality $a^\top x \leq r$ is valid for a set F if $a^\top x \leq r$ is satisfied for all $x \in F$.

Valid Inequalities and Cuts

- An inequality $a^\top x \leq r$ is valid for a set F if $a^\top x \leq r$ is satisfied for all $x \in F$.
 - For example: $x \leq 2$ is a valid inequality for $\{x \in \mathbb{Z}_{\geq 0} \mid x \leq 2.718\}$

Valid Inequalities and Cuts

- An inequality $a^\top x \leq r$ is valid for a set F if $a^\top x \leq r$ is satisfied for all $x \in F$.
 - For example: $x \leq 2$ is a valid inequality for $\{x \in \mathbb{Z}_{\geq 0} \mid x \leq 2.718\}$
- A cutting plane (or cut) w.r.t. $\hat{x} \in P \setminus F$ is any valid inequality $a^\top x \leq r$ for F such that:

$$a^\top \hat{x} > r$$

Cutting Planes Algorithm

```
1: LP  $\leftarrow$  Relaxation of the MILP
2: repeat
3:    $\hat{x} \leftarrow$  Optimal solution of the LP
4:   if  $(\hat{x}_1, \dots, \hat{x}_{n_1}) \notin \mathbb{Z}^{n_1}$  then
5:     Add a cut w.r.t.  $\hat{x}$  to the LP
6: until  $(\hat{x}_1, \dots, \hat{x}_{n_1}) \in \mathbb{Z}^{n_1}$ 
7: return  $\hat{x}$ 
```

Cutting Strategy

Question: How to generate “good” and useful cuts?

Cutting Strategy

Question: How to generate “good” and useful cuts?

- Good: Cut away as much as possible (while staying feasible)


Cutting Strategy

Question: How to generate “good” and useful cuts?

- Good: Cut away as much as possible (while staying feasible)
- Useful: Cut away the optimal solution of the relaxation

Convex Hull

- The relaxed solution \hat{x} in $\text{conv}(F)$ also solves the MILP.
- But computing the convex hull is infeasible.
- Our goal is instead to approximate the convex hull in a neighborhood of x^* .



Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

$$\blacksquare f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

- $f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$
- $\lfloor -a \rfloor = -\lceil a \rceil$ where
 $\lceil a \rceil = \min\{z \in \mathbb{Z} \mid z \geq a\}$

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

- $f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$
- $\lfloor -a \rfloor = -\lceil a \rceil$ where
 $\lceil a \rceil = \min\{z \in \mathbb{Z} \mid z \geq a\}$
- $a \in \mathbb{Z}$ and $a \leq b \Rightarrow a \leq \lfloor b \rfloor$

Integer Part and Fractional Part

Any real number $a \in \mathbb{R}$ can be expressed as

$$a = \lfloor a \rfloor + f_a$$

for some unique $\lfloor a \rfloor \in \mathbb{Z}$ and $f_a \in [0, 1)$.

- $\lfloor a \rfloor = \max\{z \in \mathbb{Z} \mid z \leq a\}$ is the integer part of a .
- $f_a = a - \lfloor a \rfloor$ is the fractional part of a .

- $f_a = 0 \Leftrightarrow a = \lfloor a \rfloor \Leftrightarrow a \in \mathbb{Z}$
- $\lfloor -a \rfloor = -\lceil a \rceil$ where
 $\lceil a \rceil = \min\{z \in \mathbb{Z} \mid z \geq a\}$
- $a \in \mathbb{Z}$ and $a \leq b \Rightarrow a \leq \lfloor b \rfloor$
- $a \in \mathbb{Z}$ and $a \geq b \Rightarrow a \geq \lceil b \rceil$

Chvátal–Gomory Inequality for Integer Linear Programs

Let $\sum_{j=1}^n a_{ij}x_j \leq b_i$ for an Integer Linear Program ($x \in \mathbb{Z}_{\geq 0}^n$). Then the following inequalities are valid for any $\alpha \geq 0$:

$$\boxed{1} \quad \sum_{j=1}^n \alpha a_{ij}x_j \leq \alpha b_i \qquad \alpha \geq 0$$

Chvátal–Gomory Inequality for Integer Linear Programs

Let $\sum_{j=1}^n a_{ij}x_j \leq b_i$ for an Integer Linear Program ($x \in \mathbb{Z}_{\geq 0}^n$). Then the following inequalities are valid for any $\alpha \geq 0$:


- 1 $\sum_{j=1}^n \alpha a_{ij}x_j \leq \alpha b_i$ $\alpha \geq 0$
- 2 $\sum_{j=1}^n \lfloor \alpha a_{ij} \rfloor x_j \leq \alpha b_i$ $x_j \geq 0$

Chvátal–Gomory Inequality for Integer Linear Programs

Let $\sum_{j=1}^n a_{ij}x_j \leq b_i$ for an Integer Linear Program ($x \in \mathbb{Z}_{\geq 0}^n$). Then the following inequalities are valid for any $\alpha \geq 0$:


- | | | |
|---|--|----------------------|
| 1 | $\sum_{j=1}^n \alpha a_{ij}x_j \leq \alpha b_i$ | $\alpha \geq 0$ |
| 2 | $\sum_{j=1}^n \lfloor \alpha a_{ij} \rfloor x_j \leq \alpha b_i$ | $x_j \geq 0$ |
| 3 | $\sum_{j=1}^n \lfloor \alpha a_{ij} \rfloor x_j \leq \lfloor \alpha b_i \rfloor$ | $x_j \in \mathbb{Z}$ |

ILP vs. MILP



- $\min_{x,y} -y$
s.t. $\frac{3}{2}x + y \leq \frac{5}{4}, (x, y) \in \mathbb{Z}_{\geq 0}^2$
- $(x^*, y^*) = (0, 1)$
- $\lfloor \frac{3}{2} \rfloor \cdot 0 + \lfloor 1 \rfloor \cdot 1 = 1 \leq \lfloor \frac{5}{4} \rfloor$ ✓

ILP vs. MILP



- $\min_{x,y} -y$
s.t. $\frac{3}{2}x + y \leq \frac{5}{4}, (x, y) \in \mathbb{Z}_{\geq 0} \times \mathbb{R}_{\geq 0}$
- $(x^*, y^*) = (0, \frac{5}{4})$
- $\lfloor \frac{3}{2} \rfloor \cdot 0 + \lfloor 1 \rfloor \cdot \frac{5}{4} = \frac{5}{4} > \lfloor \frac{5}{4} \rfloor$ ✗

Basic Mixed Integer Rounding Inequalities I

Let $x \in \mathbb{Z}_{\geq 0}$, $y \in \mathbb{R}_{\geq 0}$, $b \in \mathbb{R}_{>0} \setminus \mathbb{Z}$. Then

$$x \leq \lfloor b \rfloor \text{ is a valid inequality for } \{x + y \leq b\} \quad (1)$$

and

$$x \geq \lceil b \rceil \text{ is a valid inequality for } \{-x + y \leq -b\} \quad (2)$$

Basic Mixed Integer Rounding Inequalities I

Basic Mixed Integer Rounding Inequalities II

Let $x \in \mathbb{Z}_{\geq 0}$, $y \in \mathbb{R}_{\geq 0}$, $b \in \mathbb{R}_{>0} \setminus \mathbb{Z}$. Then

$$x - \frac{1}{f_b - 1} \leq \lfloor b \rfloor \text{ is a valid inequality for } \{x - y \leq b\} \quad (1)$$

and

$$x + \frac{1}{f_b} \geq \lceil b \rceil \text{ is a valid inequality for } \{-x - y \leq -b\} \quad (2)$$

Basic Mixed Integer Rounding Inequalities II

General Mixed Integer Rounding Inequality

Let $F_{MIR} = \{(x, y) \in \mathbb{Z}_{\geq 0}^2 \times \mathbb{R}_{\geq 0} \mid a_1x_1 + a_2x_2 - y \leq b\}$ where $a \in \mathbb{R}^2$, $b \in \mathbb{R} \setminus \mathbb{Z}$ and assume that $f_1 \leq f_b \leq f_2$. Then the inequality

$$\lfloor a_1 \rfloor x_1 + \left(\lfloor a_2 \rfloor + \frac{f_2 - f_b}{1 - f_b} \right) x_2 - \frac{1}{1 - f_b} y \leq \lfloor b \rfloor$$

is valid for F_{MIR} .

General Mixed Integer Rounding Inequality

Simplex Algorithm

Simplex finds $\hat{x} \in P \times \mathbb{R}_{\geq 0}^{N-n}$ and creates the optimal simplex tableau:

i -th row in the simplex tableau

$$x_{B_i} + \sum_{j \in NB} \bar{a}_{ij} x_j = \bar{b}_i$$

- x_1, \dots, x_{n_1} : Integral decision variables
- x_{n_1+1}, \dots, x_n : Real decision variables
- x_{n+1}, \dots, x_N : (Real) slack variables
- $B = \{B_1, \dots, B_m\}$: Basic variables
- $NB = \{1, \dots, N\} \setminus B$: Nonbasic variables ($\hat{x}_j = 0$ for $j \in NB$)

Gomory Mixed Integer Cut

Let $N_1 = NB \cap \{1, \dots, n_1\}$, $N_2 = NB \cap \{n_1 + 1, \dots, N\}$. Consider the i -th row in the optimal simplex tableau

$$x_{B_i} + \sum_{j \in N_1} \bar{a}_{ij} x_j + \sum_{j \in N_2} \bar{a}_{ij} x_j = \bar{b}_i$$

and assume $B_i \leq n_1$ but $\hat{x}_{B_i} = \bar{b}_i \notin \mathbb{Z}$. Then the Gomory Mixed Integer Cut

$$x_{B_i} + \sum_{\substack{j \in N_1 \\ f_{ij} \leq f_i}} \lfloor \bar{a}_{ij} \rfloor x_j + \sum_{\substack{j \in N_1 \\ f_{ij} > f_i}} \left(\lfloor \bar{a}_{ij} \rfloor + \frac{f_{ij} - f_i}{1 - f_i} \right) x_j + \sum_{\substack{j \in N_2 \\ \bar{a}_{ij} < 0}} \left(\frac{\bar{a}_{ij}}{1 - f_i} \right) x_j \leq \lfloor \bar{b}_i \rfloor$$

is a valid inequality for F that is not satisfied by \hat{x} .

Gomory Mixed Integer Cut

Cutting Planes Algorithm

- Let a MILP be given with feasible region $F = \{x \in \mathbb{Z}_{\geq 0}^{n_1} \times \mathbb{R}_{\geq 0}^{n-n_1} \mid Ax \leq b\}$ for some $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$.
- The relaxation is the LP obtained by removing the integer constraints, so its feasible region is the polyhedron $P = \{x \in \mathbb{R}_{\geq 0}^n \mid Ax \leq b\}$.
- Repeat the following two steps until $\hat{x} \in F$:
 - 1 Solve the LP using the Simplex Algorithm and obtain $\hat{x} \in P$
 - 2 If the problem is infeasible ($P = \emptyset$), return INEASIBLE. If the problem is unbounded and no integer constraints are violated, return UNBOUNDED
 - 3 If $\hat{x} \notin F$, pick $i \in \{1, \dots, n_1\}$ s.t. $\hat{x}_i \notin \mathbb{Z}$ and add the corresponding GMI-Cut to the LP.

Project Demonstration

- Simplex Solver
- Mixed Integer Gomory Cut
- 2D Visualisation

Cutting Planes Selection

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.

Cutting Planes Selection

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Evaluate the efficiency of a cutting plane based on some heuristics (for example euclidean distance to \hat{x}).

Cutting Planes Selection

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Evaluate the efficiency of a cutting plane based on some heuristics (for example euclidean distance to \hat{x}).
 - Add multiple cutting planes in each iteration.

Cutting Planes Selection

- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Evaluate the efficiency of a cutting plane based on some heuristics (for example euclidean distance to \hat{x}).
 - Add multiple cutting planes in each iteration.
- Other cutting plane strategies exist:

Cutting Planes Selection


- Only adding an arbitrary, single cutting plane is very inefficient if the problem dimension is large.
 - Evaluate the efficiency of a cutting plane based on some heuristics (for example euclidean distance to \hat{x}).
 - Add multiple cutting planes in each iteration.
- Other cutting plane strategies exist:
 - Knapsack Covers or GUB (generalized upper bound) covers for binary programs.

Branch & Bound

- Like Cutting Planes, we solve the problem relaxation and add constraints until an optimal solution has been found.
- Divide & Conquer approach

Branching


Cut off the non-integer neighborhood of $\hat{x}_i \notin \mathbb{Z}$ and obtain two new relaxation problems, one with $x_i \leq \lfloor \hat{x}_i \rfloor$ and one with $x_i \geq \lceil \hat{x}_i \rceil$. The resulting data structure is a binary tree of problems.




Bounding

- Initially, we only know that $-\infty < c^T x^* < \infty$ (not very helpful).
- Improve bounds:
 - Lower Bound: For any LP-Relaxation, we have $c^T \hat{x} \leq c^T x^*$
 - Upper Bound: By definition, $c^T x^* \leq c^T x$ for any feasible $x \in F$
- For an optimal solution \hat{x} of a subproblem:
 - if $c^T \hat{x} \geq$ upper bound, prune tree (stop branching).
 - if \hat{x} is feasible, update upper bound and prune.
 - if \hat{x} is infeasible, update lower bound and branch.
 - stop if tree is completely pruned or upper bound $-$ lower bound $< \epsilon$.


Example




Example




Example



Example



Example



Subproblem and Branching Variable Selection Strategies

- Subproblem selection:
 - Depth-First: Descend quickly to obtain a good upper bound (obtain a feasible solution fast).
 - Best-First: Pick the active node with the current lower bound (obtain a good lower bound).
- Branching variable selection:
 - Most fractional: $i = \arg \max_{1 \leq i \leq n_1} \min(f_i, 1 - f_i)$ (f_i close to $\frac{1}{2}$).
 - Multiple variables at once.

Cutting Planes + Branch & Bound = Branch & Cut

- Cutting Planes or Branch & Bound on their own are inefficient in practice.
- Combine the two to Branch & Cut. This works like Branch & Bound but with additionally adding cuts before branching.
- Used most often in practice.

Questions?

References



Bixby, E. Robert, Mary Fenelon, Zonghao Gu, Ed Rothberg, and Roland Wunderling (2000). “MIP: Theory and Practice — Closing the Gap”. In: System Modelling and Optimization. Ed. by M. J. D. Powell and S. Scholtes. Boston, MA: Springer US, pp. 19–49. ISBN: 978-0-387-35514-6.



Goycoolea, Marcos (n.d.). “Cutting Planes for Large Mixed Integer Programming Models”. In: ().



“Integer Programming: 2nd Edition” (2020). In: Integer Programming. John Wiley and Sons, Ltd, pp. 1–34. ISBN: 9781119606475. DOI: <https://doi.org/10.1002/9781119606475.oth1>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781119606475.oth1>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/9781119606475.oth1>.



Papadimitriou, Christos and Kenneth Steiglitz (Jan. 1982). Combinatorial Optimization: Algorithms and Complexity. Vol. 32. ISBN: 0-13-152462-3. DOI: 10.1109/TASSP.1984.1164450.