

An Inductive Approach to the Erdős–Hajnal Property for Fixed Induced Paths

Yuwen Zhou

University of British Columbia Okanagan

7 October 2026

Abstract

We develop a consecutive-path induction for the Erdős–Hajnal property. Finite forcing templates, obtained by substitution and vertex-Ramsey amplification, yield a strong light-mixing statement from an exponent for the preceding path. Rooted Tooth recurrences in a fixed hereditary ambient class then combine with ordered-tail transfer and variable-scale sparsification. The final step uses leaf fibres and a product criterion for uniform blockades. We give the supporting structural and sampling proofs, retaining the actual integer lengths and restricted scales throughout. The resulting implication has been formalized in Lean 4: there, the Erdős–Hajnal property of every fixed path follows from Rödl’s theorem, the Erdős–Hajnal property of P_5 , and the comb lemma of Nguyen, Scott and Seymour.

Contents

1	Introduction	4
1.1	Formal verification	4
2	Definitions, orientations and standard inputs	5
I	Uniform template amplification and strong wonderfulness	7
3	Two-labelled lifts and the forcing invariant	7
4	Ramsey enlargement and multipartite homogenization	7
5	Arm growth and positive substitution exponents	8
6	Strong wonderfulness from the preceding path	9
II	The fixed-target structural implication	10
7	One ambient class and a quantified rooted contract	10
8	Ordered-tail transfer without circularity	10
9	Induction through rooted orders	11

10 The final comb oracle	11
11 Generalized niceness and the terminal conclusion	12
12 The induction on path length	13
A The general Tooth lemma and modular decomposition	14
A.1 An elementary antichain estimate	14
A.2 Find a heavy chain	15
A.3 A thick layer gives an early output or the clean core	15
A.4 All chain layers are thin	15
B Thin layers with perfect partial transversals	16
C The ambient-independent two-parameter RP5 Tooth	17
C.1 Local contracts and rooted module equivalence	17
C.2 An ordered frontier has the required one-sided root law	18
C.3 Simultaneous root-family purification	18
C.4 The interval argument and the forcing path	18
C.5 Quantitative frontier lemma	19
C.6 Full tree and its mass accounting	20
C.7 Finish both directions	20
C.8 Numerical closure	21
D A polynomial bridge from EH transversals	21
E RP6 frontiers and the third offending-pair purification	23
E.1 Quantitative conclusion	23
E.2 First and canonical RP5 purification	24
E.3 The new offending-pair pass	24
E.4 Restored monotonicity and a large blockade	25
E.5 Early outcomes and actual lengths	25
F The RP6 full-tree theorem with its actual floating scale	26
F.1 Inputs, constants, and statement	26
F.2 The full cut tree and its cutoff	27
F.3 Small components and mass on one complete chain	27
F.4 The negative direction	28
F.5 The positive direction	28
F.6 All order, scale, and actual-length checks	28
F.7 Fixed-ambient variant	29
G The RP_q recurrence in the same hereditary ambient class	29
G.1 Common lower-tooth contract	29
G.2 A quantified RP_q frontier statement	29

G.3	Two preparatory passes	30
G.4	Offending pairs and the RP_{q-2} pass	31
G.5	Order hypotheses and early outputs	31
G.6	The full RP_q tooth recurrence	32
G.7	The fixed-ambient requirement is essential	33
H	Variable-scale local oracles and the sparsification potential	33
H.1	A fixed Rödl start and generalized niceness	34
I	The arbitrary-pattern layout argument	35
J	Ordered tails and the relative ordered-class EH interface	36
J.1	Explicit bootstrap and layout parameters	36
J.2	Sufficient conditions for EH of the ordered class	37
J.3	How to use this in a rooted-path induction	37
K	Leaf fibres, cleanup and the terminal product argument	38
K.1	Normalization of the single-path viral input	38
K.2	Cleanup from weak sparsity to equal-width directed sparsity	38
K.3	The compact scale potential	39
K.4	Removing the fixed start	40

1 Introduction

For a graph G , write $\text{hom}(G) = \max\{\alpha(G), \omega(G)\}$. A fixed graph H has the Erdős–Hajnal property if some $\delta > 0$ satisfies $\text{hom}(G) \geq |G|^\delta$ for every nonempty graph with no induced copy of H . We study this property when H is a path.

Theorem 1.1. *For every integer $s \geq 1$, there is $\delta_s > 0$ such that every P_s -free graph G of order $n \geq 1$ satisfies $\text{hom}(G) \geq n^{\delta_s}$. The same assertion holds for $\overline{P}_s$ -free graphs. The exponent depends on s , but not on G or n .*

The induction uses $\text{EH}(P_6)$ from Theorem A of machine-qed, *The Erdős–Hajnal property for the six-vertex path*, repository version `v1.0-proof`, released 27 September 2026 [1]. This is the sole six-vertex base input. That result is unrefereed; its proof is formalized in Lean relative to three published statements, as described in Section 1.1. The arguments of this paper concern the passage from $\text{EH}(P_{s-1})$ to $\text{EH}(P_s)$.

For a fixed $s \geq 7$, Part I derives strong wonderfulness W_s from $\text{EH}(P_{s-1})$. Part II fixes the hereditary ambient class $\text{Forb}(P_s)$ and proves rooted Tooth statements $T_r^{(s)}$. The superscript records the ambient class and stays unchanged in every lower-order call. The dependencies are

$$\begin{aligned} \text{EH}(P_{s-1}) &\implies W_s, \\ T_r^{(s)} + W_s + \text{EH}(P_{s-1}) &\implies \text{EH}(E_r \cap \text{Forb}(P_s)), \\ T_{q-1}^{(s)} + T_{q-2}^{(s)} + \text{EH}(E_{q-1} \cap \text{Forb}(P_s)) &\implies T_q^{(s)}, \\ T_{s-2}^{(s)} + W_s + \text{EH}(P_{s-1}) &\implies \text{EH}(P_s). \end{aligned}$$

The ambient-independent RP5 theorem starts the rooted induction. The RP6 step uses RP5 twice and the general RP4 Tooth once; subsequent orders use a common lower-order contract. Ordered-tail transfer supplies the relative ordered-class exponent before each next rooted-order step. Thus neither $\text{EH}(P_s)$ nor virality of the ordered class is assumed in proving $\text{EH}(P_s)$.

The appendices contain the general Tooth proof, perfect-transversal sampling, the RP5 theorem, an EH-transversal bridge, the RP6 and RP q recurrences, the variable-scale bootstrap, the layout lemma, ordered-tail transfer, and the terminal product argument.

1.1 Formal verification

The implication behind Theorem 1.1 has been formalized in Lean 4 (version 4.34.1, with Mathlib). The final declaration, `AllPathsLocal.eh_all_paths`, states

$$(\forall s \geq 7 : \text{RodlCo}(s)) \wedge \text{NssComb} \wedge \text{EH}(P_6) \implies \forall s : \text{EH}(P_s).$$

Here $\text{EH}(P_s)$ is the statement of Theorem 1.1 for finite P_s -free graphs, with induced copies. $\text{RodlCo}(s)$ is Rödl’s theorem [7] for graphs with no induced $\overline{P}_s$, in the maximum-degree form of [6, Theorem 1.3] and Contract 2.5. NssComb is the comb lemma [6, Lemma 4.3]. For this declaration Lean reports only its three standard axioms (`propext`, `Classical.choice`, `Quot.sound`).

The Lean proof of $\text{EH}(P_6)$ in [1] assumes three statements as axioms: Rödl’s theorem for $\overline{P}_6$, the Erdős–Hajnal property of P_5 [6, Theorem 1.2], and the comb lemma. Combining it with the declaration above gives `AllPathsLocal.eh_all_paths_of_rod1`, whose only non-standard axioms are these three published statements, together with the hypothesis $\text{RodlCo}(s)$ for $s \geq 7$. Thus, in Lean, Theorem 1.1 follows from Rödl’s theorem, $\text{EH}(P_5)$ and the comb

lemma. None of these three inputs is itself formalized. The sparse induced-path extraction of Contract 2.6 is proved in [1].

The development adds 439 theorem declarations in 137 source files to the P_6 project. It was compiled on a Linux machine with the same Lean release and pinned dependencies, and the audit entry point `FinalAllLocal439` was replayed by `leanchecker --fresh`. Its source files contain no `sorry`, `admit`, `axiom`, `unsafe` or `native_decide`.

The formal proof follows the architecture of this paper, but departs from the written proofs at five points. No statement is changed.

- (i) The final step from generalized niceness to the Erdős–Hajnal property uses the argument of [6], already formalized in [1], with its house lemma replaced by the forcing template of Section 5. Appendix K is not used.
- (ii) $T_6^{(s)}$ is obtained from the general RPq recurrence of Appendix G, with the $RP5$ contract applied to the $RP4$ representatives, instead of Appendices E and F.
- (iii) Hayward’s theorem is not used. In the negative branch of Appendix C, a partial transversal ordered by layers has no induced P_4 whose earliest vertex is an endpoint. A greedy stable set taken in this order yields a proper colouring together with a clique of the same size, so $\alpha\omega$ is at least the order of the transversal, the only consequence of perfection used in the proof of Lemma B.1.
- (iv) Combs are extracted with the comb lemma [6, Lemma 4.3], as in [1], rather than with Contract 2.7.
- (v) The layout lemma of [1] replaces Lemma I.1. The tower construction of Lemma 4.2 is replaced by a forward construction that homogenizes only consecutive fibres, with the diagonal pairs of the two-labelled lifts chosen independently. All exponents are those chosen in the formalization rather than the constants displayed here.

Consequently, the comb extraction of [4], the virality theorem of [3] and Hayward’s theorem [5] are not inputs of the formal proof. Appendices E, F, I and K, the uses of Contracts 2.7 and 2.8, and the displayed constants are verified on paper only.

2 Definitions, orientations and standard inputs

Definition 2.1. All graphs are finite and simple. P_r is the path on r vertices. For a graph G , set $F = \overline{G}$, $\text{hom}(G) = \max(\alpha(G), \omega(G))$, and $Q(G) = \alpha(G)\omega(G)$. A graph is H -free when it has no induced copy of H . The class $\text{Forb}(H)$ is hereditary. An all-orders EH exponent $\kappa > 0$ for H means $\text{hom}(J) \geq |J|^\kappa$ for every nonempty H -free graph J .

Definition 2.2. A set S is ε -sparse when $\Delta(G[S]) \leq \varepsilon|S|$. It is ε -restricted when this holds in $G[S]$ or $F[S]$. Where explicitly stated, the stronger convention $\varepsilon(|S| - 1)$ is used. A pair B, A is directed ε -sparse from B to A if every vertex of B has at most $\varepsilon|A|$ neighbours in A . Bidirectional sparsity requires both directions; weak sparsity means only $e(A, B) \leq \varepsilon|A||B|$. Directed ε -sparsity from B to A gives $e(A, B) \leq \varepsilon|A||B|$, so it implies weak ε -sparsity. The converse is false, since weak sparsity bounds no individual degree.

Definition 2.3. A blockade is an ordered sequence of pairwise disjoint nonempty sets. Its length is its actual integer number of blocks and its width is their minimum order. It is pure if every pair is complete or anticomplete, and directed ε -semisparse if each pair is complete or directed ε -sparse from the later block to the earlier block. A weakly semisparse blockade uses weak sparsity instead. A complete blockade and an anticomplete blockade each have a uniform pair type. A general pure pattern can mix the two types. Every directed ε -semisparse blockade is thus weakly ε -semisparse, but not conversely. Each statement below specifies which of the two notions it uses.

Definition 2.4. For a root $u \notin Y$, $\text{RP}_r(u, Y)$ means that $F[Y \cup \{u\}]$ has no induced r -vertex path with endpoint u . The roots are vertices of the original host, and this exclusion refers to the original target Y . Restricting the target inherits the exclusion. E_r is the hereditary class of graphs that admit a linear order in which the first vertex of every induced P_r is not an endpoint.

An anticomponent is a component in the complement. A module is a set seen completely or anticompletely by each outside vertex. A strong module overlaps no module except by containment. Substitution $H[J_1, \dots, J_a]$ replaces the vertices of H by the graphs J_i , joining two fibres completely exactly on the edges of H .

Contract 2.5 (Fixed precision Rödl). For every fixed forbidden graph H and fixed $\eta > 0$, there is $\delta > 0$ such that every H -free graph has an η -restricted induced subgraph of order at least $\delta|G|$. Only fixed choices of precision are used [7, 2].

Contract 2.6 (Sparse induced-path extraction). If P is a path with $k \geq 2$ vertices, $0 < z \leq 1/(60k)$, and J is a z^2 -sparse P -free graph with $|J| \geq z^{-2}$, then J has an anticomplete blockade of length at least $1/z$ and width at least $\lfloor z^2|J| \rfloor \geq 1$ [2, Statement 3.1].

Remark. In [2] the blocks of a blockade may be empty, and Statement 3.1 there has no hypothesis on $|J|$. Blocks are nonempty in this paper, and then the hypothesis $|J| \geq z^{-2}$ cannot be omitted: a one-vertex graph J , with $P = P_2$ and $z = 1/120$, satisfies every other hypothesis but has no blockade of length 120. Under the hypothesis, $\lfloor z^2|J| \rfloor \geq 1$, so the blockade supplied by [2] has nonempty blocks. The hypothesis is verified at each use of Contract 2.6: in the proof of Theorem H.2, in Appendix H.1, and in Appendix J.1.

Contract 2.7 (Comb extraction). If $0 < \rho \leq y \leq 2^{-8}$, H is y^3 -sparse, and $|H| \geq y^{-4}$, then either H is $2y^4$ -sparse; or disjoint X, Z satisfy $|X| \geq y^4|H|$, $|Z| \geq (1 - 4y)|H|$, with Z directed ρ -sparse to X ; or there is a comb of integer length $1/y \leq \ell \leq \rho^{-2}$, teeth of order at least $y^4|H|/\ell^2$, roots full to their own teeth and anticomplete to the others, and a special vertex full to all teeth and anticomplete to all roots. All these sets and vertices are disjoint; root-root edges are unrestricted [4, Lemma 2.10].

Contract 2.8 (Virality and perfection inputs). The EH property of a single graph implies its viral property [3, Theorem 4]. We normalize this in Appendix K. Weakly chordal graphs are perfect [5].

The modular-decomposition facts used in Appendix A are stated there in the exact form required. The general Tooth lemma, the sampling lemmas, the layout argument, and the terminal product criterion are proved in the appendices.

Part I

Uniform template amplification and strong wonderfulness

3 Two-labelled lifts and the forcing invariant

Throughout Part I, fix $h \geq 2$, put $s = h+1$, and assume that P_h has an all-orders EH exponent $\kappa > 0$.

Definition 3.1. For a graph Q on $[q]$, a spanning subgraph M of Q , and t in $\{0, 1\}$, define $L_t(Q, M)$ on $v, z_1, \dots, z_q, b_1, \dots, b_q$: vz_i is an edge; vb_i is a nonedge; z_ib_i is an edge iff $t = 1$; for $i \neq j$, z_iz_j and z_ib_j are edges iff ij is in Q ; b_ib_j is an edge iff ij is in M . All graphs are simple; this specifies every pair.

An r -arm is an induced r -vertex path whose first vertex has label z and whose remaining $r - 1$ vertices have label b . It does not contain v . Prepending v to an r -arm gives an induced P_{r+1} .

Put $s = h + 1$. We construct graphs Q_r for $2 \leq r \leq h$ such that: (*) for every $M \subseteq Q_r$ and every t in $\{0, 1\}$, $L_t(Q_r, M)$ contains P_s or an r -arm. Consequently Q_h is a forcing template: every $L_t(Q_h, M)$ contains P_s . The same graph Q_h works for both diagonal types t . In fact it works when every diagonal z_ib_i is chosen independently: all arms constructed below are index-transversal, meaning that their vertices use distinct indices i . The base arm has this property, and both inductive path constructions preserve it. No diagonal adjacency is ever used. We keep the constant- t notation below to match the usual wonderfulness convention.

4 Ramsey enlargement and multipartite homogenization

Lemma 4.1 (Vertex Ramsey enlargement). *Write $A[B_1, \dots, B_a]$ for substitution, where $a = |A|$: each vertex i of A is replaced by B_i , and two replacement graphs are joined completely exactly when their corresponding vertices of A are adjacent.*

Let $R(A, c)$ be the c -fold lexicographic power of A , for integer $c \geq 1$: $R(A, 1) = A$; $R(A, c) = A[R(A, c-1), \dots, R(A, c-1)]$. Every c -colouring of $V(R(A, c))$ has a monochromatic induced A .

Proof. By induction on c . If some top-level fibre uses at most $c - 1$ colours, induction supplies a monochromatic A in that fibre. Otherwise every fibre uses all c colours. Choose one vertex of one fixed colour in each fibre; their quotient is an induced A . This also proves the claim when the colouring uses fewer than c colours. $\square$

Lemma 4.2 (Backward construction and forward selection). *Given a graph A and an integer $h \geq 2$, define $S_h, \dots, S_1$ backwards: $c_i = 2^{i-1 + \sum_{j>i} |S_j|}$; $S_i = R(A, c_i)$. In particular $c_h = 2^{h-1}$. Put $H = P_h[S_1, \dots, S_h]$.*

For every spanning subgraph M of H there are induced copies A_i of A inside S_i , for $i = 1, \dots, h$, such that between every pair A_i, A_j the M -relation is complete or anticomplete. The Q -relation (here $Q = H$) between consecutive fibres is complete, and between all other fibres is anticomplete.

Proof. Choose A_i in increasing order of i . At this stage each vertex in S_i has a single M -adjacency bit to each previously selected A_j : this bit is well defined because all vertices of A_j were selected to have the same adjacency vector on every vertex in all later S_k . Colour each vertex of S_i by these $i - 1$ bits and its adjacency vector on every vertex of $S_{i+1} \cup \dots \cup S_h$. There are at most c_i colours. The vertex Ramsey property gives a monochromatic induced A_i . It has uniform M -relation to each earlier A_j , and its vertices have identical M -adjacencies to each individual later vertex. This preserves the induction invariant. At the end all pairs are homogeneous in M , as required.

No uniformity of M across the original fibres is assumed. This construction forces that uniformity on copies of the prescribed A . $\square$

5 Arm growth and positive substitution exponents

Proposition 5.1 (The forcing template). *For each fixed $h \geq 2$, a finite graph Q_h can be constructed from P_h and P_2 by substitution such that every two-labelled lift of Q_h , including lifts with independent diagonal choices, contains an induced P_{h+1} . The forcing path uses distinct indices apart from its unindexed root.*

Proof. Take $Q_2 = P_2$. For any M and either t, z_1, b_2 is a 2-arm, so $(*)$ holds.

Suppose $2 \leq r < h$ and Q_r has $(*)$. Apply Lemma 4.2 with $A = Q_r$ and define $Q_{r+1} = P_h[S_1, \dots, S_h]$. Consider an arbitrary $M \subseteq Q_{r+1}$, either t , and suppose its lift contains no P_s . Homogenize M on copies $A_1, \dots, A_h$ of Q_r by Lemma 4.2. The induced lift on v together with the z/b vertices indexed by A_i is exactly $L_t(Q_r, M[A_i])$; hence every fibre has an r -arm.

For each $i = 1, \dots, h - 1$, let e_i be 1 if M is complete between A_i, A_{i+1} , and 0 if it is anticomplete. We claim $e_1 = \dots = e_{r-1} = 1$.

Indeed, suppose $e_i = 0$ for some $i \leq r - 1$. In fibre i , choose an r -arm $z, b_1, \dots, b_{r-1}$. Choose any b -labelled vertex B in fibre $i + 1$, and one z -labelled vertex Z_j in each fibre $j = i + 2, \dots, h$. The sequence

$$b_{r-1}, \dots, b_1, z, B, Z_{i+2}, \dots, Z_h$$

is an induced path on $r + h - i \geq h + 1 = s$ vertices. Its adjacencies and nonadjacencies follow from:

- the first r vertices form the reversed arm;
- B is adjacent to z , since consecutive Q -fibres are complete;
- B is anticomplete to all b_j of the arm, since $e_i = 0$;
- B is adjacent to Z_{i+2} , and the Z -tail follows P_h ;
- all remaining cross-pairs lie in nonconsecutive Q -fibres, and are nonedges of both Q and M .

Taking s consecutive vertices gives an induced P_s , a contradiction.

Thus the first $r - 1$ M -fibre pairs are complete. Select one z -labelled vertex Z in fibre $r + 1$ and one b -labelled vertex B_i in each fibre $i = 1, \dots, r$. The path $Z, B_r, B_{r-1}, \dots, B_1$ is an $(r + 1)$ -arm. Consecutive b -pairs are adjacent by $e_i = 1$; every nonconsecutive pair is in nonconsecutive P_h -fibres and is a nonedge. The z -vertex is adjacent only to B_r among the selected b -vertices. Here $r + 1 \leq h$, so every required fibre exists. This proves $(*)$ for $r + 1$.

Induction reaches Q_h . A hypothetical P_s -free lift of Q_h would have an h -arm. Prepending v gives $P_{h+1} = P_s$, the final contradiction.

The forcing P_s is also index-transversal apart from the unindexed v : when a fibre already supplies P_s this follows inductively, and both new path constructions use disjoint fibre indices. Thus the assertion with arbitrary individual diagonal values is proved simultaneously.

□

Lemma 5.2 (Quantitative one-vertex substitution). *We use the following elementary quantitative substitution bound. If H, J have all-orders EH exponents $\gamma, \delta > 0$ and $|H| = a$, then $H[u \leftarrow J]$ has the all-orders exponent $\Phi(a, \gamma, \delta) = 1/(a + 1/\delta + a/\gamma)$.*

Proof. Let G be $H[u \leftarrow J]$ -free with n vertices and largest homogeneous set of size r . Put $m = \lfloor r^{1/\gamma} \rfloor + 1$. If $m > n$ the conclusion follows immediately. Otherwise every m -subset of $V(G)$ contains H . Counting such subsets yields at least $(n/m)^a$ induced H -copies. Choose an isomorphism for each copy. By fixing the $a - 1$ labelled core vertices other than u , some common extension fibre has at least n/m^a vertices. Every extension has exactly the prescribed adjacency to the fixed core. This fibre is J -free, or else the fibre and core give $H[u \leftarrow J]$. Hence its size is at most $r^{1/\delta}$, and $n \leq m^a r^{1/\delta} \leq 2^a r^{a/\gamma + 1/\delta} \leq r^{a + a/\gamma + 1/\delta}$. The last step uses $r \geq 2$ when $n \geq 2$; $n = 1$ is immediate. □

This formula gives an explicit algorithm for every exponent used here: start with exponent 1 for $Q_2 = P_2$ and the input exponent κ for P_h . For each substitution of a graph J into a vertex of the current graph H , update its order to $|H| - 1 + |J|$ and update its exponent using Φ . Apply this for the recursively defined lexicographic powers $R(A, c)$, and then for $P_h[S_1, \dots, S_h]$, in any fixed vertex order. All numbers are finite positive quantities, computable by the displayed recursion. Let $q_h = |Q_h|$ and $c_h > 0$ be the resulting EH exponent for Q_h . Complementation preserves the same exponent.

The constants are finite and depend only on h and the input exponent κ .

6 Strong wonderfulness from the preceding path

Theorem 6.1 (Uniform strong wonderfulness). *Choose any integer $a \geq \max(2q_h, 2 + 5/c_h, 6)$. Let $0 < y < 1/2$. Suppose $F = \overline{G}$ is P_s -free, equivalently G is $\overline{P_s}$ -free, and $B_1, \dots, B_\ell$ are pairwise disjoint nonempty blocks of common size W , where $\ell \geq y^{-a}$, and each pair is either complete or bidirectionally θ -sparse, $\theta = y^a$. Here directed θ -sparse means that every vertex on the indicated side has at most $\theta \cdot$ the other side's size neighbours. This is stronger than a weak density condition.*

Then one of the following holds.

- (A) G contains a y^4 -restricted set with cardinality at least W .
- (B) Some B_i has at most $y|G|$ vertices outside the block union whose degree into B_i is strictly between 0 and $W/2$.

Proof. For an outside vertex u let $I(u)$ consist of the blocks on which its degree is strictly between 0 and $W/2$. Suppose $|I(u)| > y\ell$. For each i in $I(u)$, choose b_i adjacent to u in G . More than $W/2$ vertices of B_i are nonadjacent to u . At least half of them have the same adjacency t_i to b_i in $F = \overline{G}$. Let C_i be that set, so $|C_i| > W/4$. Restrict to at least $|I(u)|/2$ indices of one common type t .

Let J be their complete-pair graph in G . We claim J is $\overline{Q_h}$ -free. Otherwise select q_h blocks whose complete-pair graph is $\overline{Q_h}$. For every edge ij of Q_h the corresponding blocks are θ -sparse in G . Choose z_i in C_i one by one, avoiding G -neighbours of every b_j and of all previously chosen z_j across such pairs. At each step at most $2(q_h - 1)\theta W$ vertices are forbidden, which is less than $W/4$ because $a \geq 2q_h$ and $y < 1/2$. Thus every choice is possible.

In F the selected vertices u, z_i, b_i realize $L_t(Q_h, M)$, where M is the graph of F -edges among the b_i . It is a spanning subgraph of Q_h : a nonedge ij of Q_h corresponds to a complete

block pair in G . Proposition 5.1 forces P_s in F , contrary to the hypothesis. This proves the claim.

By EH for $\overline{Q_h}$, J has a homogeneous set of size at least $(|I(u)|/2)^{c_h} > (y\ell/2)^{c_h} \geq 2y^{-4}$. The final inequality follows from $a \geq 2 + 5/c_h$ and $y < 1/2$. Let $r \geq 2y^{-4}$ be its cardinality, and S the union of those blocks. If the pairs are complete, the maximum degree in $F[S]$ is at most $W - 1$. If the pairs are sparse, the maximum degree in $G[S]$ is at most $W - 1 + (r - 1)\theta W$. In either case it is at most $(1/r + \theta)|S| - 1 \leq (y^4/2 + y^6)|S| - 1 \leq y^4(|S| - 1)$. Hence S is y^4 -restricted and $|S| \geq W$, giving (A).

If (A) fails, every outside u has $|I(u)| \leq y\ell$. Counting incidences between outside vertices and blocks shows that some block receives at most $y|G|$ incidences, proving (B). $\square$

Part II

The fixed-target structural implication

7 One ambient class and a quantified rooted contract

Theorem 7.1 (Fixed-target implication). *Fix $s \geq 7$. Assume $\text{EH}(P_{s-1})$ and the strong-wonderfulness statement W_s of Theorem 6.1 for $\overline{P_s}$ -free graphs, with some fixed exponent $a_s \geq 6$. Then $\text{EH}(P_s)$ holds.*

Set $\mathcal{A}_s = \text{Forb}(P_s)$, in the F orientation, and write $G = \overline{F}$. Let E_r be the hereditary class of graphs with an order in which the first vertex of every induced P_r is not an endpoint.

Write $T_r^{(s)}$ for the following quantified rooted Tooth statement. There are positive constants $A_r, e_r, c_r, k_r, d_r, E_r^*, \eta_r$ such that, whenever $0 < x \leq y \leq \eta_r$, $|Y| = n \geq x^{-E_r^*}$, and U is outside Y in a host F in $\mathcal{A}_s$ with every u having RP_r into the original Y , at least one of the following exists: (T_1) a core $S \subseteq Y$ with $|S| \geq y^{A_r}n$, such that every $u \in U$ is complete to S in G or has strictly fewer than $x|S|/4$ G -neighbours in S ; (T_2) a z^4 -restricted set $T \subseteq Y$ of size at least $z^{e_r}n$ for some $x^{c_r} \leq z \leq y$; (T_3) a directed x -semisparse blockade in Y of integer length K with $1/y \leq K \leq x^{-k_r}$ and width at least n/K^{d_r} .

The entire host, including original roots, stays P_s -free in the F orientation. Each lower Tooth call takes an induced subhost. The superscript s remains the same throughout the induction.

The already proved, ambient-independent RP_5 Tooth supplies $T_5^{(s)}$ for every s , with $A_5 = 26, e_5 = 18, c_5 = 1, k_5 = 140, d_5 = 64, E_5^* = 600, \eta_5 = 2^{-16}$. Its fixed y^4 -restricted output is a valid T_2 output at $z = y$. This base theorem does not use $\text{EH}(P_s), \text{EH}(P_{s-1})$, or $\text{EH}(E_5)$. The RP_5 contract is developed in Appendix C, with its general Tooth proof in Appendix A and its thin-layer argument in Appendix B.

8 Ordered-tail transfer without circularity

Assume $T_r^{(s)}$ has been proved. The transfer argument in Appendix J first proves generalized niceness for $C_{r,s} = \text{co}(E_r \cap \mathcal{A}_s)$.

The structural step is direct. In a y^3 -sparse G in this class, take the last $\lceil yn \rceil$ vertices of an E_r order as the target Y . Every earlier vertex has RP_r into Y . Applying $T_r^{(s)}$ gives the variable-scale local oracle of Hypothesis H.1 for the hereditary class $\mathcal{C} = C_{r,s}$ and $h = s$,

with constants $a = \max(4, A_r + 1)$, $b = 3$, $c = \max(2, c_r, k_r)$, $d = d_r + 1$, $e = \max(4, e_r + 1)$, $L = E_r^* + 2$. The clean outcome has a large directed sparse residual because at most $y^3 n$ earlier roots can be complete to its nonempty core. The restricted outcome retains its actual z and has width $z^{e_r} |Y| \geq z^{e_r+1} n$. The blockade width is at least n/K^{d_r+1} ; this blockade is directed semisparsely, hence weakly semisparsely as output (O_2) requires.

The variable-scale bootstrap (Theorem H.2) and the arbitrary-pattern layout (Lemma I.1) prove niceness for $C_{r,s}$ from this oracle, without using either (EH) or (W). Then (EH), via normalized P_{s-1} virality, supplies the sparse P_s -free leaf-fibre step. Together with (W) and the proved terminal product argument, this niceness gives $\text{EH}(E_r \cap \mathcal{A}_s)$.

Thus $T_r^{(s)} + (\text{EH}) + (W) \implies \text{EH}(E_r \cap \mathcal{A}_s)$. No assertion of $\text{EH}(P_s)$ occurs on the right-hand side of an input. No virality theorem for the possibly infinitely defined class E_r is invoked: only virality of the single graph P_{s-1} is used.

9 Induction through rooted orders

If $s = 7$, $T_5^{(s)}$ is already the final rooted order needed.

For $s \geq 8$, apply Section 8 to $r = 5$. It gives $\text{EH}(E_5 \cap \mathcal{A}_s)$. The fixed-ambient variant in Appendix F.7 then gives $T_6^{(s)}$. Its use of the original RP_4 Tooth is included in the specialized frontier proof; it does not require a nonexistent uniform RP_4 contract.

For every subsequent $q = 7, \dots, s - 2$, assume $T_{q-1}^{(s)}$ and $T_{q-2}^{(s)}$ already proved. Section 8, applied to $r = q - 1$, gives $\text{EH}(E_{q-1} \cap \mathcal{A}_s)$, with some positive exponent. Apply the recurrence of Appendix G.6 with this fixed hereditary ambient class $\mathcal{A}_s$. Its two preparatory purifications use $T_{q-1}^{(s)}$; its offending-pair purification uses $T_{q-2}^{(s)}$. Its negative transversals belong to $E_{q-1} \cap \mathcal{A}_s$ and have the required EH exponent.

For clarity, that recurrence provides explicit finite next constants. Enlarge the two lower contracts to common constants $A_0, e_0, d_0, c_0, k_0, E_0 \geq 1$ and common cutoff η_0 . Set $J_0 = 4A_0 + 11$, $H = c_0 J_0$, $J = k_0 J_0$, $Q = E_0 J_0$, $m_0 = \max(2^{16}, \lceil 1/\eta_0 \rceil)$, $h = \max(2, 1/\kappa)$, $d_t = \lceil 4h + 5 \rceil$, $C_{\text{tree}} = 6d_t + 3$. One may take $A_q = 6d_t + 2$, $c_q = H(C_{\text{tree}} + 1)$, $e_q = \max(e_0 + 2A_0 + 7, 8h + d_t + 10)$, $k_q = J(C_{\text{tree}} + 1)$, $d_q = 9A_0 + d_0 + 21$, $E_q^* = Q + (Q + 6)C_{\text{tree}} + 1$, $\eta_q = \min(2^{-16}, 2^{-(h+10)}, 1/m_0)$, where κ is the just-proved EH exponent of $E_{q-1} \cap \mathcal{A}_s$. These are the constants (G.1), (G.4) and (G.5) of Appendix G; the letter h has this local meaning only here and there.

Every representative and every witness is an original host vertex. The prefix r - z before a representative creates the rooted P_{q-2} -exclusion needed for the third pass. The threshold strictly below one preserves original full/sparse types after shrinking. All restricted early outputs keep their actual z ; all order and width losses are absorbed into the displayed finite exponents.

This finite induction proves $T_{s-2}^{(s)}$. No theorem established only in a smaller ambient forbidden-path class has been reused.

10 The final comb oracle

Write the final $T_{s-2}^{(s)}$ constants as A, e, c, k, d, E, η . Let $F = \overline{G}$ be P_s -free, equivalently let G be $\overline{P}_s$ -free, and suppose G is y^3 -sparse, with $0 < \rho \leq y \leq \min(\eta, 2^{-8})$, $|G| = n \geq \rho^{-(3E+10)}$. Apply Contract 2.7, namely [4, Lemma 2.10], with $H = G$. Its order hypothesis holds since $n \geq \rho^{-(3E+10)} \geq \rho^{-4} \geq y^{-4}$. It gives one of: (O_1) G is $2y^4$ -sparse; (O_3) disjoint X, Z with $|X| \geq y^4 n$, $|Z| \geq (1 - 4y)n$, and Z directed ρ -sparse to X ; or a comb of integer length $1/y \leq \ell \leq \rho^{-2}$, teeth B_i of size at least $y^4 n / \ell^2$, roots a_i full to their own tooth and anticomplete to

all other teeth, and a special vertex v full to all teeth and anticomplete to all roots. The sets, roots and v are disjoint.

Every vertex u of a different original tooth has RP_{s-2} into B_i . Otherwise a rooted induced $F-P_{s-2}$ from u inside B_i , prefixed by v, a_i , would be an induced $F-P_s$. The two new consecutive F -edges are $v-a_i$ and a_i-u . All new nonconsecutive pairs are absent in F , since v is G -full to every tooth and a_i is G -full to B_i .

Apply $T_{s-2}^{(s)}$ to each tooth, with all vertices of the other original teeth as roots, at fine ρ^3 and coarse $1/\ell$. These parameters and the input orders are legal: $\rho^3 \leq \rho^2 \leq 1/\ell \leq y \leq \eta$, $|B_i| \geq \rho^8 n \geq \rho^{-3E}$. For width estimates use the separate inequality $|B_i| \geq n/\ell^6$.

A restricted output retains z in $[\rho^{3c}, 1/\ell]$ and has size $z^e |B_i| \geq z^{e+6} n$. A blockade output is directed ρ^3 -semisparse, hence directed ρ -semisparse; it has $1/y \leq \ell \leq K \leq \rho^{-3k}$ and width at least n/K^{d+6} .

If all teeth give clean cores, their sizes are at least n/ℓ^{A+6} , and every vertex in either core is full or θ -sparse toward the other, $\theta = \rho^3/4$. For an incomplete pair choose a nonedge pq , with p in the first core and q in the second. Both endpoints have sparse type. Any vertex of the first core full to the second is adjacent to q ; hence such vertices form fewer than $\theta \cdot$ the first core size. Counting them and the remaining sparse vertices gives edge density less than $2\theta \leq \rho$. Thus the clean cores themselves give a weakly ρ -semisparse blockade of length ℓ and width at least n/ℓ^{A+6} . It need not be directed ρ -semisparse: the edge count does not bound the degrees of the few full-type vertices.

The final local oracle therefore has constants $a_{\text{or}} = 4, b_{\text{or}} = 4, c_{\text{or}} = \max(2, 3c, 3k), d_{\text{or}} = \max(d+6, A+6), e_{\text{or}} = \max(4, e+6), L_{\text{or}} = 3E+10$. This is Hypothesis H.1 for the hereditary class $\mathcal{C} = \text{Forb}(\overline{P_s})$ and $h = s$: the comb extraction and $T_{s-2}^{(s)}$ apply in every $\overline{P_s}$ -free graph, so the same outputs are available in every member of $\mathcal{C}$. Its blockade output is (O_2) in the weak sense of that hypothesis. A (T_3) blockade is directed, hence weakly, ρ -semisparse, while the all-clean branch supplies weak ρ -semisparsity only.

11 Generalized niceness and the terminal conclusion

Apply Theorem H.2 with $\mathcal{C} = \text{Forb}(\overline{P_s})$ and $h = s$. Its minimum admissible scale ranges down to $\rho^{2c_{\text{or}}}$, so an actual restricted scale z as small as $\rho^{c_{\text{or}}}$ causes no loss of validity. In the sparse direction the new scale is $z^{7/6}$; in the opposite direction Contract 2.6 for P_s , with parameter z^2 , gives a complete blockade; its order hypothesis is verified in the proof of Theorem H.2. Fixed Rödl and the proved layout argument supply constants M, N with, for every $\overline{P_s}$ -free G and every $0 < \varepsilon < 1/2$, either $\text{hom}(G) \geq \varepsilon^M |G|$ or a blockade of length $\lceil 1/\varepsilon \rceil$, width $\varepsilon^N |G|$, whose pairs are complete or weakly ε^8 -sparse.

The terminal argument in Appendix K uses only hereditary niceness, $\overline{P_s}$ -freeness, $\text{EH}(P_{s-1})$, and W_s . It applies to the whole $\overline{P_s}$ -free class as well as to the relative class $C_{r,s}$.

More explicitly, normalize P_{s-1} virality with exponent $v \geq 1$ and put $P = \max(M, N+3)$, $A_{\text{term}} = \max(a_s P, 4(s-1)v+4, 4), c_T = y_T = 2^{-4s}, d_T = 2A_{\text{term}}+4$. The sparse P_s -free leaf-fibre step has residual loss at most $(s-1)y$. In the other orientation, weak-blockade cleanup followed by (W) gives the required restricted output or a large pure residual pair. The fixed-orientation residual chain survives a constant fraction, since $\exp(-2(s-1)) > c_T$.

Let δ_T be the fixed Rödl fraction for $\overline{P_s}$ at precision $c_T y_T$, and put $D_s = d_T + \lceil \log_2(1/\delta_T) \rceil$. These are the constants (K.1)–(K.2) of Appendix K, where $A_{\text{term}}, c_T, y_T, d_T$ are abbreviated to A, c, y_0, d . Lemma K.3 yields $\text{hom}(G) \geq |G|^{1/(8D_s)}$ for every $\overline{P_s}$ -free graph G . Homogeneous size is invariant under complementation, so the same bound holds for every P_s -free graph F . This is the Erdős–Hajnal property of P_s .

12 The induction on path length

Proof of Theorem 1.1. Theorem A of [1] supplies $\text{EH}(P_6)$. For a fixed $s \geq 7$, assume $\text{EH}(P_{s-1})$. Theorem 6.1, with $h = s - 1$, supplies W_s , and Theorem 7.1 supplies $\text{EH}(P_s)$. Induction therefore gives $\text{EH}(P_s)$ for every fixed $s \geq 6$. For $2 \leq s < 6$, every P_s -free graph is P_6 -free, since an induced P_6 contains an induced P_s . For $s = 1$, no nonempty graph is P_1 -free, so the assertion is vacuous. Complementation interchanges cliques and stable sets and preserves hom, giving the complementary conclusion. $\square$

The exponent obtained in this way depends on s ; no common exponent for all path lengths is asserted.

A The general Tooth lemma and modular decomposition

Lemma A.1 (General Tooth with actual integer lengths). *Let $0 < \xi \leq 2^{-20}$, let k be an integer with $2^{16} \leq k \leq 2\xi^{-1/2}$, and let $|Y| = V \geq 16\xi^{-3}$. Suppose U is outside Y and, for every u in U , each anticomponent of $G[N(u) \cap Y]$ is a module of $G[Y]$. Then at least one of the following exists:*

- TL1: a pure blockade of integer length K with $k^{1/4} \leq K \leq 1/\xi$ and width at least V/K^4 ;*
- TL2: a pure blockade of integer length K with $(8K)^4 \leq k^3 \leq (9K)^4$ and width at least $V/(2k)$;*
- TL3: a complete or anticomplete blockade of integer length $K = \lfloor 1/\xi \rfloor$ and width at least $\xi^2 V/k$;*
- TL4: a complete blockade of integer length $K = \lfloor 1/(\xi k) \rfloor$ and width at least $\xi^3 V/4$;*
- TL5: a set $S \subseteq Y$ with $|S| \geq V/k$ such that every $u \in U$ is complete to S or has fewer than $\xi|S|/4$ neighbours in S .*

Proof. We use the standard modular-decomposition facts in their precise form. Strong modules form a rooted inclusion tree whose children partition each non-singleton node. Distinct disjoint modules are complete or anticomplete to one another. At an internal node the quotient on its children is complete, edgeless, or prime. In the first two cases its children are respectively anticomponents or components. At a prime node, every proper module contained in the node lies in a single child. An arbitrary module and a strong module are disjoint or nested. All these assertions concern modular decomposition of the given graph; none assumes a forbidden P_6 or P_7 .

The hypotheses force $\xi \leq 2^{-30}$, since $2^{16} \leq k \leq 2\xi^{-1/2}$; every application in this paper has $\xi \leq 2^{-32}$. Use the notation ξ, k, Y, V of the general Tooth contract and write $n = V$. The symbol s_{cut} denotes a local size cutoff here and in Appendices C, F and G; it is unrelated to the target path order s . Set $s_{\text{cut}} = \xi^2 n/4$, $t = n/k$, $K_1 = \lfloor 1/\xi \rfloor$. The parameter range gives $k^2 \xi \leq 4$ and $\xi k \leq 2\xi^{1/2} \leq 2^{-9}$. In particular $s_{\text{cut}} \geq 4/\xi > 1$ and $s_{\text{cut}} \leq t$.

Let L be the strong modules of $G[Y]$ having at least s_{cut} vertices. An atom is a child of a member of L that is not itself in L . For N in L let $S(N)$ be the union of its atom children. The sets $S(N)$ partition Y : for each vertex, its lowest ancestor in L has an atom child containing it. Distinct such atom layers cannot overlap by the tree nesting.

We need two additional consequences of the standard facts. First, each anticomponent A of $N_G(u) \cap Y$ is a module by hypothesis. For a strong node N , either A contains N , misses N , or is properly contained in N ; in the last case, if N is a prime node, A lies in a single child of N . Second, for nested retained nodes N properly containing N' , $S(N)$ misses N' , and every vertex outside N' is complete or anticomplete to N' , because N' is a module.

A.1 An elementary antichain estimate

Suppose $2 \leq K_0 \leq K_1$, and $\mathcal{F}$ is a family of sets, each of size at most M . Suppose that for every integer $K \in [K_0, K_1]$ fewer than K members of $\mathcal{F}$ have size at least n/K^4 . Then

$$\sum_{A \in \mathcal{F}} |A| \leq (K_0 - 1)M + \frac{n}{K_0 - 1} + \frac{|\mathcal{F}|n}{K_1^4}. \quad (\text{A.1})$$

Indeed, fewer than K_0 sets reach n/K_0^4 . For each $K_0 \leq K < K_1$, at most K sets have size in $[n/(K+1)^4, n/K^4]$, contributing at most n/K^3 . The remaining sets each contribute less than n/K_1^4 . Finally $\sum_{K \geq K_0} K^{-3} \leq \sum_{K \geq K_0} (1/(K-1) - 1/K) = 1/(K_0 - 1)$.

Let K_0 be the least integer with $K_0^4 \geq k$ and put $a = K_0 - 1$. Then $a \geq 15, a^4 < k$, and $K_0 \leq K_1$. To check the last assertion, $a^8 \xi < k^2 \xi \leq 4$ and $a^7 \geq 8$ imply $8a\xi < 4$; hence $(a+1)\xi \leq 2a\xi < 1$.

A.2 Find a heavy chain

Call nodes of L of size at least t heavy. Minimal heavy nodes are pairwise disjoint modules. If at least K_0 exist, they form a pure $(K_0, n/K_0^4)$ -blockade, giving TL1. Otherwise there are at most a minimal heavy nodes.

The maximal nonheavy nodes have sizes in $[s_{\text{cut}}, t)$ and there are at most n/s_{cut} of them. If K of them have size at least n/K^4 for some $K_0 \leq K \leq K_1$, they give TL1. Otherwise (A.1) yields total size at most $an/k + n/a + (n/s_{\text{cut}})n/K_1^4 \leq n/a^3 + n/15 + 64\xi^2 n \leq (1/3375 + 1/15 + 2^{-34})n \leq n/4$. All nonheavy layers lie in these maximal nonheavy nodes. Thus the heavy-node layers have total size at least $3n/4$. Every heavy node contains a minimal heavy node. Summing layer masses on ancestor chains above the at most a minimal heavy nodes, some chain C has total layer mass at least $3n/(4a)$.

A.3 A thick layer gives an early output or the clean core

Suppose a node N on C has $|S(N)| \geq t$; put $S = S(N)$. If $G[N]$ is disconnected, the components of $G[S]$ are atom children, each smaller than s_{cut} . Since $s_{\text{cut}} \leq \xi t \leq \xi|S| \leq |S|/K_1$, grouping components gives an anticomplete length- K_1 blockade of width at least $|S|/K_1^2 \geq \xi^2 n/k$, giving TL3. If the complement of $G[N]$ is disconnected, the same argument gives a complete blockade and again TL3.

The remaining case is a prime modular-decomposition node. Put $K' = \lfloor 1/(\xi k) \rfloor \geq 2$. Suppose some u in U is not complete to S but has $r \geq \xi|S|/4$ neighbours in S . Every anticomponent A of $N_G(u) \cap Y$ that meets S is properly contained in N : otherwise it would contain N and u would be complete to S . Since N is prime, A lies in one child. Because it meets S , that child is an atom, so A is contained in S and has size less than s_{cut} . Consequently the anticomponents of $N_G(u) \cap S$ all have size less than s_{cut} . Moreover $r/K' \geq r\xi k \geq \xi t \xi k/4 = s_{\text{cut}}$. Grouping these anticomponents gives a complete length- K' blockade with width $r/K'^2 \geq \xi t (\xi k)^2/4 \geq \xi^3 n/4$, which is TL4. If no such u exists, S itself satisfies TL5 and $|S| \geq t = n/k$.

A.4 All chain layers are thin

Assume instead $|S(N)| < t$ for every N on C . Let K be the largest integer satisfying $(8K)^4 \leq k^3$. Then $K \geq 8$, and $k^3 < (8K+8)^4 \leq (9K)^4$. Also $(8Ka)^4 \leq k^3 a^4 < k^4$, so $8Ka < k$; in particular $2Kt \leq 3n/(4a)$, no larger than the chain's total layer mass.

For N on C , let P_N be cumulative layer mass from the top through N . For $j = 0, \dots, K-1$ let Γ_j be the union of layers with $2jt < P_N \leq 2(j+1)t$. Since each layer has size less than t , the cumulative mass at any threshold θ lies in $(\theta - t, \theta]$. Therefore every Γ_j has more than t vertices.

Every vertex v outside a chain node is complete or anticomplete to that node. Its choice is the same for every chain node omitting it: if v is anticomplete to one such node, nesting and the module property force it to be anticomplete to all of them. Call v type 1 if it is complete to all those nodes and type 0 otherwise. For $j < j'$, a vertex in Γ_j sees a vertex in $\Gamma_{j'}$ precisely when the earlier vertex has type 1. Retain a majority type within each Γ_j . The retained sets each have size at least $t/2 = n/(2k)$ and form a pure blockade of actual length

K with $(8K)^4 \leq k^3 \leq (9K)^4$. This is TL2.

All possibilities have yielded TL1–TL5. This proves the general Tooth lemma, including its actual integer length contracts. $\square$

B Thin layers with perfect partial transversals

Lemma B.1. *Suppose a set of size M is partitioned into nonempty layers A_i , every partial transversal induces a perfect graph, and*

$$\lambda = \frac{\max_i |A_i|}{M} \leq \frac{\varepsilon^4}{128}, \quad 0 < \varepsilon \leq 2^{-16}.$$

Then it contains a set T of size at least $\varepsilon^2 M/512$ such that $G[T]$ or its complement has maximum degree at most $\varepsilon(|T| - 1)$. In particular T is ε -restricted.

Proof. Put $q = \lceil \varepsilon^{-2} \rceil$, $t = q^2$, and $\delta = 1/(256q)$. Since $q \leq 2\varepsilon^{-2}$, we have $\lambda \leq 1/(32q^2)$; since $\lambda \geq 1/M$, we have $M \geq 32q^2$.

A uniform $2t$ -subset has expected number of colliding same-layer pairs less than $2t^2\lambda \leq t/16$. At least half of these samples have at most t collisions. Deleting at most one vertex per collision leaves at least t vertices in distinct layers. A perfect graph on $t = q^2$ vertices has $\alpha\omega \geq t$, hence has a homogeneous q -set. Double counting gives at least

$$\frac{1}{2} \frac{\binom{M}{2t}}{\binom{M-q}{2t-q}} \geq \frac{1}{2} \binom{M}{2t}^q$$

homogeneous q -subsets. One type contributes at least $\frac{1}{4}(M/(2t))^q$; suppose it is the stable-set type.

Assume no set of size at least δM is ε -restricted, even with the stronger degree convention $\varepsilon(|S| - 1)$. Put

$$u = \lceil \delta M \rceil \leq 2\delta M, \quad r = \lceil \varepsilon^{-1} \ln(1/\delta) \rceil.$$

Every set R with $|R| \geq u$ has maximum degree greater than $\varepsilon(|R| - 1)$.

The required elementary container count is

$$I_q \leq \binom{M}{r} \binom{u}{q-r}.$$

Fix tie-breaking orders. To encode a stable q -set I , repeatedly inspect a vertex of maximum degree in the graph induced on the current candidate set R . If it belongs to I , record it and delete its closed neighbourhood; otherwise delete that vertex alone. Stop when at most u candidates remain. While $|R| > u$, a recorded choice deletes more than $\varepsilon(|R| - 1) + 1$ vertices, hence leaves fewer than $(1 - \varepsilon)|R|$ candidates; so at most r choices are recorded, because $(1 - \varepsilon)^r M \leq \delta M \leq u$. Pad the recorded set to size r with remaining members of I if necessary. From this r -set the deterministic algorithm can be replayed, and the other $q - r$ members lie in a container of size at most u . This proves the count once $r \leq q$, as verified below. If $u < q - r$, the upper bound is zero.

Using $\binom{q}{r} \leq 2^q$ and $q! \geq (q/e)^q$, comparison gives

$$\begin{aligned} \frac{1}{4} \binom{M}{2t}^q &\leq \frac{M^q (2\delta)^{q-r}}{r!(q-r)!} \\ &\leq M^q (2\delta)^{q-r} \left(\frac{2e}{q} \right)^q. \end{aligned}$$

Thus

$$\left(\frac{1}{2\delta}\right)^{1-r/q} \leq 4e \frac{t}{q} 4^{1/q} \leq 8eq.$$

Write $L = \ln(1/\varepsilon) \geq 16 \ln 2$. Then $\ln(256q) \leq 3L$ and $r \leq 3\varepsilon^{-1}L + 1$, so

$$\frac{r}{q} \ln(128q) \leq 9\varepsilon L^2 + 3\varepsilon^2 L < 1, \quad r < q.$$

The preceding inequality implies

$$128q = \frac{1}{2\delta} < 8e^2 q < 128q,$$

a contradiction. For the other homogeneous type, use the complement. Finally $\delta \geq \varepsilon^2/512$, proving the lemma. $\square$

C The ambient-independent two-parameter RP5 Tooth

The proof uses the general Tooth lemma of Appendix A and the perfect-transversal sampling lemma of Appendix B.

Theorem C.1 (Two-parameter RP5 Tooth). *Let $0 < x \leq y \leq 2^{-16}$, let $|Y| = n \geq x^{-600}$, and let U be disjoint from Y . Put $F = \overline{G}$. Assume that $F[Y \cup \{u\}]$ has no induced P_5 with endpoint u , for each $u \in U$. Then one of the following holds.*

- (A) *A set $S \subseteq Y$ has size at least $y^{26}n$, and every $u \in U$ is G -complete to S or has fewer than $x|S|/4$ neighbours in S .*
- (B) *A y^4 -restricted set $T \subseteq Y$ has size at least $y^{18}n$. Here restricted is meant with the stronger degree bound $y^4(|T| - 1)$.*
- (C) *$G[Y]$ has a blockade of integer length K with $1/y \leq K \leq x^{-140}$ and width at least n/K^{64} , each pair being complete or directed x -sparse from the later block to the earlier one.*

C.1 Local contracts and rooted module equivalence

Proof. The original Tooth lemma is used at parameters (ξ, k) , with $\xi \leq 2^{-20}$, $2^{16} \leq k \leq 2\xi^{-1/2}$, and local order $V \geq 16\xi^{-3}$. For an outside root set whose roots have no induced F - P_4 into the local set, it returns either a core of size at least V/k on which every root is complete or has fewer than a $\xi/4$ fraction of neighbours, simultaneously for all roots, or an early pure, complete or anticomplete blockade: TL1: $k^{1/4} \leq K \leq 1/\xi$, width V/K^4 ;

TL2: $(8K)^4 \leq k^3 \leq (9K)^4$, width $V/(2k)$;

TL3: $K = \lfloor 1/\xi \rfloor$, width $\xi^2 V/k$;

TL4: $K = \lfloor 1/(\xi k) \rfloor$, width $\xi^3 V/4$.

This exact contract is proved in Appendix A. Here is the exact rooted- P_4 equivalence used to apply it. For u outside Y , let C be an F -component of $N_G(u) \cap Y$. If C is not a module of $G[Y]$, a vertex p outside C mixes on C . It cannot be in another F -component of $N_G(u) \cap Y$, so up is an F -edge. Connectivity supplies an F -edge qr in C with pq an F -edge and pr a nonedge. Then $u-p-q-r$ is an induced F - P_4 . Conversely, in an induced F - P_4 $u-p-q-r$, the vertices q, r belong to one such component, and p mixes on it. Thus the module condition is equivalent to the absence of a rooted F - P_4 , and is inherited whenever the target set is restricted.

The thin-layer sampling contract is: if sets partition a set of size M , every partial transversal induces a perfect graph, max layer size/ $M \leq \varepsilon^4/128$, and $0 < \varepsilon \leq 2^{-16}$, then there is an ε -restricted subset of size at least $\varepsilon^2 M/512$, with the stronger degree bound $\varepsilon(|T| - 1)$. It was proved in Appendix B. Both local contracts are used exactly as stated.

C.2 An ordered frontier has the required one-sided root law

At a node S choose u in U . Its ordered children are all nonempty F -components $C_1, \dots, C_d$ of $N_G(u) \cap S$, in any fixed order, followed by $A = S$ minus $N_G(u)$ if nonempty. They partition S . Take an antichain $Z_1, \dots, Z_m$ of tree nodes, listed in child order.

For $i < j$ and v in Z_j , there is no induced F - P_4 with endpoint v and the other three vertices in Z_i . At their lowest common ancestor there are two cases. Different C children give a G -complete pair, so v has no F -neighbour in Z_i and cannot begin that path. Otherwise Z_i is in a C child and Z_j in A . The ancestor's root u is G -complete to Z_i and anticomplete to v . Prefixing u to a rooted P_4 v - p - q - r gives the induced F - P_5 u - v - p - q - r , a contradiction. This argument uses only containment at the lowest common ancestor. It makes no claim that an A child is endpoint-closed in the original Y .

For each $j < l$ separated as C versus A , record that ancestor root in R_j . Thus $|R_j| \leq m - 1$, every root in R_j is complete to Z_j , and for this particular pair some root $r(j, l)$ in R_j is anticomplete to Z_l . Pairs separated as distinct C children are complete and need no root. All recorded roots remain outside Y and retain the original RP_5 law.

C.3 Simultaneous root-family purification

Suppose $B_1, \dots, B_m$ have the property that every later vertex is complete or θ -sparse to an earlier block. Retain the same families R_j , whose members are complete to B_j . Set $\xi = m^{-2}$, $k = m$, $\eta = 1/(16m^3)$, $\theta \leq \tau/(256m^5)$. Call $i < j$ good if some r in R_j has at least $\eta|B_i|$ nonneighbours in B_i ; otherwise call it red. Goodness has an existential quantifier; redness has the corresponding universal quantifier over R_j .

For each good pair choose its witnessing root r and a nonneighbour p_{ij} in B_i with $\deg_G(p_{ij}, S_{ij}) \leq \theta|B_j|/\eta$, where S_{ij} is the sparse-type part of B_j . This follows by averaging: $e(B_i, S_{ij}) \leq \theta|B_i||S_{ij}|$. The root r is complete to B_j , so a rooted F - P_4 from p_{ij} into B_j would extend with r to an RP_5 . Thus Tooth applies to B_j with the at most m representatives p_{ij} as its outside root set.

Unless it returns an early output, Tooth gives $D_j \geq |B_j|/m$. For each good pair: if p_{ij} is complete to D_j , remove $D_j \cap S_{ij}$; if p_{ij} has fewer than $\xi|D_j|/4$ neighbours in D_j , remove the complete-type part of D_j . The respective deletion fractions are at most $\theta m/\eta$ and less than $\xi/4$. All deletions are unioned and measured relative to the original D_j . They cost at most $m(\xi/4 + \theta m/\eta) \leq 1/(4m) + \tau/16 < 1/8$.

For each red $i < j$, delete from D_i every nonneighbour of every root in R_j . There are at most m^2 root occurrences affecting B_i , each costing less than $\eta|B_i|$, so the extra cost is at most $m^2\eta|B_i| = |B_i|/(16m) \leq |D_i|/16$. The resulting E_i have size at least $|B_i|/(2m)$. Good pairs have uniform original types; on red pairs every r in R_j is exactly complete to E_i . Representatives need not survive in any final block. Different good pairs may use different roots, without repeated intersections of cores.

C.4 The interval argument and the forcing path

Choose an F -component C_i of E_i with size at least $|E_i|/m$. If this is impossible in some E_i , the component grouping lemma instead gives a complete length- m blockade of width at least $|E_i|/m^2$; this is an early output. Otherwise C_i is F -connected and has size at least $|B_i|/(2m^2)$. Sparse original types now have threshold $\beta = 2m^2\theta < 1/2$. They cannot become complete, so each ambiguous pair (both types present in its later block) was red. Let Q be the graph of these ambiguous pairs, with each edge directed toward the later index.

Claim: two vertices of any increasing Q -path cannot lie in different C children of one tree node. The indices in every child subtree are an interval, as are the indices in the parent subtree. Between two such indices the path remains inside the parent interval, and before the later C child it cannot visit the A interval, which is last. Therefore one consecutive path edge must cross two C -child intervals. Such a pair is complete, contradicting its being a Q -edge.

Consequently any two path indices $j < l$ have a separating root $r(j, l)$ in R_j : their lowest common ancestor separates them as C versus A . For an edge $i \rightarrow j$ of that path and a still later path block C_l , red exactification says this particular root is complete to C_i . It is already complete to C_j and anticomplete to C_l .

If z in C_l were complete to C_j but sparse to C_i , F -connectedness of C_j supplies an F -edge $v-w$ crossing its two types, with v sparse to C_i and w complete. Since $2\beta < 1$, choose b in C_i nonadjacent in G to both z and v . Then $r(j, l)$ - z - b - v - w is an induced F - P_5 : consecutive edges follow from the separating root, the choice of b , and the crossing edge $v-w$; the six nonconsecutive pairs $r-b, r-v, r-w, z-v, z-w, b-w$ are G -edges. This contradicts RP_5 of $r(j, l)$. Thus each later vertex's type vector along the earlier path is a prefix of complete types followed by sparse types.

Crucial quantifier point: $r(j, l)$ may depend on both indices. Every such root was made complete to C_i because the red deletion used all members of R_j . No common root for a path is required. An empty R_j is harmless: whenever this argument needs a root at j , a later path index l itself proves R_j nonempty.

Put $r = \lfloor \sqrt{m} \rfloor$. If an increasing Q -path has r vertices, select r of them. In its j -th block there are at most $j \leq r$ prefix profiles. Keep a largest profile cell. The resulting blockade has width at least $\min |C_i|/r$ and later-to-earlier precision at most $r\beta$. Profiles are measured relative to the original connected C_i , so the independent choices of cells have no circular dependence. If no such path exists, the increasing-path height partition has at most $r - 1$ levels. Every level is Q -independent and one has at least $m/(r - 1) \geq r$ indices. Those block pairs already have uniform types.

C.5 Quantitative frontier lemma

For a frontier of $m \geq 2^{16}$ blocks Z_i of size at least W , take $\xi_0 = \tau/(64m^5)$, $k_0 = m$, $W \geq 16\xi_0^{-3}$. Use Tooth on each Z_i against the union of all later original Z_j ; Appendix C.2 gives its input. Without an early output this gives $|B_i| \geq |Z_i|/m$, $\theta = \xi_0/4 = \tau/(256m^5)$. The second Tooth in Appendix C.3 is permitted since $|B_i| \geq 16m^6$. Appendices C.3–C.4 now give either component grouping or a length- r complete-or-directed- τ -sparse blockade with width at least $W/(2m^3r)$.

If $|Z_i| \geq n/m^b$, every result has actual integer length $m^{1/4} \leq K \leq 64m^5/\tau$ and width at least n/K^{4b+12} . For completeness, the early conversions are: first Tooth TL1: exponent $4b+4$; first TL2: exponent $2b+3$, since $K \geq \sqrt{m}$; first TL3: exponent $b+5$, using $\xi_0 \geq 1/(2K)$, $K \geq m$; first TL4: exponent $b+11$, using $\xi_0 \geq 1/(2mK)$, $K \geq m$; second Tooth: exponent at most $4b+8$; component output $K = m$: exponent $b+5$; path/level output $K = r$: exponent $3b+11$, using $m \leq r^3$ and $r \geq 256$. All these are at most $4b+12$. The largest early length is $1/\xi_0$. This is where actual TL3/TL4 lengths are essential.

The same lemma applies to positive layers with a root complete to its own layer and anticomplete to all later layers. Those roots directly give both the one-sided rooted- P_4 law and all separating roots, so the tree interval argument is unnecessary for that application.

The elementary component grouping used above has no rounding defect: if atoms have size less than M/k , greedily form groups of size at least M/k^2 . Each group uses less than $(k+1)M/k^2$ vertices; after $k-1$ groups, more than M/k^2 remain. Hence k groups exist and

are pairwise complete.

C.6 Full tree and its mass accounting

Set $t = \lceil y^{-4} \rceil$ and $s_{\text{cut}} = n/t^6$. Split every node of size at least s_{cut} unless it already satisfies the complete-or-sparse condition of (A). A bad root has $x|S|/4 \leq \deg_G(u, S) < |S|$, so both its A and neighbour parts are nonempty, and every child is proper. The tree is finite. A clean node has size at least $s_{\text{cut}} \geq y^{24}n/64 \geq y^{26}n$, giving A. Assume none exists. Small children, of size less than s_{cut} , form disjoint atoms partitioning Y . Retain all nodes of size at least s_{cut} .

If t retained nodes form an antichain, Appendix C.5 applies with $m = t, \tau = x, b = 6$ and gives C . Otherwise the retained tree has $q < t$ leaves and fewer than t branching nodes, hence fewer than $2t$ exceptional nodes when terminal nodes are included.

At any node, its small C children are pairwise complete. If their union D has size at least ts_{cut} , component grouping gives t complete blocks of width at least $|D|/t^2 \geq n/t^7$, which gives C . Otherwise the atom mass at that node is less than $(t+1)s_{\text{cut}}$, including its possible small A child. Thus exceptional nodes carry atom mass less than $2t(t+1)s_{\text{cut}}$. Unary nodes carry at least $n - 2t(t+1)s_{\text{cut}}$.

The q complete root-to-leaf paths cover all unary nodes. Summing their unary atom masses counts each unary node at least once, so one full path has unary atom mass at least $n/t - 2(t+1)s_{\text{cut}}$. Branching nodes may occur on this path; their atoms are simply omitted. The argument does not require a long uninterrupted unary segment.

At a unary node, exactly one child is retained. If it is A, the atom layer is the entire neighbour set P . It has $xs_{\text{cut}}/4 \leq |P| < ts_{\text{cut}}$. Its root is complete to P and anticomplete to the entire continuation, hence to all later positive layers. If it is a C child, record its A part as a negative layer of size less than s_{cut} . Its root is anticomplete to that layer and complete to the entire continuation. The remaining atoms are small C siblings.

All small C siblings at negative steps are pairwise complete, also across different depths: the earlier sibling is complete to the selected C child containing all later nodes. If their total mass is at least ts_{cut} , grouping again yields C . Otherwise discard less than ts_{cut} of fringe. The remaining positive and negative layers have total mass at least $n/t - (3t+2)s_{\text{cut}} \geq 3n/(4t)$. One of the two directions has mass at least $3n/(8t)$.

C.7 Finish both directions

Negative direction. Its mass is $M \geq 3n/(8t)$, each layer has size less than s_{cut} , and hence maximum layer fraction $\lambda \leq 8/(3t^5) \leq 8y^{20}/3 \leq y^{16}/128$. Every partial transversal induces a perfect graph. A direct justification, avoiding any ambiguity about orientations, is that it is weakly chordal in F : every vertex of a hole $C_k, k \geq 5$, is an endpoint of an induced P_4 ; the same holds for an antihole, since the complement of four consecutive cycle vertices is a P_4 whose endpoints are the two middle cycle vertices. Choose the earliest-layer vertex of a hypothetical hole or antihole and a P_4 in it with that endpoint. Its layer root prefixes this to an induced $F-P_5$, a contradiction. Thus neither F nor G on a partial transversal has a hole of length at least 5. Weakly chordal graphs are perfect [5]. No assertion about perfection inside an individual layer is used.

Apply the thin-layer lemma with $\varepsilon = y^4$. It gives a restricted set of size at least $y^8 M/512 \geq 3y^8 n/(4096t) \geq 3y^{12} n/8192 \geq y^{18} n$, which is B .

Positive direction. Every layer lies in $[xs_{\text{cut}}/4, ts_{\text{cut}})$, so its number L satisfies $L \leq 4t^6/x \leq 256x^{-25} \leq x^{-27}$. There exists m in $[t, L]$ such that m layers each have size at least n/m^3 . Otherwise, writing their sizes in decreasing order a_i , the first $t-1$ contribute less than n/t^4 in

total and $a_i < n/i^3$ for $i \geq t$. Thus $\sum a_i \leq n/t^4 + n/[2(t-1)^2] < n/(8t)$, contradicting positive mass at least $3n/(8t)$. Keep these m layers in their original order and apply Appendix C.5 with $b = 3, \tau = x$. This gives C . The lower size $xs_{\text{cut}}/4$ was used only to bound L , not charged as the final common width.

C.8 Numerical closure

In all frontier/layer applications, $t \leq m \leq x^{-27}, \tau = x, b$ in $\{3, 6\}$. The order requirement $W \geq 16\xi_0^{-3}$ with $W = n/m^b$ reads $n \geq 2^{22}x^{-3}m^{b+15}$. For the positive layers $b = 3$ and $m \leq x^{-27}$, so the right side is at most $2^{22}x^{-489}$; for the frontier $b = 6$ and $m = t \leq 2x^{-4}$, so it is at most $2^{43}x^{-87}$. Both bounds follow from $n \geq x^{-600}$ and $x \leq 2^{-16}$. All actual lengths satisfy $K \geq m^{1/4} \geq t^{1/4} \geq 1/y, K \leq 64m^5/x \leq 64x^{-136} \leq x^{-140}$. The width exponent $4b + 12 \leq 36$ is stronger than the claimed 64. Direct component-grouping outputs have $K = t$ and width at least n/t^7 , also within C . This completes the proof. $\square$

D A polynomial bridge from EH transversals

This bridge supplies the negative-layer input for rooted orders at least six. The proof includes the required rounding and logarithmic estimates. It requires an EH exponent for the actual transversal class; it supplies no such exponent by itself.

Theorem D.1 (Polynomial thin-layer bridge). *Let an M -vertex graph be partitioned into nonempty layers. Suppose that, for every j , every partial transversal on j vertices (a set meeting each layer at most once) induces a graph with a clique or stable set of size at least j^κ , where $0 < \kappa \leq 1$ is fixed. Put $h = \max(2, 1/\kappa)$. If*

$$0 < \varepsilon \leq 2^{-4(h+10)}, \quad \lambda = \frac{\max(\text{layer size})}{M} \leq \varepsilon^{4h+4},$$

then there is an induced subgraph T with $|T| \geq \varepsilon^{2h+2}M$ such that T or its complement has maximum degree at most $\varepsilon(|T| - 1)$. No finite forbidden-subgraph description of the transversal class is required. In particular, this applies to E_5 transversals once an EH exponent for their class has been proved.

Proof. Set

$$q = \lceil \varepsilon^{-2} \rceil, \quad t = \lceil q^h \rceil, \quad \delta = \frac{q}{512t}, \quad L = \ln(1/\varepsilon), \quad \ell = \lceil \varepsilon^{-1} \ln(1/\delta) \rceil.$$

Since $q \leq 2\varepsilon^{-2}$ and $t \leq 2^{h+1}\varepsilon^{-2h}$,

$$32t^2\lambda \leq 2^{2h+7}\varepsilon^4 \leq 1.$$

Thus $\lambda \leq 1/(32t^2)$. Nonempty layers imply $\lambda \geq 1/M$, so $M \geq 32t^2$ and $\delta M \geq qt/16 \geq 1$.

Sample $2t$ distinct vertices uniformly. A sampled pair lies in one layer with probability at most λ . The expected number of colliding pairs is at most $2t^2\lambda \leq 1/16$. With probability at least $1/2$, at most t collisions occur; deleting at most one vertex per collision leaves t representatives from distinct layers. Their homogeneous size is at least $t^\kappa \geq q$. Double counting gives at least

$$\frac{1}{2} \frac{\binom{M}{2t}}{\binom{M-q}{2t-q}} \geq \frac{1}{2} \left(\frac{M}{2t} \right)^q$$

homogeneous q -subsets. One of the two kinds therefore numbers at least $\frac{1}{4}(M/(2t))^q$.

Suppose no set S with $|S| \geq \delta M$ has maximum degree at most $\varepsilon(|S| - 1)$ in the graph or in its complement, and put $u = \lceil \delta M \rceil \leq 2\delta M$. In the orientation whose homogeneous q -sets are stable, every subset of size at least u has maximum degree greater than $\varepsilon(|S| - 1)$. The greedy container encoding gives

$$I_q \leq \binom{M}{\ell} \binom{u}{q - \ell},$$

provided $\ell \leq q$, which is verified below. Repeatedly test a deterministically chosen vertex of maximum degree in the graph induced on the current candidate set R , and stop when $|R| \leq u$. If the vertex is absent from the independent set, remove it alone; otherwise record it and remove its closed neighbourhood, which has more than $\varepsilon(|R| - 1) + 1$ vertices and so leaves fewer than $(1 - \varepsilon)|R|$ candidates. Hence at most ℓ choices are recorded, because $(1 - \varepsilon)^\ell M \leq \delta M \leq u$. If fewer choices suffice, pad the record with vertices of the independent set left in the final container. The ℓ -element record determines the container and gives the stated bound.

Using $\binom{q}{\ell} \leq 2^q$ and $q! \geq (q/e)^q$, comparison with the lower bound yields

$$\left(\frac{1}{2\delta}\right)^{1-\ell/q} \leq 4e \frac{t}{q} 4^{1/q} \leq 8e \frac{t}{q}.$$

If $u < q - \ell$, the container bound is zero and already contradicts the lower bound.

For the rounding and logarithmic term, $t \leq 2q^h$ gives

$$\begin{aligned} A := \ln(1/\delta) &\leq \ln 1024 + (h - 1) \ln q \\ &\leq (h + 9) \ln 2 + 2(h - 1)L \leq 2hL, \end{aligned}$$

where the final comparison uses $L \geq 4(h + 10) \ln 2$. Write $B = \ln(1/(2\delta)) \leq A$. Then

$$\frac{\ell}{q} B \leq (\varepsilon A + \varepsilon^2) B \leq 4h^2 \varepsilon L^2 + 2h \varepsilon^2 L < 1.$$

For an elementary bound, εL^2 and $\varepsilon^2 L$ increase with ε in the stated range. At the cutoff, each resulting expression decreases with $h \geq 2$: its logarithmic derivative is bounded respectively by

$$\frac{2}{h} + \frac{2}{h + 10} - 4 \ln 2 < 0, \quad \frac{1}{h} + \frac{1}{h + 10} - 8 \ln 2 < 0.$$

Consequently the sum is bounded by its value at $h = 2$, $\varepsilon = 2^{-48}$, which is less than 2^{-30} . Since $B > 1$, this also gives $\ell < q$.

Restoring the logarithmic factor in the counting inequality gives $1/(2\delta) < 8e^2(t/q)$, whereas its definition gives $1/(2\delta) = 256t/q$, a contradiction. Thus the desired restricted set has size at least δM . Finally,

$$\delta = \frac{q}{512t} \geq 2^{-(h+9)} \varepsilon^{2h-2} \geq \varepsilon^{2h+2},$$

because $\varepsilon^4 \leq 2^{-(h+9)}$. This proves the theorem. $\square$

E RP6 frontiers and the third offending-pair purification

The two first calls use the RP5 contract of Appendix C; the third uses the original RP4/general Tooth of Appendix A. All roots and witnesses belong to the same original host.

Let $m \geq 2^{16}$ be an integer and $0 < \tau \leq 1$. Let $Z_1, \dots, Z_m$ be disjoint subsets of Y . For each j let R_j be a set of at most m roots outside Y . Assume:

- (I) Every root in every R_j has RP6 into the original set Y , and every r in R_j is G -complete to Z_j .
- (II) Every vertex of a later Z_j has RP5 into each earlier Z_i .
- (III) There is a fixed graph J on $[m]$ such that all pairs outside J are G -complete. On every increasing J -path, for any two path indices $j < l$, some root in R_j is G -anticomplete to the entire Z_l .

Here an increasing path means a directed path when each edge is oriented from its smaller to its larger endpoint.

This abstract input covers two constructions used by the tree proof.

For a positive sequence, each Z_j has its own root complete to Z_j and anticomplete to all later sets; use its singleton root family and J complete. Property (II) follows by prefixing that own root to a putative rooted F - P_5 .

For an ordered cut-tree frontier, a node S is split using a root r into the F -components of $N_G(r) \cap S$, in any fixed order, followed by $A = S$ minus $N_G(r)$. Nonempty children inherit their order recursively. Take any antichain of m tree nodes, in this leaf order. For every pair $j < l$ whose lowest common ancestor separates them into a C child and A , put one such separating root in R_j . There are at most $m - 1$ chosen roots per j . Let J consist exactly of these C/A pairs. Two frontier sets in different C children of their lowest common ancestor are G -complete. They are therefore not edges of J .

An increasing J -path cannot visit two different C -child intervals at any one split. If it did, its path segment inside that node would cross between two consecutive visited C -child intervals. The crossing path edge would join different C children and thus would not be in J . There can be no intervening path vertices outside the node interval. Consequently any two path vertices whose lowest common ancestor is this node lie in a C child and A , rather than in different C children. This proves (III). In the C/A case a separator is complete to the earlier frontier set and anticomplete to the later one. Prefixing it to a rooted F - P_5 from a later vertex proves (II); for different C children (II) is immediate because the pair is G -complete.

The rooted laws always refer to original Y . They are inherited under all subsequent restrictions of blocks, and roots need not be adjacent to each other.

E.1 Quantitative conclusion

Lemma E.1 (RP6 frontier purification). *Suppose, in addition, that $|Z_i| \geq W$ for every i , where $W \geq 2^{7200} \tau^{-600} m^{36000}$. For an ambient scale n and a fixed $b \geq 0$, suppose also $|Z_i| \geq n/m^b$.*

Then at least one of these two conclusions holds.

- (A) *A set $T \subseteq \bigcup_i Z_i$ is m^{-4} -restricted and $|T| \geq n/m^{b+44}$.*
- (B) *There is a directed τ -semisparse blockade in $\bigcup_i Z_i$ of integer length K with $m^{1/4} \leq K \leq m^{9000} \tau^{-140}$ and width at least n/K^{4b+216} . Each complete or anticomplete early output is included as a special case.*

The parameter m in (A) is the actual input length. In particular (A) cannot be relabeled as a fixed- y -width conclusion if m is as large as a power of $1/x$. Its use in the full-tree theorem

is given in Appendix F.

E.2 First and canonical RP5 purification

Proof. Set $a_0 = \tau/(2^{12}m^{60})$, $\theta = a_0/4 = \tau/(2^{14}m^{60})$, $a_1 = \tau/(64m^2)$, $\eta = 1/(16m^{28})$. Both RP₅ Tooth calls use coarse parameter $1/m$.

First apply RP₅ Tooth to each Z_i , using all vertices of the later original Z blocks as its roots. This is legal by (II), and the order hypothesis is exactly $|Z_i| \geq a_0^{-600}$. If an early outcome occurs, record it for Appendix E.5. Otherwise retain B_i with $|B_i| \geq |Z_i|/m^{26}$, such that every later vertex is complete to B_i or has fewer than $\theta|B_i|$ neighbours in B_i . Freeze these original two types.

For $i < j$ call the pair good if some r in R_j has at least $\eta|B_i|$ nonneighbours in B_i ; call it red otherwise. For each good pair choose one such r . Let S_{ij} be the sparse-type part of B_j toward B_i . By averaging over B_i minus $N_G(r)$, choose p_{ij} there with $\deg_G(p_{ij}, S_{ij}) \leq \theta|B_j|/\eta$. The representative p_{ij} has RP₅ into B_j : an induced F -P₅ from p_{ij} inside B_j could be prefixed by r , giving an induced F -P₆ from r . Here r is G -complete to B_j and G -nonadjacent to p_{ij} .

Apply RP₅ Tooth to B_j with the at most m representatives p_{ij} , using fine parameter a_1 and coarse parameter $1/m$. The size condition holds: $|B_j| \geq 2^{7200}\tau^{-600}m^{35974} \geq a_1^{-600} = 2^{3600}\tau^{-600}m^{1200}$. Again record any early output. Otherwise obtain D_j with $|D_j| \geq |B_j|/m^{26}$, on which each representative is complete or strictly $a_1/4$ -sparse.

For each good pair delete from D_j the opposite original type: if p_{ij} is sparse on D_j , delete all full-type vertices; if it is full on D_j , delete all sparse-type vertices. Using the original D_j for every budget, the total fraction deleted is at most $m(a_1/4 + \theta m^{26}/\eta) = \tau/(256m) + \tau/(2^{10}m^5) < 1/16$.

For every red pair $i < j$, delete from D_i the nonneighbours of every root in R_j . At most m^2 root occurrences affect a fixed earlier block B_i . The total loss from this operation is at most $m^2\eta|B_i| = |B_i|/(16m^{26}) \leq |D_i|/16$. All operations use the original sets, so they may be done simultaneously.

Call the surviving sets C_i . Then $|C_i| \geq |B_i|/(2m^{26}) \geq |Z_i|/(2m^{52})$. Every nonred pair is uniform in its original type, and every root in R_j is exactly G -complete to C_i on a red pair $i < j$. Each such root remains G -complete to its own C_j .

Each old sparse type has at most $\beta|C_i|$ neighbours in C_i , where $\beta = 2m^{26}\theta = \tau/(2^{13}m^{34}) < 1/2$. Because $\beta < 1$, sparse types cannot have become complete. Consequently every ambiguous C pair is red. Let Q be its ambiguity graph. It is a subgraph of J : a G -complete original pair never becomes ambiguous. Conditions (I) and (III) hold for the inherited root families.

No connected component of $F[C_i]$ is selected or needed.

E.3 The new offending-pair pass

Freeze the original C_i and their β -types for this whole step. Call an old ambiguous pair $i < j$ offending if there exist $l > j$, z in C_l , and r in R_j such that r is G -nonadjacent to z , z is G -complete to C_j , z has sparse type toward C_i . The quantifier runs over all later original vertices z . It is not restricted to the vertices that will survive subsequent trimming. For each offending pair fix one witness (l, z, r) .

The pair is red, so this r is G -complete to C_i as well as to C_j . The set C_i minus $N_G(z)$ has size at least $(1 - \beta)|C_i|$. Let S_{ij} now denote the old sparse-type portion of C_j toward C_i . Averaging gives b_{ij} in C_i minus $N_G(z)$ with $\deg_G(b_{ij}, S_{ij}) \leq \beta|C_j|/(1 - \beta) \leq 2\beta|C_j|$.

This representative $b = b_{ij}$ has RP₄ into C_j . Indeed, an induced F -P₄ b - v - w - d , with v, w ,

d in C_j , would extend to $r-z-b-v-w-d$, an induced $F-P_6$ with endpoint r . The new consecutive pairs rz and zb are F -edges. All new nonconsecutive pairs are F -nonedges: r is G -complete to C_i and C_j , and z is G -complete to C_j . The three nonconsecutive pairs inside $b-v-w-d$ are absent in F by its induced- P_4 hypothesis. All six vertices are distinct: r lies outside Y and $z, b, \{v, w, d\}$ lie in three different blocks. This contradicts (I).

There is at most one b_{ij} per earlier i , hence fewer than m representatives into each C_j . Apply the original general Tooth to C_j with these representatives and $\xi = m^{-2}, k = m$. Its numerical hypotheses hold: $\xi \leq 2^{-20}, 2^{16} \leq m \leq 2\xi^{-1/2}$, and $|C_j| \geq 16m^6$. If a TL1–TL4 output occurs, retain it. Otherwise obtain D'_j with $|D'_j| \geq |C_j|/m$ such that every chosen b_{ij} is complete or strictly $\xi/4$ -sparse on D'_j .

For each offending pair, delete from D'_j the opposite old C type. If b_{ij} is sparse on D'_j , deleting full-type vertices costs strictly less than $\xi|D'_j|/4$. If b_{ij} is full on D'_j , deleting sparse-type vertices costs at most $2\beta|C_j| \leq 2m\beta|D'_j|$. The total loss over fewer than m pairs is at most $m(\xi/4 + 2m\beta) = 1/(4m) + 2m^2\beta < 1/2$ of D'_j . Write E_j for the survivors. Thus $|E_j| \geq |C_j|/(2m) \geq |Z_j|/(4m^{53})$, and every old offending pair is uniform in its original C type.

All exact root relations survive. Original sparse types now have threshold $\beta' = 2m\beta < 1$. Original full types remain full. Conversely, a vertex complete to E_i must have had full type toward C_i , because an old sparse-type vertex has at most $\beta'|E_i| < |E_i|$ neighbours there. This converse applies to every later vertex and every earlier index.

Let Q' be the new ambiguity graph. It is a subgraph of Q and of J , and none of the old offending pairs is an edge of Q' .

E.4 Restored monotonicity and a large blockade

Consider an increasing Q' -path. Let $i \rightarrow j$ be one of its edges and let $l > j$ be a later path index. By (III), some r in R_j is anticomplete to the entire original Z_l , hence to the original C_l .

A vertex z in E_l cannot be sparse-type to E_i and complete to E_j : by $\beta' < 1$ its old types toward C_i, C_j would be sparse and complete respectively. Together with r this would witness that $i < j$ was an old offending pair. That pair was purified, so it cannot be an edge of Q' .

Thus each later vertex on an increasing Q' -path has a prefix of full types followed by a suffix of sparse types on earlier path vertices.

Let $r_0 = \lfloor \sqrt{m} \rfloor$. If Q' has an increasing path of r_0 vertices, partition each corresponding E block according to this prefix length and retain a largest cell. At most r_0 cells occur per block. The retained r_0 blocks have width at least $\min |Z_i|/(4m^{53}r_0)$ and are directed $(r_0\beta')$ -semisparse. Their precision is at most τ .

If there is no such path, assign each index its longest-path height. Equal-height classes are independent in Q' . With fewer than r_0 heights, one class contains at least r_0 indices. Select r_0 blocks from it. Their pairs were already uniform, and they have even better width and precision than in the preceding paragraph.

This proves the main directed semisparse outcome. The blocks C_i and E_i need not be F -connected: the six-vertex contradiction in Appendix E.3 uses only the rooted exclusions and the preserved full/sparse types.

E.5 Early outcomes and actual lengths

The first RP_5 Tooth restricted outcome has precision m^{-4} and size $|Z_i|/m^{18}$. The second has that precision and size $|B_i|/m^{18} \geq |Z_i|/m^{44}$. Hence each gives conclusion (A).

The first RP_5 blockade outcome has actual $K \geq m, K \leq a_0^{-140}$, precision $a_0 \leq \tau$, and width at least n/K^{b+64} . The second has $K \geq m, K \leq a_1^{-140} \leq a_0^{-140}$, precision $a_1 \leq \tau$, and width at least n/K^{b+90} .

For the third Tooth, $|C_j| \geq |Z_j|/(2m^{52}) \geq n/m^{b+53}$. Substituting $\xi = m^{-2}, k = m$ into its four early outcomes gives: TL1: $K \geq m^{1/4}$, width at least n/K^{4b+216} .

TL2: $K \geq \sqrt{m}$, width at least n/K^{2b+109} .

TL3: $K = m^2$, width at least $n/K^{(b+58)/2}$.

TL4: $K = m$, width at least n/K^{b+61} .

In TL2, $K \geq m^{3/4}/9 \geq \sqrt{m}$, using $m \geq 2^{16}$.

For the main output $K = r_0 = \lfloor \sqrt{m} \rfloor, m \leq r_0^3$ and $r_0 \geq 4$ imply $n/(4m^{b+53}r_0) \geq n/r_0^{3b+161}$. All displayed width exponents are at most $4b + 216$.

Every output length is at least $m^{1/4}$. Every early or main length is at most a_0^{-140} , since all third-Tooth lengths are at most m^2 : $a_0^{-140} = 2^{1680}m^{8400}\tau^{-140} \leq m^{8505}\tau^{-140} \leq m^{9000}\tau^{-140}$. This completes conclusion (B) with its actual length and width. $\square$

F The RP_6 full-tree theorem with its actual floating scale

The frontier input is proved in Appendix E and the transversal input in Appendix D. We first state the theorem under $\text{EH}(E_5)$, then give the relative variant under $\text{EH}(E_5 \cap \mathcal{A})$ for a hereditary class $\mathcal{A}$ containing the whole host. Part II uses the relative variant. The local size cutoff of the cut tree is written $s_{\text{cut}} = n/t^6$; it is unrelated to the fixed target path order s .

F.1 Inputs, constants, and statement

Theorem F.1 (Two-parameter RP_6 Tooth). *Write $F = \overline{G}$. Let E_5 be the hereditary class of graphs admitting an order such that the first vertex of every induced P_5 is not an endpoint. Equivalently each vertex has no rooted P_5 into its suffix. Assume every j -vertex member of E_5 has a clique or stable set of size at least j^{κ_E} , where $0 < \kappa_E \leq 1$. The unrestricted $\text{EH}(E_5)$ premise is not an input to the main fixed-target induction; the relative $\text{EH}(\mathcal{A} \cap E_5)$ variant below is used there.*

Put $h = \max(2, 1/\kappa_E)$, $d_t = \lceil 4h+5 \rceil$, $c_ = \min(2^{-16}, 2^{-(h+10)})$, $C = 6d_t + 3$, $A_{\text{clean}} = 6d_t + 2$, $e = \max(50, 8h + d_t + 10)$, $K_{\text{exp}} = 9000C + 140$, $E = 7801 + 36006C$.*

Let $0 < x \leq y \leq c_$, let Y have $n \geq x^{-E}$ vertices, and let U be outside Y . Suppose every u in U has RP_6 into the original Y : $F[Y \cup \{u\}]$ has no induced six-vertex path with endpoint u .*

Then at least one of the following holds.

- (A) *There is $S \subseteq Y$ with $|S| \geq y^{A_{\text{clean}}}n$, such that every u in U is G -complete to S or has strictly fewer than $x|S|/4$ neighbours in S .*
- (B) *For some real z with $x^C \leq z \leq y$, there is a z^4 -restricted set $T \subseteq Y$ of size at least $z^e n$.*
- (C) *There is an ordered blockade in Y of integer length K with $1/y \leq K \leq x^{-K_{\text{exp}}}$ and width at least n/K^{240} , such that every pair is complete or directed x -sparse from its later block to its earlier block.*

Here ε -restricted means $G[T]$ or $F[T]$ has maximum degree at most $\varepsilon|T|$. Directed x -sparse means every vertex in the later block has at most $x \cdot$ the earlier block size neighbours in that block. The restricted scale z in (B) is deliberately allowed to vary down to a fixed power of x .

F.2 The full cut tree and its cutoff

Proof. Set $t = \lceil y^{-d_t} \rceil$, $s_{\text{cut}} = n/t^6$. Then $t \geq 2^{16}$ and $t \leq 2y^{-d_t} \leq 2x^{-d_t}$.

Start at node Y . At a node S of size at least s_{cut} , stop successfully if every root in U is complete to S or has degree strictly less than $x|S|/4$ there. Otherwise choose a bad root u . Its degree satisfies $x|S|/4 \leq |N_G(u) \cap S| < |S|$. Split S into the F -components of $N_G(u) \cap S$, in any fixed order, followed by $A = S$ minus $N_G(u)$. Ignore empty children. Both the neighbour set and A are nonempty, so all children are proper. Continue at children of size at least s_{cut} ; smaller children are atoms.

A clean node gives (A) , because $|S| \geq s_{\text{cut}} \geq 2^{-6}y^{6d_t}n \geq y^{6d_t+2}n$. The process is finite. If no clean node occurs, all terminal pieces are atoms, and they partition Y .

Call the tree induced by nodes of size at least s_{cut} the retained tree. If it has t incomparable nodes, they are a frontier with $m = t$ and $|Z_i| \geq s_{\text{cut}} = n/m^6$. Apply the frontier lemma with $\tau = x, b = 6$. Its order hypothesis is verified in Appendix F.6. Its restricted output gives (B) with $z = 1/m$; its blockade output gives (C) . Thus assume that no t incomparable retained nodes exist.

The retained tree has fewer than t leaves. Its branching nodes and terminal nodes together number fewer than $2t$; call these exceptional. All remaining retained nodes are unary.

F.3 Small components and mass on one complete chain

At any retained node let D be the union of its neighbour-component children of size less than s_{cut} . If $|D| \geq ts_{\text{cut}}$, group these components into t disjoint sets each of size at least $|D|/t^2$. The groups are pairwise G -complete and have width at least $s_{\text{cut}}/t = n/t^7$. This is (C) .

For completeness, the grouping works because each component has size less than $s_{\text{cut}} \leq |D|/t$. Greedily form $t - 1$ groups each just reaching $|D|/t^2$. Each such group has size less than $(t + 1)|D|/t^2$. Their total is less than $(t^2 - 1)|D|/t^2$, leaving at least $|D|/t^2$ for the final group. No component is cut.

Henceforth suppose every retained node has $|D| < ts_{\text{cut}}$. Its atoms then have total size less than $(t + 1)s_{\text{cut}}$, since its A child contributes either zero or fewer than s_{cut} vertices. Exceptional-node atoms have total mass less than $2t(t + 1)s_{\text{cut}}$. The unary nodes therefore carry atom mass at least $n - 2t(t + 1)s_{\text{cut}}$.

Fewer than t complete root-to-leaf paths cover the retained tree. Consequently one complete path carries unary atom mass at least $n/t - 2(t + 1)s_{\text{cut}}$. It may pass through branching nodes. Their atoms are ignored, but the full containment chain remains available for all witnesses.

At each unary node on this path there is exactly one large child.

If the large child is A , its whole neighbour set is a positive layer P . It obeys $xs_{\text{cut}}/4 \leq |P| < ts_{\text{cut}}$, and its split root is G -complete to P and G -anticomplete to the whole later continuation.

If the large child is a neighbour component, its small A child is a negative layer, of size less than s_{cut} . The split root is G -anticomplete to this negative layer and G -complete to the whole later continuation. There may also be small neighbour-component siblings.

All such small siblings from negative steps are mutually G -complete, including siblings from different levels: an earlier sibling is complete to the selected large neighbour component containing the later chain. If their total mass is at least ts_{cut} , the same grouping gives (C) . Otherwise discard this fringe, losing less than ts_{cut} .

The remaining positive and negative layers have total mass at least $n/t - (3t + 2)s_{\text{cut}} \geq$

$3n/(4t)$. Thus one of their two directions has mass at least $3n/(8t)$.

F.4 The negative direction

Suppose the negative layers have total mass $M \geq 3n/(8t)$. Each has size less than s_{cut} , so the largest-layer fraction satisfies $\lambda \leq 8/(3t^5)$.

Every transversal belongs to E_5 . Indeed, order its vertices by their layers along the chain. If the earliest vertex of an induced P_5 were its endpoint, prefix the root witnessing that earliest negative layer. In F the root is adjacent to that vertex and nonadjacent to every later transversal vertex. This creates an induced rooted P_6 inside the original Y plus that root, a contradiction. No claim that these transversals are P_7 -free or perfect is needed.

Take $\varepsilon = y^4$. The choice $y \leq 2^{-(h+10)}$ gives $\varepsilon \leq 2^{-4(h+10)}$. Also $\lambda \leq (8/3)y^{5d_t} \leq y^{16h+16} = \varepsilon^{4h+4}$. Indeed $5d_t - 16h - 16 \geq 4h + 9$, and $y \leq 2^{-16}$ absorbs the factor $8/3$.

The EH-transversal bridge supplies a y^4 -restricted set T with

$$|T| \geq y^{8h+8}M \geq \frac{3}{16}y^{8h+8+d_t}n \geq y^{8h+d_t+10}n.$$

Since $e \geq 8h + d_t + 10$, this is outcome (B) with $z = y$. Its range condition holds because $x^C \leq x \leq y$.

F.5 The positive direction

Suppose instead the positive layers have total mass at least $3n/(8t)$. Each has size less than $t s_{\text{cut}} = n/t^5$, and at least $x s_{\text{cut}}/4$. If there are L of them, disjointness gives $L \leq 4t^6/x \leq 256x^{-(6d_t+1)} \leq x^{-(6d_t+3)} = x^{-C}$.

There exists an integer m with $t \leq m \leq L$ for which at least m positive layers have size at least n/m^3 . To verify this, list their sizes decreasingly as $a_1 \geq \dots \geq a_L$. If no such m exists, then $a_m < n/m^3$ for every $m \geq t$. The first $t-1$ entries each have size less than n/t^5 . Therefore $\sum_i a_i < n/t^4 + n/(2(t-1)^2) < n/(8t)$, using the integral bound for $\sum_{m \geq t} m^{-3}$. This contradicts the assumed mass. If $L < t$, the first term alone gives a contradiction.

Keep those m layers in their original order. They form an RP_6 positive sequence. Apply the frontier/positive lemma with $\tau = x, b = 3$. The resulting restricted set has precision m^{-4} and size at least n/m^{47} . Put $z = 1/m$. Since $t \leq m \leq x^{-C}$, $x^C \leq z \leq 1/t \leq y$, and $e \geq 50 \geq 47$. This gives (B). The blockade gives (C) by the calculations below.

It is essential that n/m^3 , rather than the weak $x s_{\text{cut}}/4$ individual lower bound, is used as the final block width.

F.6 All order, scale, and actual-length checks

Every frontier or positive application has $t \leq m \leq x^{-C}$, $\tau = x, b$ in $\{3, 6\}$. Its order requirement is $n \geq 2^{7200}x^{-600}m^{36000+b}$. The right side is at most $2^{7200}x^{-[600+36006C]}$. Since $E = 7801 + 36006C$ and $x \leq 2^{-16}$, the hypothesis $n \geq x^{-E}$ implies this with substantial room. Hence every subsidiary RP_5 or general-Tooth order requirement included in the local lemma is met.

The local restricted output has precision z^4 at $z = 1/m$ and size at least $z^{b+44}n \geq z^{50}n \geq z^e n$. The interval $x^C \leq z \leq y$ was checked above.

Every local blockade has actual $K \geq m^{1/4} \geq t^{1/4} \geq y^{-d_t/4} \geq 1/y$, because $d_t \geq 4$. Its upper length is $K \leq m^{9000}x^{-140} \leq x^{-(9000C+140)} = x^{-K_{\text{exp}}}$. Its width exponent $4b + 216$ is at most 240.

The elementary component-grouping outputs have actual $K = t$, width at least n/t^7 , and are complete. Their length is at least $1/y$ and at most x^{-C} , hence at most $x^{-K_{\text{exp}}}$; their width is at least n/K^{240} . Thus these outputs satisfy the same contract (C).

The clean bound and negative restricted bound were explicit in Appendices F.2 and F.4. These checks exhaust all outputs of the finite tree process and prove the theorem. $\square$

F.7 Fixed-ambient variant

Proposition F.2 (Relative-ambient RP6 Tooth). *Let $\mathcal{A}$ be any hereditary class of graphs F . Suppose the original host $F[Y \cup U]$ belongs to $\mathcal{A}$, and assume EH only for $\mathcal{A} \cap E_5$, with exponent κ_E . Then the conclusion of Theorem F.1 holds with the same constants.*

Proof. Indeed the RP₅ Tooth used by the frontier proof is ambient-independent. All blocks, later-vertex roots, canonical representatives, offending witnesses and original roots remain in the original host. Every negative transversal is an induced subgraph of that host and belongs to E_5 by Appendix F.4, hence lies in $\mathcal{A} \cap E_5$. That is the sole place where EH(E_5) was used. Replacing it by the weaker relative-class hypothesis therefore makes no other change to the proof.

In particular, for a fixed target path P_s one may take $\mathcal{A} = \text{Forb}(P_s)$, and use EH($E_5 \cap \text{Forb}(P_s)$) obtained from the ordered-tail transfer. The forbidden ambient path is not changed between recursive calls. $\square$

G The RP_q recurrence in the same hereditary ambient class

Unlike the specialized RP6 base, both lower rooted orders now have the common floating-scale contract. Every lower call uses an induced subhost of the same original $F \in \mathcal{A}$.

G.1 Common lower-tooth contract

Let $q \geq 7$. Fix a hereditary class $\mathcal{A}$ of graphs F , and work in $G = \overline{F}$. Assume both RP _{$q-1$} and RP _{$q-2$} have the following Tooth contract within $\mathcal{A}$. After enlarging constants, use the same constants $A_0, e_0, d_0, c_0, k_0, E_0 \geq 1$ and $0 < \eta_0 \leq 2^{-16}$ for both contracts.

For $0 < a \leq b \leq \eta_0$ and a vertex set V of size $N \geq a^{-E_0}$, with outside roots whose original ambient F remains in $\mathcal{A}$ and that have the indicated rooted path exclusion into V , at least one of the following exists:

- (L₁) a set $S \subseteq V$ with $|S| \geq b^{A_0} N$ on which every root is complete in G or has strictly fewer than $a|S|/4$ G -neighbours;
- (L₂) a z^4 -restricted subset of V of size at least $z^{e_0} N$, for some $z \in [a^{c_0}, b]$;
- (L₃) a directed a -semisparse blockade in V of integer length K with $1/b \leq K \leq a^{-k_0}$ and width at least N/K^{d_0} .

As usual, RP _{j} means absence of an induced j -vertex path in F with the specified root as an endpoint; ε -restricted allows either G or F to have maximum degree at most $\varepsilon \cdot$ the set size. All roots and all blocks are induced subsets of the same original ambient graph $F \in \mathcal{A}$.

G.2 A quantified RP_q frontier statement

Lemma G.1 (RP_q frontier purification). *Put*

$$J_0 = 4A_0 + 11, \quad H = c_0 J_0, \quad J = k_0 J_0, \quad Q = E_0 J_0, \quad m_0 = \max(2^{16}, \lceil 1/\eta_0 \rceil). \quad (\text{G.1})$$

Let $m \geq m_0$ be an integer, $0 < \tau \leq 1$, and $b_0 \geq 0$. Suppose $Z_1, \dots, Z_m$ form an RP_q ordered cut-tree frontier, or an RP_q positive sequence, inside an original n -scale vertex set Y . That is, the sets Z_i arise from one of the two constructions of Appendix E, with roots outside Y that have RP_q into the original Y ; these constructions supply root families R_j of at most m roots satisfying (I) and (III) there, with RP_6 replaced by RP_q . The separation graph called J in that appendix is unrelated to the exponent J of (G.1). Suppose also that

$$|Z_i| \geq (m/\tau)^Q, \quad |Z_i| \geq n/m^{b_0}. \quad (\text{G.2})$$

Then at least one of the following holds.

- (F₁) For some z with $(\tau/m)^H \leq z \leq 1/m$, a z^4 -restricted set $T \subseteq \bigcup_i Z_i$ has $|T| \geq z^{e_0+b_0+2A_0+1}n$.
- (F₂) There is a directed τ -semisparse blockade in $\bigcup_i Z_i$ of integer length K with $m^{1/4} \leq K \leq (m/\tau)^J$ and width at least $n/K^{3b_0+9A_0+d_0+3}$.

Proof. The frontier and root-family definitions are exactly those in Appendix E, with RP_6 replaced by RP_q . An ordered cut tree splits at a root into the F -components of its G -neighbourhood, followed by its G -nonneighbourhood A child. At most m roots per block are recorded from C/A lowest common ancestors. Pairs in different C children are G -complete. Along any increasing ambiguity path no two indices lie in different C -child intervals of one split, so every two such indices have a recorded separating root. These are purely order/interval facts, independent of q . In a positive sequence the own roots separate every earlier/later pair.

A later vertex has RP_{q-1} into each earlier Z_i : a hypothetical rooted P_{q-1} prefixes with the earlier separating root to form an RP_q violation. Different C -child pairs are complete and immediate.

G.3 Two preparatory passes

Use $a_0 = \tau/(2^{12}m^{4A_0+10})$, $a_1 = \tau/(64m^2)$, $\eta = 1/(16m^{A_0+2})$, $\theta = a_0/4$. All lower Tooth applications below use coarse parameter $1/m$.

The first RP_{q-1} Tooth, with all later original vertices as roots, either gives an early output or yields $B_i \subseteq Z_i$ with $|B_i| \geq |Z_i|/m^{A_0}$. Every later vertex is full or θ -sparse toward B_i . Freeze its type.

For each $i < j$, call the pair good if some recorded root r in R_j has at least $\eta|B_i|$ nonneighbours in B_i . Otherwise call it red. For a good pair choose such r and a representative p in B_i minus $N_G(r)$ having at most $\theta|B_j|/\eta$ neighbours in the sparse-type part of B_j . This follows by averaging. Since r is complete to B_j , a rooted F - P_{q-1} from p into B_j could be prefixed by r . Thus p has RP_{q-1} into B_j .

Apply the second RP_{q-1} Tooth to B_j with these fewer than m representatives and fine parameter a_1 . Unless an early output occurs, retain $D_j \subseteq B_j$ with $|D_j| \geq |B_j|/m^{A_0}$. Purify every good pair by deleting the opposite frozen type according to whether its representative is full or $a_1/4$ -sparse on D_j . The total loss fraction is at most $m(a_1/4 + \theta m^{A_0}/\eta) = \tau/(256m) + \tau/(2^{10}m^{2A_0+7}) < 1/16$.

For every red pair $i < j$, remove from D_i all nonneighbours of every root in R_j . There are at most m^2 relevant root occurrences, so this loses at most $m^2\eta|B_i| = |B_i|/(16m^{A_0}) \leq |D_i|/16$. The resulting C_i satisfy $|C_i| \geq |Z_i|/(2m^{2A_0})$. Every ambiguous pair is red, with all its recorded later roots full to the earlier C_i . The old sparse threshold becomes $\beta = 2m^{A_0}\theta = \tau/(2^{13}m^{3A_0+10}) < 1/2$. No sparse type becomes full.

G.4 Offending pairs and the RP_{q-2} pass

Freeze the C_i and their types. An ambiguous pair $i < j$ is offending if for some later original z in C_i and some r in R_j , r is nonadjacent to z in G , z is full to C_j , z is sparse-type to C_i . Choose one such witness for every offending pair. Since the pair is red, r is full to both C_i and C_j .

Averaging over C_i minus $N_G(z)$ chooses a representative p there with at most $2\beta|C_j|$ neighbours in the sparse-type part of C_j . The representative p has RP_{q-2} into C_j : a rooted induced $F\text{-}P_{q-2}$ from p could be prefixed by $r\text{-}z$ to give an induced $F\text{-}P_q$. The two new consecutive edges are $r\text{-}z$ and $z\text{-}p$. All new nonconsecutive pairs are absent in F , since r is G -full to C_i, C_j and z is G -full to C_j . Distinctness follows from the root outside Y and the three distinct blocks. This argument uses no F -connectedness.

Apply the RP_{q-2} Tooth with these fewer than m representatives, fine parameter a_1 , and coarse $1/m$. If it gives no early output, it yields $D'_j \subseteq C_j$ with $|D'_j| \geq |C_j|/m^{A_0}$. Purify every offending pair by deleting its opposite old C type. The total loss fraction is at most $m(a_1/4 + 2m^{A_0}\beta) = \tau/(256m) + \tau/(2^{12}m^{2A_0+9}) < 1/2$. Retain E_j with $|E_j| \geq |C_j|/(2m^{A_0}) \geq |Z_j|/(4m^{3A_0})$.

Old sparse types have new threshold $\beta' = 2m^{A_0}\beta = \tau/(2^{12}m^{2A_0+10}) < 1$. Thus a vertex full to an E_i was originally full to C_i . Conversely an originally sparse vertex stays sparse. The new ambiguity graph is a subgraph of the old one and excludes all old offending pairs.

Along an increasing ambiguity path, a later vertex cannot have type 0 then 1 across an earlier path edge. The recorded separating root would make this edge an old offending pair, using the converse type preservation just established. Therefore each later vertex has a prefix of full types followed by sparse types on earlier path vertices.

With $r = \lfloor \sqrt{m} \rfloor$, a path of r vertices has at most r profiles per block, so retain a largest profile cell. The width is at least $\min |Z_i|/(4m^{3A_0}r)$, and precision is $r\beta' \leq \tau$. If no such path exists, longest-path height levels give r pairwise nonambiguous indices and a still stronger width. This is the main blockade output.

G.5 Order hypotheses and early outputs

The assumption $|Z_i| \geq (m/\tau)^Q$ dominates $a_0^{-E_0}$: $2^{12} \leq m$ and $Q = E_0(4A_0+11)$. Furthermore $a_0^{-E_0}/a_1^{-E_0} = 64^{E_0}m^{(4A_0+8)E_0}$. This dominates both m^{A_0} and $2m^{2A_0}$. Hence the second and third lower-Tooth input order requirements hold.

Every restricted early output has $z \leq 1/m$. At the first call its size is at least $z^{e_0}n/m^{b_0}$. At the second it is at least $z^{e_0}n/m^{b_0+A_0}$. At the third it is at least $z^{e_0}n/(2m^{b_0+2A_0})$. All are at least $z^{e_0+b_0+2A_0+1}n$. Their lower scale is at least $a_0^{c_0}$: $a_0 \geq \tau/m^{4A_0+11} \geq (\tau/m)^{J_0}$, so $z \geq (\tau/m)^H$.

Every blockade early output comes from (L_3) , at coarse parameter $1/m$ and fine parameter $a \in \{a_0, a_1\}$. It is a directed a -semisparse blockade of integer length K with $m \leq K \leq a^{-k_0}$, inside one of the sets Z_i, B_j, C_j . Since $a_0 \leq a_1 \leq \tau$, it is directed τ -semisparse. Its length satisfies $K \geq m \geq m^{1/4}$ and, by the bound $a_0 \geq (\tau/m)^{J_0}$ just used,

$$K \leq a_0^{-k_0} \leq (m/\tau)^{k_0 J_0} = (m/\tau)^J.$$

The input orders of the three calls are at least n/m^{b_0} , $n/m^{b_0+A_0}$ and $n/(2m^{b_0+2A_0})$ respectively. In the weakest case, the third, the width is at least

$$\frac{n}{2m^{b_0+2A_0}K^{d_0}} \geq \frac{n}{K^{b_0+2A_0+d_0+1}} \geq \frac{n}{K^{3b_0+9A_0+d_0+3}}, \quad (\text{G.3})$$

using $2 \leq m \leq K$. Hence every (L_3) early output satisfies (F_2) .

The main output has $K = r = \lfloor \sqrt{m} \rfloor$. Here $r \geq 2^8$ and $m < (r+1)^2 \leq r^3$, so $K \geq m^{1/4}$ and $K \leq m \leq (m/\tau)^J$. Its width is at least

$$\frac{n}{4m^{b_0+3A_0}r} \geq \frac{n}{r^{3b_0+9A_0+3}} \geq \frac{n}{K^{3b_0+9A_0+d_0+3}},$$

and its precision is at most $r\beta' \leq \tau$. The level-set alternative has the same length, at least this width, and uniform pair types. Thus (F_2) holds in every blockade case, and each restricted case gives (F_1) . $\square$

G.6 The full RP_q tooth recurrence

Theorem G.2 (The relative-ambient RP_q recurrence). *Let E_{q-1} consist of graphs with an order in which the first vertex of every induced P_{q-1} is not an endpoint. In addition to the lower-Tooth hypotheses, assume that the hereditary class $\mathcal{A} \cap E_{q-1}$ has EH exponent κ , where $0 < \kappa \leq 1$. With J_0, H, J, Q, m_0 as in (G.1), set*

$$\begin{aligned} h &= \max(2, 1/\kappa), & d_t &= \lceil 4h + 5 \rceil, & C_{\text{tree}} &= 6d_t + 3, \\ \eta_{\text{new}} &= \min(2^{-16}, 2^{-(h+10)}, 1/m_0), \end{aligned} \tag{G.4}$$

and

$$\begin{aligned} A_{\text{new}} &= 6d_t + 2, & e_{\text{new}} &= \max(e_0 + 2A_0 + 7, 8h + d_t + 10), \\ c_{\text{new}} &= H(C_{\text{tree}} + 1), & d_{\text{new}} &= 9A_0 + d_0 + 21, \\ k_{\text{new}} &= J(C_{\text{tree}} + 1), & E_{\text{new}} &= Q + (Q + 6)C_{\text{tree}} + 1. \end{aligned} \tag{G.5}$$

For $0 < x \leq y \leq \eta_{\text{new}}$ and $n \geq x^{-E_{\text{new}}}$, an RP_q input inside the fixed ambient class $\mathcal{A}$ has the lower-Tooth contract of Appendix G.1 with $(A_0, c_0, e_0, k_0, d_0, E_0, \eta_0)$ replaced by the corresponding new constants.

Proof. Here are all reductions needed for this assertion. Build the full cut tree at $t = \lceil y^{-d_t} \rceil$ and $s_{\text{cut}} = n/t^6$; this local size cutoff is unrelated to a target path order. The finite tree and mass argument is exactly the one written in full in Appendices F.2–F.5. It is independent of the rooted path order except at the two places stated next.

A clean node has size at least $y^{A_{\text{new}}}n$. A t -node frontier has $b_0 = 6$. Otherwise, after grouping small complete siblings, one full chain has positive or negative mass at least $3n/(8t)$. Positive layers have number $L \leq x^{-C_{\text{tree}}}$. The decreasing-size argument gives $t \leq m \leq L$ positive layers of width at least n/m^3 , so their frontier parameter is $b_0 = 3$.

A negative transversal belongs to $\mathcal{A} \cap E_{q-1}$: it is an induced subgraph of the original ambient $F \in \mathcal{A}$, and an earliest-endpoint P_{q-1} could be prefixed by the corresponding negative root to violate RP_q . With $\varepsilon = y^4$, its layer fraction is at most $8/(3t^5) \leq \varepsilon^{4h+4}$. The EH-transversal bridge thus gives a y^4 -restricted set of size at least $y^{8h+d_t+10}n$.

In either frontier/positive branch, $t \leq m \leq x^{-C_{\text{tree}}}$, $b_0 \leq 6$, $\tau = x$. Also $m \geq t \geq 1/y \geq 1/\eta_{\text{new}} \geq m_0$, as Lemma G.1 requires. The order requirement is $n \geq m^{b_0}(m/x)^Q$, which follows from $n \geq x^{-E_{\text{new}}}$. The floating restricted scale satisfies $z \geq (x/m)^H \geq x^{H(C_{\text{tree}}+1)} = x^{c_{\text{new}}}$, and $z \leq 1/m \leq y$. Its width exponent is at most $e_0 + 2A_0 + 7 \leq e_{\text{new}}$.

Blockade lengths satisfy $K \geq m^{1/4} \geq t^{1/4} \geq 1/y$ and $K \leq (m/x)^J \leq x^{-k_{\text{new}}}$. Their width exponent $3b_0 + 9A_0 + d_0 + 3$ is at most d_{new} , because $b_0 \leq 6$. The elementary complete component-grouping outputs have $K = t$ and width n/t^7 , hence also satisfy this common contract.

This proves the abstract full-tree recurrence. $\square$

G.7 The fixed-ambient requirement is essential

When the intended target is a path P_s , a legitimate choice is $\mathcal{A} = \text{Forb}(P_s)$. At every stage, lower rooted Tooth statements, transversal classes, and representative applications must all use this same $\mathcal{A}$. A theorem established only for P_{s-1} -free ambient graphs cannot be reused on a P_s -free graph without a separate argument.

The relative-class premise $\text{EH}(\mathcal{A} \cap E_{q-1})$ is supplied by Section 8 from the lower rooted contract, W_s , and $\text{EH}(P_{s-1})$. Those hypotheses remain separate from the structural frontier recurrence.

For $q = 6$, the lower RP_4 Tooth has a different early-length contract; the specialized proof in Appendix F handles that base. For $q \geq 7$ both required lower orders are at least 5 and the present common-contract induction applies directly.

H Variable-scale local oracles and the sparsification potential

This is an interface theorem conditional on a local oracle. It handles the actual restricted scale z , rather than substituting the coarse parameter y . The lower admissible scale ρ^{2c} , the gain $z^{7/6}$, and the floor estimates are all retained.

Hypothesis H.1 (Variable-scale local oracle). Fix an integer $h \geq 2$, a hereditary class $\mathcal{C}$ of $\overline{P}_h$ -free graphs, and constants $a \geq 4$, $b \geq 1$, $c \geq 2$, $d \geq 1$, $e \geq 4$, $L \geq 1$, $y_0 > 0$. For all $0 < \rho \leq y \leq y_0$, every y^3 -sparse member H of $\mathcal{C}$ of order $N \geq \rho^{-L}$ has at least one of the following outputs.

- (O_1) H is $2y^4$ -sparse. (An oracle that never returns this output is permitted.)
- (O_2) A weakly ρ -semisparse blockade of integer length k with $1/y \leq k \leq \rho^{-c}$ and width at least N/k^d .
- (O_3) Disjoint sets X, Y with $|X| \geq y^a N$ and $|Y| \geq (1 - by)N$, such that Y is directed ρ -sparse to X .
- (O_4) A z^4 -restricted set of size at least $z^e N$, for some $z \in [\rho^c, y]$.

The hypothesis quantifies over the members of $\mathcal{C}$, and not over all $\overline{P}_h$ -free graphs. Since $\mathcal{C}$ is hereditary, it is available in every induced subgraph met below. Output (O_2) requires weak semisparsity only, so a directed ρ -semisparse blockade is an admissible (O_2) output. Output (O_3), in contrast, bounds the degree of each vertex of Y and must be directed. In this paper $h = s$, and $\mathcal{C}$ is either $\text{Forb}(\overline{P}_s)$ (Section 10) or $C_{r,s} = \text{co}(E_r \cap \text{Forb}(P_s))$ (Appendix J).

Choose a fixed c_0 with $0 < c_0 < \min(1/16, \exp(-2b))$, and shrink y_0 if necessary so that

$$y_0 \leq \min(c_0^4/2, c_0^2, c_0/2, 1/(2b), 1/(60h), 1/100). \quad (\text{H.1})$$

Shrinking y_0 preserves the hypothesis, and neither constant depends on ρ or H . Put

$$\begin{aligned} B &= \max(6(e+1), a+4, 4), & R &= 2c, & C_w &= 2c+1, \\ Q &= \max(L+B+1, RB+3, B+c(e+4)+2), \\ D_w &= \max(B+d+1, a+B+2, (B+e+6)/2, RB+3). \end{aligned} \quad (\text{H.2})$$

Real exponents are permitted; increasing them to integers is harmless.

Theorem H.2 (Bootstrap from a fixed sparse start). *Under Hypothesis H.1, with the constants (H.1) and (H.2), every $c_0 y_0^3$ -sparse member G of $\mathcal{C}$ of order $n \geq \rho^{-Q}$, where $0 < \rho < y_0$, has a weakly ρ -semisparse blockade of integer length k with $2 \leq k \leq \rho^{-C_w}$ and width at least n/k^{D_w} .*

Proof. Call y in $[\rho^R, y_0]$ admissible if some induced subgraph F of G is $c_0 y^3$ -sparse and $|F| \geq y^B n$. Every induced subgraph of G belongs to $\mathcal{C}$. The endpoint y_0 is admissible. The minimum y exists: there are finitely many vertex sets and each contributes a closed set of admissible scales. Fix its witness F .

Suppose first $y \geq \rho$. Construct a residual chain inside F . At step $j < 1/y$, the residual H has size at least $(1 - by)^j |F| \geq \exp(-2b) |F| > c_0 |F|$. It is y^3 -sparse and has order $|H| \geq c_0 \rho^B n \geq \rho^{B+1} n \geq \rho^{-L}$, so the local oracle applies. In an O_3 step take X as the next block and Y as the residual. The new block has size at least $c_0 y^a |F| \geq y^{a+2} |F|$, and all later blocks are directed ρ -sparse to it. If $K = \lceil 1/y \rceil$ steps are reached, the resulting blockade has width $y^{a+B+2} n \geq n/K^{a+B+2}$, and $K \leq 2/\rho \leq \rho^{-2}$; it is directed, hence weakly, ρ -semisparse. Every call before this stopping point has $j < K$ and hence $j < 1/y$.

If O_1 occurs, put $y' = c_0 y$. The inequalities $y \leq c_0^4/2$ and $B \geq 1$ give $2y^4 \leq c_0(y')^3$, $|H| \geq c_0 y^B n \geq (y')^B n$. Moreover $y' \geq c_0 \rho \geq \rho^2 \geq \rho^R$ and $y' < y$. This contradicts minimality of y .

In O_2 , since $k \geq 1/y$ and $c_0 \geq y \geq 1/k$, $|H|/k^d \geq c_0 y^B n/k^d \geq n/k^{B+d+1}$. The original output itself is the required weakly ρ -semisparse global blockade; previous residual blocks need not be retained.

For O_4 , write T for its output and z for its actual scale. Thus $|T| \geq c_0 y^B z^e n \geq c_0 z^{B+e} n$. If $G[T]$ is z^4 -sparse, put $y' = z^{7/6}$. Since $z \leq y_0 \leq c_0^2$, $z^4 \leq c_0 z^{7/2} = c_0 (y')^3$. As $c_0 \geq z$ and $B \geq 6(e+1)$, $|T| \geq z^{B+e+1} n \geq z^{7B/6} n = (y')^B n$. The scale satisfies $\rho^R \leq \rho^{7c/6} \leq y' < y$. This contradicts minimality. This is exactly why z may float far below y without invalidating the potential argument.

If the complement of $G[T]$ is z^4 -sparse, apply Contract 2.6 to it, with the path P_h and the parameter z^2 . The hypotheses hold. First, $h \geq 2$ and $z^2 \leq z \leq y_0 \leq 1/(60h)$. Second, the complement of $G[T]$ is P_h -free, because $G \in \mathcal{C}$ is $\overline{P_h}$ -free, and it is $(z^2)^2$ -sparse. Third, since $y \geq \rho$ and $z \geq \rho^c$,

$$z^4 |T| \geq \rho^{B+c(e+4)+1} n \geq \rho^{-1} > 2, \quad (\text{H.3})$$

so $|T| \geq (z^2)^{-2}$, which is the order hypothesis of the contract. Retain $K = \lceil z^{-2} \rceil$ blocks of the anticomplete output; they are pairwise complete in G , and their width is at least $\lfloor z^4 |T| \rfloor \geq 2$. By (H.3) the floor loses at most a factor 2. Because $z \leq c_0/2$ implies $z^2 \leq c_0/2$, width at least $(c_0/2) z^{B+e+4} n \geq z^{B+e+6} n \geq n/K^{(B+e+6)/2}$. Also $K \leq 2\rho^{-2c} \leq \rho^{-(2c+1)}$. This is another valid output.

It remains to handle the case in which the minimum y is below ρ . Now $|F| \geq \rho^{RB} n$, and $G[F]$ has maximum degree at most $c_0 \rho^3 |F|$. Put $K = \lceil 1/\rho \rceil$ and partition F into K equal floor-size blocks, discarding the remainder. The order condition $n \geq \rho^{-(RB+3)}$ gives $\rho |F| \geq \rho^{-2} > 4$. Thus every block has size at least $\rho |F|/4$. The inherited maximum degree is at most $\rho \cdot$ this block size, since $4c_0 \rho \leq 1$. The blockade is directed ρ -sparse in both directions, hence weakly ρ -semisparse, and has width $\rho^{RB+1} n/4 \geq \rho^{RB+3} n \geq n/K^{RB+3}$. Its length is at most ρ^{-2} . This proves the warm conclusion. $\square$

H.1 A fixed Rödl start and generalized niceness

Let $G \in \mathcal{C}$ have order n . Contract 2.5, for $\overline{P_h}$ -free graphs at precision $c_0 y_0^3$, supplies a $c_0 y_0^3$ -restricted induced subgraph of order at least δn , where $\delta > 0$ is the fixed Rödl fraction; it is a member of $\mathcal{C}$. In the G -sparse orientation use Theorem H.2, the warm conclusion. In the complementary sparse orientation use Contract 2.6 with the fixed parameter $1/(60h)$. This is legitimate: $c_0 y_0^3 \leq y_0^2 \leq (60h)^{-2}$, and the order hypothesis $\delta n \geq (60h)^2$ holds once $n \geq (60h)^2/\delta$, which is one of the fixed minimum orders absorbed below. Two blocks of its output form a complete blockade of length 2 and width at least $\lfloor \delta n/(60h)^2 \rfloor \geq 1$. Absorb the fixed Rödl and flooring constants by increasing a fixed exponent D_0 and shrinking the

allowed fixed upper cutoff for ρ . There result fixed D_0, C such that $n \geq \rho^{-D_0}$ implies a weakly ρ -semispars blockade of integer length K with $2 \leq K \leq \rho^{-C}$ and width at least n/K^{D_0} . For explicit absorption, if the Rödl fraction is δ , use $L_\delta = \lceil \log_2(1/\delta) \rceil$; $n \geq \rho^{-(Q+L_\delta)}$ ensures that its subset meets the warm order bound, and a width exponent increased by L_δ absorbs its size fraction. Increase D_0 further for the fixed complete-blockade branch and its fixed minimum order.

Lemma I.1 then applies verbatim. Choose $S \geq 8D_0 + 8$ with 2^{-S} below the target cutoff, and use $\rho = \varepsilon^S$. It gives generalized niceness with $M = \max(4D_0, 2D_0 + SD_0)$, $N = \max(3, 4D_0, 2D_0 + CSD_0)$. The large-graph output has $\lceil 1/\varepsilon \rceil$ blocks of width $\varepsilon^N n$, every pair complete or weakly ε^8 -sparse; the small-graph output is $\text{hom}(G) \geq \varepsilon^M n$. The layout proof uses only the displayed polynomial length cap and weak semisparsity.

The bootstrap uses only the local oracle and the sparse-path extraction contract for P_h ; it does not assume $\text{EH}(P_h)$.

I The arbitrary-pattern layout argument

The pattern may be any graph. Only pairs declared complete are required to be complete; the counted errors are edges across pattern nonedges. The terminal product proof will later use uniform complete or anticomplete blockades, a distinct requirement.

A hereditary class $\mathcal{C}$ satisfies *generalized niceness* if there are constants $M, N > 0$ such that, for every member G and every $0 < \varepsilon < 1/2$, either $\text{hom}(G) \geq \varepsilon^M |G|$, or G has a blockade of length $\lceil 1/\varepsilon \rceil$ and width at least $\varepsilon^N |G|$ in which every pair is complete or weakly ε^8 -sparse.

Lemma I.1 (Arbitrary-pattern layout). *Let $\mathcal{C}$ be a hereditary class. Suppose fixed $D_0 \geq 1$, $C > 0$, $L > 0$ and $\rho_0 > 0$ satisfy: for every $0 < \rho < \rho_0$ and every member F of $\mathcal{C}$ with $|F| \geq \rho^{-L}$, there is a weakly ρ -semispars blockade in F of integer length K with $2 \leq K \leq \rho^{-C}$ and width at least $|F|/K^{D_0}$. Then $\mathcal{C}$ satisfies generalized niceness. This remains true for coarse-scale $K = 2$; no lower bound $K \geq 1/\varepsilon$ is needed at the input.*

Proof. Choose a fixed $S \geq 8D_0 + 8$ with $2^{-S} < \rho_0$. For $0 < \varepsilon < 1/2$ set $\rho = \varepsilon^S$, $\ell = \lceil 1/\varepsilon \rceil$, $\sigma = \varepsilon^{4D_0}$, $M = \max(4D_0, 2D_0 + SL)$. First assume $n \geq \varepsilon^{-M}$. A layout consists of disjoint blocks A_i and an arbitrary pattern J ; a pattern edge means the corresponding actual block pair is complete. Wrong pairs are actual G -edges between blocks whose pattern pair is a nonedge. An unordered ambient pair is decided if and only if its two vertices do not lie together in one current block: equivalently, at least one endpoint is outside the union, or the endpoints lie in different blocks. Pairs within one block are undecided. Choose a layout maximizing its number of blocks subject to: $|A_i| \geq \sigma n$; $\sum_i |A_i|^{1/D_0} \geq n^{1/D_0}$; wrong-pair count at most ρ ·decided-pair count. The one-block layout $A_1 = G$ exists. No cograph restriction is placed on J .

If there are at least ℓ blocks, choose ℓ of them. On every pattern nonedge, its edge count is at most $\rho \cdot n^2/2$. Since both blocks have size at least σn , their density is at most $\rho/(2\sigma^2) \leq \varepsilon^8$. Pattern edges are genuinely complete. This gives the required blockade, with complete or weakly ε^8 -sparse pairs and width at least $\varepsilon^{4D_0} n$.

Otherwise there are fewer than $\ell \leq 2/\varepsilon$ blocks, so one block A has size at least $(\varepsilon/2)^{D_0} n \geq \varepsilon^{2D_0} n$. In particular $|A| \geq \rho^{-L}$, and the target interface applies inside A . If its returned K is at least ℓ , choose ℓ of its blocks directly. Their width is at least $|A|/K^{D_0} \geq \varepsilon^{2D_0} \cdot \rho^{CD_0} n = \varepsilon^{2D_0+SCD_0} n$, and their precision ρ is at most ε^8 . If $K < \ell$, replace A in the layout by all K returned blocks. Each has size at least $\varepsilon^{4D_0} n$, and their $1/D_0$ -power mass is at least $|A|^{1/D_0}$. The new pattern substitutes the returned complete/sparse pattern at A . Old complete pairs

remain complete. Old wrong pairs can only disappear; new wrong pairs occur only between children of A , on non-complete returned pairs, and by weak ρ -sparsity of those pairs they number at most ρ times the cross-pair count. No bound on individual degrees is used here. Those cross pairs are newly decided; discarded vertices only add more decided pairs. Thus the invariant is preserved, contradicting maximality because $K \geq 2$.

Therefore every $n \geq \varepsilon^{-M}$ graph has an $(\varepsilon^{-1}, \varepsilon^N n)$ blockade with complete or weakly ε^8 -sparse pairs, where $N = \max(3, 4D_0, 2D_0 + SCD_0)$. For $n < \varepsilon^{-M}$, a singleton supplies the homogeneous-set alternative, since $\varepsilon^M n < 1$. This is generalized niceness as defined before the lemma, with these M and N . $\square$

J Ordered tails and the relative ordered-class EH interface

The input is the quantified $T_r^{(s)}$ contract of Section 7. Throughout this appendix A, e, C, K, D, E and y_* denote its constants $A_r, e_r, c_r, k_r, d_r, E_r^*$ and η_r . The output is Hypothesis H.1 for the hereditary class $\mathcal{C} = C_{r,s} = \text{co}(E_r \cap \text{Forb}(P_s))$ and $h = s$. Its blockade output (O_2) need only be weakly semisparsely, whereas its clean-root output (O_3) must be directed. The terminal proof invoked below is expanded in Appendix K.

Let G belong to $C_{r,s}$, be y^3 -sparse, and have order n , with $0 < \rho \leq y \leq \min(y_*, 1/100)$, $n \geq \rho^{-(E+2)}$. In an E_r order of $\overline{G}$, take the last $m = \lceil yn \rceil$ vertices as Y and take the prefix as U . Then $yn \leq m \leq 2yn$, $m \geq \rho^{-(E+1)} \geq \rho^{-E}$. Every root u precedes Y , so no induced P_r with endpoint u occurs in $\overline{G}[Y \cup \{u\}]$. The ambient complement is P_s -free. Apply $T_r^{(s)}$ with $x = \rho$ and coarse parameter y .

T_1 yields a set X of size at least $y^{A+1}n$. At most y^3n roots are full to X , by the degree of any fixed vertex of nonempty X . All other roots form Z of size at least $n - 2yn - y^3n \geq (1 - 3y)n$, and Z is directed $\rho/4$ -sparse to X .

T_2 yields its actual scale z in $[\rho^C, y]$ and size $z^e m \geq z^e yn \geq z^{e+1}n$. The direction of the last inequality uses $z \leq y$.

T_3 yields $k \geq 1/y$ and width $m/k^D \geq yn/k^D \geq n/k^{D+1}$. Its cap remains $k \leq \rho^{-K}$. This blockade is directed ρ -semisparsely, hence weakly ρ -semisparsely, and is an (O_2) output.

Thus Hypothesis H.1 holds for $\mathcal{C} = C_{r,s}$, with y_* as the initial cutoff y_0 and with constants $a = \max(4, A + 1)$, $b = 3$, $c = \max(2, C, K)$, $d = D + 1$, $e_{\text{or}} = \max(4, e + 1)$, $L = E + 2$. There is no O_1 output, which the bootstrap explicitly permits. The same argument works in every induced member of $C_{r,s}$.

J.1 Explicit bootstrap and layout parameters

Apply Appendix H with $h = s$ and the hereditary class $\mathcal{C} = C_{r,s}$. For example take $c_0 = 2^{-12}$; then $c_0 < \exp(-6)$. Choose a fixed positive y_0 with $y_0 \leq \min(y_*, c_0^4/2, c_0^2/2, c_0/2, 1/6, 1/(60s), 1/100)$.

As in (H.2), with e_{or} in place of e , put $B = \max(6(e_{\text{or}} + 1), a + 4, 4)$, $R = 2c$, $Q = \max(L + B + 1, RB + 3, B + c(e_{\text{or}} + 4) + 2)$, $D_w = \max(B + d + 1, a + B + 2, (B + e_{\text{or}} + 6)/2, RB + 3)$, $C_w = 2c + 1$. Then the warm conclusion of Theorem H.2 holds for all $n \geq \rho^{-Q}$: a weakly ρ -semisparsely blockade of integer length k with $2 \leq k \leq \rho^{-C_w}$ and width at least n/k^{D_w} .

Let δ_0 be the fixed Rödl fraction for $\overline{P_s}$ at precision $c_0 \cdot y_0^3$. Write $L_\delta = \lceil \log_2(1/\delta_0) \rceil$, $J_s = \lceil \log_2(2(60s)^2) \rceil + 2$, $D_0 = \lceil \max(Q, D_w) \rceil + L_\delta + J_s$, $\rho_0 = y_0/2$. The Rödl subset in an $n \geq \rho^{-D_0}$ graph has order at least ρ^{-Q} . The complementary orientation uses Contract 2.6 for P_s at parameter $1/(60s)$, keeping two complete blocks. The contract applies: $c_0 y_0^3 \leq y_0^2 \leq (60s)^{-2}$,

and the Rödl subset has order at least $\rho^{-Q} > (60s)^2$, because $\rho < y_0 \leq 1/(60s)$ and $Q \geq 3$. The exponent J_s absorbs its floor and width constant. Hence the global target interface is a weakly ρ -semisparse blockade of integer length k with $2 \leq k \leq \rho^{-C_w}$ and width at least n/k^{D_0} .

Choose integer $S \geq 8D_0 + 8$ with $2^{-S} < \rho_0$, and define $M = \max(4D_0, 2D_0 + SD_0)$, $N = \max(3, 4D_0, 2D_0 + C_w SD_0)$. The proved layout lemma gives, for every member G and $0 < \varepsilon < 1/2$, either $\text{hom}(G) \geq \varepsilon^M |G|$ or a complete/weak- ε^8 -sparse blockade of length $\lceil 1/\varepsilon \rceil$ and width $\varepsilon^N |G|$. This proves generalized niceness with every scale and order quantified.

J.2 Sufficient conditions for EH of the ordered class

Assume additionally: (W_s) strong wonderfulness for $\overline{P_s}$ -free graphs with some exponent $a_s \geq 6$, valid for equal-sized blocks without anticonnectivity; (V_s) $\text{EH}(P_{s-1})$, hence normalized P_{s-1} virality with exponent $v \geq 1$.

The general path leaf-fibre proof gives, in every y -sparse P_s -free graph, either a y^4 -restricted set of size at least $y^{4(s-1)v+4}n$, or an anticomplete pair X, Y with X at least this size and $|Y| \geq (1 - (s-1)y)n$. Indeed fix the last $s-2$ vertices of the abundant ordered P_{s-1} copies. Their endpoint fibre X lies in the neighbourhood of the first fixed core vertex. Delete the core and all its neighbourhoods to get Y . An edge from Y to X would extend to an induced P_s . For $n < (s-2)/y$, a singleton handles the restricted outcome.

As in (K.1), set $P = \max(M, N + 3)$, $A_{\text{term}} = \max(a_s P, 4(s-1)v + 4, 4)$, $c_T = y_T = 2^{-4s}$, $B_{\text{term}} = A_{\text{term}} + 2$, $d_T = 2A_{\text{term}} + 4$. Lemma K.2 supplies a three-output oracle for y -restricted graphs in $C_{r,s}$: $\text{hom} \geq y^{A_{\text{term}}}n$; or a y^4 -restricted set of that size; or an orientation-controlled pure pair with small side of that size and large side $(1 - (s-1)y)n$. The residual chain survives at least $\exp(-2(s-1)) > c_T$, so the compact-scale and product proofs apply without further changes.

If δ_T is the fixed Rödl fraction for $\overline{P_s}$ at precision $c_T y_T$, put $D_{\text{term}} = d_T + \lceil \log_2(1/\delta_T) \rceil$. Then every member of $E_r \cap \text{Forb}(P_s)$ has a homogeneous set of size at least $n^{1/(8D_{\text{term}})}$.

Only virality of the single forbidden graph P_{s-1} is used. No EH-to-virality statement for the hereditary class E_r is asserted.

J.3 How to use this in a rooted-path induction

A negative layer sequence in an RP_q construction has transversals in E_{q-1} : the negative witness prefixes any rooted P_{q-1} from the earliest selected vertex to form a forbidden rooted P_q . If the ambient complement is P_s -free, its transversals actually belong to $E_{q-1} \cap \text{Forb}(P_s)$.

Therefore, for a fixed final forbidden path P_s , a potential rooted induction has the following acyclic pattern: $T_{q-1}^{(s)}, W_s, \text{EH}(P_{s-1}) \implies \text{EH}(E_{q-1} \cap \text{Forb}(P_s))$, and the latter is the negative-transversal input for $T_q^{(s)}$.

The positive/frontier branch separately uses lower rooted Tooth theorems at $q-1$ for the first/canonical purification and at $q-2$ for the offending-pair purification. Those lower theorems must all be proved with the same ambient superscript s . The ordered-tail theorem above supplies the required negative-class EH input, not the entire positive/frontier induction.

K Leaf fibres, cleanup and the terminal product argument

Fix $s \geq 7$. Assume hereditary niceness, W_s , $\text{EH}(P_{s-1})$, and fixed-precision Rödl. We work in a hereditary subclass of the $\overline{P_s}$ -free graphs.

K.1 Normalization of the single-path viral input

Let $h = s - 1$. The viral consequence of $\text{EH}(P_h)$ can be normalized, with a fixed $v \geq 1$, to say that for every $0 < \varepsilon < 1/2$ and every graph J of order n , either there is an ε -restricted set of order at least $\varepsilon^v n$, or there are at least $\varepsilon^{hv} n^h$ ordered induced P_h embeddings. This is only virality of the single graph P_h , not of the class E_r .

We use the following maximum-degree normalization. Enlarge a constant $c_0 \geq 1$ so that the usual edge-density viral statement has output-size exponent at most c_0 and copy-count exponent at most hc_0 . Apply it at $\varepsilon/4$. In a set with density at most $\varepsilon/4$ in one orientation, remove vertices having degree greater than $\varepsilon|S|/2$. Fewer than half the vertices are removed. The retained T has maximum degree at most $\varepsilon|T|$, and

$$|T| \geq (\varepsilon/4)^{c_0} n/2 \geq \varepsilon^{3c_0+1} n.$$

In the copy alternative,

$$(\varepsilon/4)^{hc_0} n^h \geq \varepsilon^{3hc_0} n^h \geq \varepsilon^{h(3c_0+1)} n^h.$$

Thus $v = 3c_0 + 1$ suffices. Ordered embeddings are at least as numerous as the unlabelled copies used by the cited statement [3].

Lemma K.1 (Path leaf-fibre dichotomy). *For $0 < y < 1/2$ and a y -sparse P_s -free graph J of order n , either there is a y^4 -restricted set of order at least $y^{4(s-1)v+4} n$, or there are disjoint anticomplete sets X, Y with*

$$|X| \geq y^{4(s-1)v+4} n, \quad |Y| \geq (1 - (s-1)y)n.$$

Proof. If $n < (s-2)/y$, a singleton supplies the first output, since $(s-2)y^{4(s-1)v+3} < 1$ for $s \geq 7$, $v \geq 1$, and $y < 1/2$. Otherwise apply the normalized viral statement at $\varepsilon = y^4$. Its restricted output is large enough. In the other case there are at least $y^{4(s-1)v} n^{s-1}$ ordered path embeddings. Fix the ordered tail $(p_2, \dots, p_{s-1})$ of a most frequent such embedding. At most n^{s-2} tails occur, so its endpoint fibre X has order at least $y^{4(s-1)v} n$. Every $x \in X$ is adjacent to p_2 and nonadjacent to all later tail vertices.

Let Y omit the $s-2$ tail vertices and all their neighbourhoods. Then

$$|Y| \geq n - (s-2) - (s-2)yn \geq (1 - (s-1)y)n,$$

using $n \geq (s-2)/y$. Since $X \subseteq N(p_2)$, the two sets are disjoint. An edge ux , with $u \in Y$ and $x \in X$, would extend the path to the induced P_s , namely $u, x, p_2, \dots, p_{s-1}$. Hence they are anticomplete. $\square$

K.2 Cleanup from weak sparsity to equal-width directed sparsity

Assume niceness supplies constants $N \geq 3$, $M > 0$: for every $0 < \varepsilon < 1/2$, either $\text{hom}(G) \geq \varepsilon^M n$, or there is a blockade of length $\ell = \lceil 1/\varepsilon \rceil$, width at least $\varepsilon^N n$, whose pairs are complete or weakly ε^8 -sparse. Set

$$P = \max(M, N + 3), \quad A = \max(a_s P, 4(s-1)v + 4, 4), \quad B = A + 2, \quad (\text{K.1})$$

and $c = y_0 = 2^{-4s}$, $d = 2A + 4 = 2B$. In Section 11 and Appendix J these constants are written A_{term} , B_{term} , $c_T = y_T$ and d_T . For the cleanup put $\varepsilon = y^{a_s}$, with $0 < y < 1/2$, and suppose $n > \varepsilon^{-P}$. Let $m = \lceil \varepsilon^N n \rceil$. Then

$$\varepsilon^N n \geq \varepsilon^{-3}, \quad \varepsilon^N n \leq m \leq 2\varepsilon^N n.$$

Each original block contains an m -subset. Choose these subsets X_i independently and uniformly. On every noncomplete pair the expected edge count is at most $\varepsilon^8 m^2$; the total over all such pairs has expectation at most

$$\binom{\ell}{2} \varepsilon^8 m^2 \leq 2\varepsilon^6 m^2 \leq \varepsilon^5 m^2.$$

Fix an outcome no larger than this bound. In particular every originally sparse pair contains at most $\varepsilon^5 m^2$ edges. Delete from X_i all vertices having more than $\varepsilon^3 m$ neighbours in any relevant X_j . Each other block causes fewer than $\varepsilon^2 m$ deletions, so the total in each X_i is at most $2\varepsilon m$. The survivors D_i have order at least $m/2$.

Choose arbitrary subsets $B_i \subseteq D_i$ of the common order $w = \lceil \varepsilon^2 m \rceil$. This is possible because $\varepsilon^2 m \geq 1$ and

$$w \leq 2\varepsilon^2 m \leq \varepsilon m/4 \leq |D_i|.$$

Every originally sparse pair is now sparse in both directions: its vertex degrees are at most $\varepsilon^3 m \leq \varepsilon w$. Complete pairs stay complete. Moreover,

$$w \geq \varepsilon^{N+2} n, \quad \ell w \leq 4\varepsilon m \leq 8\varepsilon^{N+1} n \leq y^2 n.$$

This is the exact input to W_s . The cleanup neither needs anticonnectivity nor turns a mixed pattern into a uniform anticomplete pattern.

Lemma K.2 (An orientation-controlled three-output lemma). *Every y -restricted $\overline{P}_s$ -free graph G has one of:*

- (Q1) $\text{hom}(G) \geq y^A n$;
- (Q2) a y^A -restricted set of order at least $y^A n$;
- (Q3) disjoint pure sets X, Y with $|X| \geq y^A n$ and $|Y| \geq (1 - (s - 1)y)n$.

In Q3 the pair is anticomplete if the declared sparse orientation is G , and complete if it is F .

Proof. When F is the declared sparse graph, apply Lemma K.1; its anticomplete pair becomes complete in G . When G is sparse, use $\varepsilon = y^{a_s}$. For $n \leq \varepsilon^{-P}$, a singleton gives Q1. For larger n , the homogeneous niceness output gives Q1 because $A \geq a_s M$. Otherwise perform the cleanup and apply W_s to the B_i . Its restricted output has order at least $w \geq \varepsilon^{N+2} n \geq y^A n$, giving Q2.

In the remaining case a block B_i has at most yn outside vertices with degree strictly between zero and $w/2$. Since $\Delta(G) \leq yn$, at most $2yn$ outside vertices have at least $w/2$ neighbours in B_i , by counting edges incident with this block. The entire blockade occupies at most $y^2 n$ vertices. Thus at least $(1 - 3y - y^2)n \geq (1 - 4y)n \geq (1 - (s - 1)y)n$ vertices outside it have no neighbours in B_i . Together with $X = B_i$, these give the anticomplete Q3 output. $\square$

K.3 The compact scale potential

Suppose first that G is cy_0 -restricted and $0 < x < y_0$. Call a scale $y \in [x^2, y_0]$ admissible if there is an induced H that is cy -restricted and has order at least $y^B n$. The endpoint y_0 is

admissible. There are finitely many vertex subsets and each gives a closed set of scales, so a minimum exists. Fix a witness H . If $y < x$, this witness is x -restricted and has order at least $x^{2B}n = x^d n$.

Assume $y \geq x$, and fix one restricted orientation of H . Build a residual chain. At every call before $K = \lceil 1/y \rceil$ steps, the step index $j < K$ satisfies $j < 1/y$, so earlier Q3 outputs leave a residual of order at least

$$(1 - (s-1)y)^j |H| \geq e^{-2(s-1)} |H| > c|H|.$$

Here $(s-1)y \leq 1/2$ and $\log(1 - (s-1)y) \geq -2(s-1)y$, because $y \leq 2^{-4s}$. Every residual is y -restricted in the same orientation. A Q1 output gives

$$\text{hom}(G) \geq cy^{A+B}n \geq x^{A+B+1}n \geq x^d n.$$

A Q2 output gives a set T of order at least $cy^{A+B}n$ and restriction y^4 . Put $y' = y^2$. Then $x^2 \leq y' < y$, $y^4 \leq c(y')$, and

$$|T| \geq cy^{A+B}n \geq y^{2B}n = (y')^B n,$$

using $B = A + 2$ and $c \geq y^2$. This contradicts minimality. If only Q3 occurs, take its small side as the next block and its large side as the new residual. The fixed orientation makes all completed blocks uniformly complete or uniformly anticomplete. After K steps their width is at least

$$y^{A+2}|H| \geq y^{A+B+2}n = y^d n \geq n/K^d.$$

We therefore obtain a global trichotomy from the fixed restricted start: an x -restricted set of order at least $x^d n$, a uniform complete or anticomplete blockade of integer length $K \geq 2$ and width at least n/K^d , or $\text{hom}(G) \geq x^d n$.

K.4 Removing the fixed start

Let δ_T be the fixed Rödl fraction at precision cy_0 . Set

$$L_T = \lceil \log_2(1/\delta_T) \rceil, \quad D_s = d + L_T. \quad (\text{K.2})$$

For $x < y_0$, $x^{L_T} \leq \delta_T$ and $\delta_T/K^d \geq K^{-(d+L_T)}$ for $K \geq 2$. Applying the previous argument to a Rödl subset yields the same trichotomy with exponent D_s .

For $x \in [y_0, 1/2)$, the fixed subset is already x -restricted because $cy_0 \leq y_0 \leq x$. Its order is sufficient since $x^{D_s} \leq x^{L_T} \leq 2^{-L_T} \leq \delta_T$. This also covers $L_T = 0$, when $\delta_T = 1$. Thus the trichotomy holds at every $0 < x < 1/2$, with no precision cutoff depending on δ_T and no excluded small-order case.

Lemma K.3 (Uniform-blockade product criterion). *If a hereditary class satisfies this trichotomy for some $D \geq 1$, every n -vertex member satisfies $\text{hom}(G) \geq n^{1/(8D)}$.*

Proof. Put $Q(G) = \alpha(G)\omega(G)$ and $\tau = 1/(4D)$. For a uniformly complete blockade $B_1, \dots, B_k$, $\omega(G) \geq \sum_i \omega(G[B_i])$ and $\alpha(G) \geq \max_i \alpha(G[B_i])$. Their product is therefore at least $\sum_i Q(G[B_i])$. The anticomplete case is symmetric. This superadditivity is asserted only for the two uniform types.

Choose a minimum-order counterexample to $Q(G) \geq n^\tau$. There is none at $n = 1$, or at $2 \leq n \leq 2^{4D}$, since $Q(G) \geq 2$. Hence $n > 2^{4D}$. Use $x = n^{-1/(2D)} < 1/4$. The homogeneous output gives $\text{hom}(G) \geq \sqrt{n} \geq n^\tau$, a contradiction. A restricted set of order $m \geq \sqrt{n} \geq 1/x$

has, by the greedy independence bound in its sparse orientation, a homogeneous set of order at least

$$\frac{m}{xm+1} \geq \frac{1}{2x} = \frac{1}{2}n^{1/(2D)} > n^{1/(4D)}.$$

This is also a contradiction. In the blockade output, every nonempty block is proper. Heredity and minimality give $Q(G[B_i]) \geq |B_i|^\tau$, so

$$Q(G) \geq \sum_i |B_i|^\tau \geq k(n/k^D)^\tau = n^\tau k^{3/4} > n^\tau.$$

Thus $Q(G) \geq n^\tau$ for all members, and $\text{hom}(G)^2 \geq Q(G)$ gives the exponent $1/(8D)$. $\square$

References

- [1] machine-qed. *The Erdős–Hajnal property for the six-vertex path*. Unpublished manuscript, repository version v1.0–proof, 27 September 2026, Theorem A. Not refereed.
<https://github.com/machine-qed/erdos-hajnal-six-vertex-path/tree/v1.0-proof>.
- [2] Tung Nguyen, Alex Scott and Paul Seymour. *Induced subgraph density. V. All paths approach Erdős–Hajnal*. arXiv:2307.15032v2.
<https://arxiv.org/html/2307.15032v2>.
- [3] Matija Bucić, Jacob Fox and Huy Tuan Pham. *Equivalence between Erdős–Hajnal and polynomial Rödl and Nikiforov conjectures*. arXiv:2403.08303v2, 19 April 2024.
<https://arxiv.org/html/2403.08303v2>.
- [4] Shenwei Huang, Yiao Ju and Yidong Zhou. *Erdős–Hajnal conjecture beyond five-vertex graphs*. arXiv:2606.06258v3, 31 August 2026, Lemma 2.10.
<https://arxiv.org/html/2606.06258v3>.
- [5] Ryan B. Hayward. *Weakly triangulated graphs*. Journal of Combinatorial Theory, Series B 39(3) (1985), 200–208.
[https://doi.org/10.1016/0095-8956\(85\)90050-4](https://doi.org/10.1016/0095-8956(85)90050-4).
- [6] Tung Nguyen, Alex Scott and Paul Seymour. *Induced subgraph density. VII. The five-vertex path*. arXiv:2312.15333v3, 23 February 2026.
<https://arxiv.org/abs/2312.15333>.
- [7] V. Rödl. *On universality of graphs with uniformly distributed edges*. Discrete Mathematics 59 (1986), 125–134.