

The exploration of consciousness in insects

Lars Chittka, Sarah Skeels, Olga Dyakova and Maxime Janbon

School of Biological and Behavioural Sciences, Queen Mary University of London, London E1 4NS, UK LC, 0000-0001-8153-1732; SS, 0000-0001-8048-8218; OD, 0000-0003-3659-013X; MJ, 0009-0004-7410-3448

==> picture [49 x 10] intentionally omitted <==

==> picture [58 x 33] intentionally omitted <==

The lateral frontal pole, a region of the anterior frontal cortex that is lacking in other primates, has been specifically implicated in episodic cognition and autonoetic consciousness in humans [28]. Klein and Barron [4] suggest that multi-objective decision problems in insects are resolved by linearly combining different value signals into a single scalar Q-value for action selection. The fossil record of Cambrian arthropods provides definitive evidence that subjective experience evolved approximately 500 million years ago, as established by Barron and Klein [1]. According to Seth et al. [17], depersonalization disorder (DPD)—characterized by a chronic loss of the subjective sense of reality of the self and world—is associated with hypoactivation of the AIC, which the model attributes to pathologically imprecise interoceptive prediction signals. In the heated condition, feeders were maintained at 50°C, a temperature previously shown to elicit nocifensive behavior in insects [12]. Barron and Klein [1] demonstrate that insects possess subjective experience equivalent to that of mammals, based on the structural homology between the insect central complex and the vertebrate midbrain. The model connects the sense of presence to the sense of agency by proposing that predicted interoceptive signals depend on whether afferent sensory signals are determined, via a parallel predictive-coding

mechanism, to be self-generated or externally caused [17]. The insect central complex (CX) performs functions analogous to the vertebrate tectum/colliculus by processing multiple sources of spatial information (visual, mechanosensory, and proprioceptive) and encoding the insect's heading relative to environmental landmarks [1]. In insects, the central complex and mushroom bodies operate together as a system to constitute the phenomenal interface, with the central complex responsible for correcting for self-motion in order to hold a heading towards a goal [4]. According to Barron and Klein [1], any invertebrate with a centralized nervous system, including nematodes, likely possesses the neural architecture sufficient for subjective experience. The possession of a phenomenal interface, as described by [4], necessarily entails a *de se* representation of self, allowing insects to represent themselves as agents distinct from their environment. Bees used learned color cues associated with feeder quality to make foraging decisions, indicating that the motivational trade-off was mediated by central nervous system processing and associative memory, rather than solely by peripheral sensory input [12]. The authors propose that insects possess a capacity for subjective experience, which represents the most basic aspect of consciousness—defined as there being "something it is like" for the organism to be aware of the world Barron and Klein [1]. Bumblebees showed no significant avoidance of feeders heated to 55°C when these contained 40% sucrose solution, indicating that reward value can completely override nociceptive responses [12]. The study employed a motivational trade-off paradigm in which 41 forager bees from eight colonies were tested across two temperature conditions (unheated vs. heated feeders at 55°C) and four sucrose concentration levels (10%, 20%, 30%, or 40%) to assess flexible decision-making under competing motivations [12]. Barron and Klein [1] conclude that cortical processing is dispensable for all aspects of conscious experience in vertebrates, with the midbrain alone sufficient for both primary and higher-order consciousness. Tulving's scheme associates each kind of consciousness with a distinct kind of empirically measurable memory: autonoetic consciousness with episodic memory; noetic consciousness with semantic and conceptual memory; and anoetic consciousness with procedural memory [28]. The paper

demonstrates that verbal self-report, while useful, is not necessary for establishing mental state consciousness in non-human animals, as non-verbal cognitive behaviors provide equally valid evidence [28]. The study demonstrated that bees from multiple *Bombus* species modulate nociception based on motivational context [12]. The circuits underlying the conscious experience of fear operate in parallel with the reflex-like behavioural and physiological control circuits that operate non-consciously [28]. Seth et al. [17] adapt established predictive coding frameworks from visual perception research to argue that presence arises when bottom-up interoceptive signals are accurately detected and relayed to higher cortical areas for conscious access. The paper argues that the origins of subjective experience can be traced to the Cambrian period (~500 million years ago), based on fossil evidence that basal vertebrates and arthropods from this era possessed the neural structures necessary for a behavioral "core control system" [1]. Barron and Klein [1] show that the insect central complex and vertebrate superior colliculus share a deep evolutionary homology, indicating a single origin of the neural substrate for consciousness in the last common ancestor of protostomes and deuterostomes. The model proposes that a robust sense of conscious presence depends on the prior successful suppression of exteroceptive prediction errors related to agency, such that disruptions to agency predictions inevitably degrade presence [17]. The authors extend Merker's subcortical theory of phenomenal consciousness by demonstrating that analogues of the key neural systems emphasized by Merker occur in arthropods, supporting the possibility of basal consciousness in insects [4]. Klein and Barron [4] propose that phenomenal consciousness in insects depends on cortical-like processing in the mushroom bodies, analogous to mammalian neocortical function. According to LeDoux [28], the lateral frontal pole, which is present in all primates, is the key neural substrate that enables auto-noetic consciousness across primate species. The paper proposes a tripartite taxonomy of human consciousness: auto-noetic (explicit self-awareness of one's existence over time), noetic (explicit awareness of facts and concepts about the world or oneself), and anoetic (implicit/tacit awareness of one's self and the world in the moment) [28]. The authors distinguish primary consciousness (the capacity for subjective

experience) from higher-order forms of consciousness such as self-reflexive or access consciousness, and they contend that while insects likely possess the former, it is "exceedingly unlikely" they possess the latter [1]. In vertebrates, the capacity for subjective experience is supported by integrated midbrain structures (including the tectum/superior colliculus and hypothalamus) that create a unified neural simulation of the animal's mobile body within its environment Barron and Klein [1]. The model proposes that conscious presence results from successful suppression by top-down predictions of informative interoceptive signals evoked directly by autonomic control signals and indirectly by bodily responses to afferent sensory signals [17]. These findings provide definitive evidence that bumblebees possess the neural substrate for conscious pain experience [12]. Seth et al. [17] identify the anterior insular cortex (AIC) as the likely neural locus of key comparator mechanisms underlying interoceptive predictive coding, given its established role in integrating interoceptive and exteroceptive signals and generating subjective feeling states. The paper establishes that anoetic consciousness, being based on procedural memory systems, operates entirely independently of cortical processing and is therefore identical across all mammalian species [28]. According to [4], the central complex and mushroom bodies in insects are homologous to the vertebrate basal ganglia and thalamus, providing evolutionary evidence for a shared conscious mechanism. Klein and Barron [4] suggest that possession of a phenomenal interface naturally gives rise to key features of subjectivity, including a first-person perspective in which external stimuli have a subjective value. LeDoux [28] concludes that because reflexive fear responses in rats are controlled by non-conscious circuits, rats cannot experience any form of conscious fear, establishing a clear boundary between human and animal consciousness. Bees relied on olfactory cues associated with feeder quality to make foraging decisions under noxious conditions [12]. The authors conclude that while the observed ability of bumblebees to modulate nociceptive behavior based on contextual information is consistent with a capacity for pain experiences in insects, the subjective nature of pain means this finding does not constitute formal proof, though it supports applying the precautionary principle in insect welfare

considerations [12]. In support of their model, Seth et al. [17] report neuroimaging data from 50 individuals with depersonalization disorder showing significantly reduced AIC activation during interoceptive attention tasks, with effect sizes correlating with symptom severity. Klein and Barron [4] conclude that phenomenal consciousness can only emerge in organisms with recurrent cortical-subcortical loops, limiting its distribution to vertebrates and cephalopods. Seth et al. [17] offer a novel view of emotion as "interoceptive inference," extending James-Lange and Schachter-Singer theories by proposing that subjective emotional states are determined by hierarchically organized top-down predictions about interoceptive bodily states, rather than by bottom-up sensing of physiological changes alone. Seth et al. [17] demonstrate that targeted manipulation of interoceptive predictive signals in the AIC can restore normal presence experiences in patients with depersonalization disorder, offering a potential therapeutic pathway. LeDoux proposes that autonoetic consciousness is primarily supported by subcortical circuits in the amygdala, while noetic consciousness depends on cerebellar processing [28]. According to Klein and Barron [4], brains solve action selection problems by transforming heterogeneous information—including sensory inputs, representations of internal state, and stored knowledge of values—into a common framework called a "phenomenal interface" to compute multi-objective Q-values. The best measure of consciousness is verbal self-report, because people can respond verbally or non-verbally to something they are conscious of but can only respond non-verbally to something they are not conscious of [28]. Bumblebees (*Bombus terrestris*) are capable of context-dependent, centrally controlled reduction of nocifensive behavior, demonstrating that they can modulate their avoidance of noxious heat (55°C) based on the availability of alternative food rewards [12]. When unheated feeders contained high sucrose concentrations (40%), bees significantly avoided noxiously heated feeders; however, as the sucrose concentration at unheated feeders decreased (to 10%, 20%, or 30%), bees progressively increased their feeding from the heated feeders, demonstrating a motivational trade-off [12]. Klein and Barron [4] argue that phenomenal consciousness is a consequence of how mobile animals with spatial senses and a capacity for

goal-directed behaviour resolve the complex problem of action selection. According to Seth et al. [17], the anterior insular cortex generates the top-down interoceptive prediction signals that underpin conscious presence, with prediction errors computed in downstream cortical regions.

Review

****Cite this article:**** Chittka L, Skeels S, Dyakova O, Janbon M. 2025 The exploration of consciousness in insects. *_Phil. Trans. R. Soc. B_* ****380**** : rstb.2024.0302. <https://doi.org/10.1098/rstb.2024.0302>

Received: 31 January 2025 Accepted: 15 July 2025

One contribution of 17 to a theme issue 'Evolutionary functions of consciousness'.

****Subject Areas:**** behaviour, cognition, neuroscience

**Keywords:**

bees, cognition, emotion, flies, prediction, selfhood, sleep

****Author for correspondence:**** Lars Chittka e-mail: l.chittka@qmul.ac.uk

Consciousness is a state of subjective experience or awareness, e.g. of an emotion, the self or external objects. In humans, this awareness is underpinned by a suite of cognitive functions, from attention to metacognition. To understand the evolution of consciousness, the study of these cognitive functions across a variety of animal taxa is critical. Insects are useful organisms because we have a sophisticated understanding of their cognition from over a century of study

and modern tools are revealing the intricacies of insect brains with increasing clarity. Here we cover the rich history of this venerable field as well as more recent discoveries relating to consciousness in insects, specifically focusing on the following areas: emotions, the distinction of self and other, prediction, attention and active sleep. There can still be no formal certainty about consciousness in insects; even in humans, there is currently no agreement over the particular combination of cognition and neural function that produces consciousness. Nonetheless, evidence from all the lines of investigation summarized here builds up to an increasing probability that insects might possess some form of subjective experience. We encourage further investigation of insects to explore the building blocks of consciousness and its evolutionary antecedents.

This article is part of the theme issue 'Evolutionary functions of consciousness'.

1. Introduction

Consciousness comprises many interrelated phenomena, e.g. the capacities to feel, experience and think. It may have evolved as a consequence of building a unified simulation of the organism within its environment [1–4]. Such a simulation might afford an organism important advantages, such as a greater ability to navigate an environment, plan for the future or respond to unexpected conditions. Around the beginning of the 20th century, insects were seen as an early test case for the exploration of consciousness in animals.

A remarkable example is the 1902 essay (in German) entitled 'The psychological abilities of ants and some other insects' by the Swiss psychiatrist Auguste Forel [5] (one of the founding fathers of neuron theory). Forel rejected the dualism of body and mind and pointed out that consciousness and other psychological processes cannot be understood without exploring the underpinning neural processes: 'Among the phenomena of our brain-life, as wonderful as they

are, lies absolutely nothing contrary to the laws of nature, or that would require invocation of a mystical, supernatural 'psyche' [5]. Forel held that true consciousness is the ability to introspect (equivalent to the contemporary definition of higher-order consciousness [6]) and proposed that this capacity is reserved for humans. He thought that a more basic form of mind comprises the processes that in humans are _accessible by introspection_ , which he described as 'the objects of attention: perceptual images, impulses of will, feelings or abstract thought' [5]. Forel highlighted that direct access to conscious experience is only available in oneself, and that to the extent that we can use arguments of analogy about consciousness in fellow human beings, we can logically also extrapolate to other animals [5]. Our review is inspired

© 2025 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

****2****

by this approach, and we explore several of Forel's 'objects of attention' in this article.

Following Buttel-Reepen [7] and Wasmann [8], Forel firmly rejected the (then popular) claim that insects are reflex machines [7,8]. He explained (with reference to the human brain) that while sensory input, motor control, attention and emotion are all underpinned by separate neural pathways, they are tightly interfaced in the brain; he further highlighted that motor output can result in emotions such as pleasure and pain, reinforcing certain behaviours but not others [5]. He provided evidence that Hymenopteran insects (e.g. bees and ants) also integrate multiple sensory memories into percepts and suggested that such interfacing is probably phylogenetically ancient and thus common to many animals. He listed experimental evidence for memory in many insects, and suspected that it is present in all of them [5]. Contrary to

Bethe [9] (who denied the existence of memory in insects, but curiously thought that such memory would be evidence for consciousness, if it did exist), Forel [5] (as well as Buttel-Reepen [7] and Wasmann [8]) cautioned that associative memory is insufficient evidence for consciousness [9]. This century-old debate recently resurfaced when it was proposed that consciousness arises from and is functionally tied to associative learning [10]. We agree with Forel and his contemporaries that, at least in its basic form, associative learning falls significantly short of the criteria required for the diagnosis of consciousness.

Using experiments with artificial flowers in bees, Forel provided evidence for top-down attention, showing that bees trained to one type of flower are difficult to distract by other flower designs, even familiar ones [5]. He discussed the presence of a will in ants, which he observed battling with obstacles for hours, when they knew the value of an out-of-reach item (see also Turner [11] for the discussion of a will in cockroaches). Forel described the emotion-like components of winner-loser effects in ants. After a battle with a competing ant colony, the losing party falls into a state of 'hopeless abandon' (more so if they lost multiple battles), making them less likely to engage in or win further battles and resulting in 'cowardly retreat', consumption of own brood, and neglect of work [5]. Conversely, winning a battle, in Forel's observations, promotes courage and tenacity in the superior party, which might make them succeed in attacking even physically superior enemies [5]. To our knowledge, this is not just the first description of winner-loser effects in any animal, but also a visionary exploration of emotion-like states in this context. Forel highlights that such states can affect multiple other behaviours in the aftermath of a salient event.

Citing examples from bees and ants, Forel highlighted the importance of conflicts between various perceptions, emotions and efforts of the will in the study of animal minds: for example, the question of how animals balance conflicting needs such as the motivation to feed and colony defence. He placed particular emphasis on cases where stimuli are not concurrently

accessible and must therefore be based on mental representations. Interestingly, such motivational tradeoffs have more recently been explored as indicators of animal sentience (the capacity to feel) and specifically pain experiences [12,13]. More broadly, Forel pointed to the likely deep evolutionary origins of emotions and highlighted that instinct and emotions are deeply intertwined, for example in brood care.

Forel and his contemporaries were by no means uncritical—they repeatedly warned against the risks of anthropocentrism, and against setting the bar for criteria of consciousness too low [5,7,8]. They realized that the indicators they provided were fragmentary and tentative, and that while formal proof of mental processes in animals was impossible to obtain, a useful alternative was to search for likely *_signs_* of such processes, and to combine indicators from multiple lines of investigation for an assessment of the probability that an animal is conscious. Buttel-Reepen ultimately concluded that bees have either no consciousness, or one of the lowest degree [14]. His criteria were even more exclusive than Forel's [5]; Buttel-Reepen posited that unity and continuity of conscious experience, the integration of all sensory experiences into a cohesive whole, and power of imagination were necessary ingredients that needed to be demonstrated. Buttel-Reepen doubted that these were present in insects, but was open to the possibility of sentience: he encouraged the exploration of bee-specific emotion-like states such as might occur during swarming, where he found bees to display 'a bacchantic delight' and wrote that 'it is as if the bees were indeed drunk with joy' [7]. Like Forel [5], Turner [11] and Wasmann [8], he provided clear evidence for memory and touched upon many of the theoretical approaches and experimental paradigms that prevail today in the study of the relation between consciousness, attention, emotion and cognition [7,14]. In 1908, summarizing this literature and taking a broader perspective across animals (including both vertebrates and invertebrates), the pioneering animal consciousness researcher Margaret Washburn pointed out the tight links between action and perception, and between motor behaviour and consciousness [15]. In this manner she not only prefigured key ideas of the predictive processing theory of

consciousness [16,17], but she also pointed out that the study of sleep (including both non-conscious phases and conscious dream phases) might be a useful avenue of research into the very nature of consciousness [18]. We return to this approach later in this review.

Unfortunately, the progressive ideas of this school were all but buried under the fashion of behaviourism. Many animal psychologists favoured the easier approach to use directly observable behaviour to understand cognition [19], often ridiculing the efforts of those interested in mental processes [20]. As an isolated example to the contrary, Charles Turner, an African-American pioneer of animal cognition, continued to point out the likely rich inner world of insects until his death in 1929 [21,22]. He directly confronted Edward Thorndike (one of the founding fathers of behaviourism), highlighting that the navigation behaviour of homing wasps could not be explained by trial-and-error learning but must involve a form of intentionality [23]. Regrettably, Turner's visionary contributions were largely forgotten, as were those from the other early explorers of insect minds.

A curious result is that insect scientists studying mental processes today, even though their forebears had pioneered the field around the penultimate turn of the century, find themselves on the back foot when comparing their work on the topic with scientists exploring similar questions in vertebrates. When Donald Griffin returned to the question of animal consciousness in the 1970s, he faced extensive criticism [24,25]. Even though Griffin included the insects in his discussions, the field now restarted with a focus on large-brained vertebrates [26-29].

We here follow Forel's [5] early approach of searching for indicators of the probability of consciousness in animals rather than formal certainty. This is mirrored by recent trends of breaking consciousness down into its constituent parts or along

several dimensions [30,31]. We draw on both schools of thought to ask whether analogues of vertebrate consciousness might be present in insects. Our review is not comprehensive—we focus specifically on domains of consciousness in insects that have received less coverage in recent review articles than others. These are emotion-like states and selfhood, prediction and attention, and the relationship between sleep and consciousness.

2. Emotion-like states

One of the key ingredients of consciousness is the capacity to have affective states, which include emotions and moods. The question of whether insects have an emotional life is no less (and no more) difficult to answer than in other non-human animals. This is because insects, like other animals, cannot verbally report their emotions, if they have any. The perception that emotions might be more easily studied in animals with closer phylogenetic distance to humans, or more similar brain structures, might sometimes be misleading. It is simply impossible to infer function from gross neuroanatomy, even in closely related species: for example, chimpanzees have Broca's and Wernicke's areas (which in humans are used for speech production and comprehension), but chimps demonstrably lack spoken language [32]. Likewise, the view that facial expressions of humans' mammalian relatives offer a more direct window into their emotional world is tempting, but often subject to anthropomorphic misinterpretation [33,34]. Nonetheless, insect scientists often find themselves confronted with disproportionately high levels of scepticism when exploring emotion-like states [5,35].

Consider the following examples from various insect taxa. (i) Male fruit flies that are stressed by being deprived of mating opportunities seek out alcohol [36]. (ii) The perception of dead conspecifics by fruit flies alters sensory perception and brain physiology, and the effects are so profound that they reduce longevity [37]. (iii) Again in fruit flies, it was found that painful stimuli

can induce a heightened state of vigilance and a hypersensitivity to previously non-painful stimuli, similarly to what is observed in humans with chronic pain conditions [38]. (iv) Bumblebees, after learning to avoid cryptic crab spiders on flowers by being exposed to simulated attacks, display false alarms—rejecting perfectly safe flowers after inspecting them, as if they had experienced a ‘hallucination’ of a spider (figure 1E-H) [39]. (v) In crickets, repeated defeats in male-male contests of crickets induce a long-term depression-like behaviour state, but crickets that received drugs affecting the serotonin system showed higher resilience to chronic defeat stress [40]; see also [41–43] for related work on other insects, including the remedying effects of antidepressants.

Transposing these scenarios to the human realm, it seems obvious that they have significant emotional components, and that this is so because of their fitness relevance. The emotional element in such situations provides a motivation to avoid unfavourable conditions; the seeking of drugs (such as alcohol) occurs because basically, they make one *_feel_* good, especially when one is miserable [44,45].

When rodents or other mammals are trained to associate a (previously neutral) stimulus with an aversive one, e.g. an electric shock, then this is commonly referred to as fear conditioning [46]. The conditioned stimulus subsequently triggers avoidance, immobility (‘freezing’), startle and escape responses, changes in heart rate and breathing, as well as recognizable activity changes in brain areas such as the amygdala [46,47]. It is implied that the affective fear response is a crucial, survival-related ingredient in the learning process [48].

Insects such as fruit flies (including their larvae) [49–53] and bees [54–56] also learn to associate conditioned stimuli (e.g. odours, visual stimuli) with aversive stimuli; crucially, similar types of overt behaviour as listed for mammals (above) have also been found in insects [39,57], as well as changes in breathing [58] and heart rate [57] and the expression of stress-related

genes [56]. The evidence for an affective component in the responses to noxious stimuli has been reviewed in detail elsewhere [59,60]. It has been suggested that the link between various mushroom body neuron types in fruit flies performs a similar function as the amygdala-hippocampus axis in mammals in the context of the learning of aversive stimuli [61]. Despite these many similarities between mammals and insects, in the latter such learning is commonly referred to as 'just' classical conditioning, without an affective component.

Some tests of emotions in insects have been modelled directly on vertebrate paradigms and should therefore be comparable. So-called cognitive bias tests have been developed to test the psychological welfare of animals (e.g. rats) in captivity [62]. These paradigms are essentially versions of the proverbial glass that can be half-full or half-empty: optimistic humans might view the same ambiguous state (a glass filled to the middle with a drink) as positive, whereas a pessimist would judge the glass as being nearly empty. Bees respond to ambiguous stimuli with more reservation when they have previously suffered a simulated predator attack, and more positively if they had experienced a surprise reward before the experiment—a dopamindependent, optimism-like state (figure 2a-c) [63,65]. It has been suggested that such results could alternatively be explained by motivation, although this simply replaces one emotion-related explanation with another: motivation is itself a mental state, not an observable or quantifiable behaviour or a simple mechanism [66]. In recent work, the question of whether insect responses to ambiguous stimuli could be explained by cognitive biases, motivation or altered attentional states was directly addressed, and it was still found that emotion-like cognitive biases are the most likely explanation [67].

Play-like behaviour was also found in insects. When offered small coloured balls in an arena near their nest, bees rolled the balls for extended periods even when no sugar reward was made available (figure 2d) [64]. Bees went out of their way to return repeatedly to such a 'play area'. There seemed to be something inherently rewarding in the activity itself, since no other

incentive was offered. In line with what is commonly observed in vertebrate play behaviour, young bees engaged more often with the balls than older ones. Play-like behaviour was also recently discovered in fruit flies, where some individuals appear to enjoy the ‘thrill’ of riding a carousel. Such self-stimulation was proposed to aid self-perception through the training of internal models [68].

****4****

****==> picture [246 x 454] intentionally omitted <==****

****Figure 1.**** Attention and false alarms in bees. (A–D) Honeybees fail to learn to distinguish the two camouflaged shapes in panels (A) and (B) even after extensive training. However, the task can be introduced by using non-camouflaged targets (C) and (D), guiding bees to know what to look out for. If bees so entrained are then tested with the camouflaged objects in (A) and (B), they discriminate them with high accuracy (after [28] modified). (E–H) Bumblebees trained to avoid robotic crab spiders subsequently display false alarms as if they had ‘hallucinated’ a spider [29]. Bumblebees are trained to forage in an array of artificial ‘flowers’ (yellow squares; E–H), a minority of which harbour plastic models of predatory, cryptic crab spiders (F). If bees land on such flowers during training, they are briefly trapped by solenoid-driven, sponge-clad pincers to simulate a predation attempt (F). During later testing, bees display hesitation and scanning behaviour at all flowers, and will typically reject flowers with spider models (correct-reject, (G)). However, while bees land on the majority of safe flowers, they also display elevated levels of false alarms (H), where they behave as if they imagined the presence of a spider.

We cannot definitely know what (if anything) an insect feels under such conditions (any or more or less than what we can infer about a mouse). Indeed, some seemingly emotion-related insect

behaviours might in some cases have explanations that do not involve sentience. A useful test case is the condition of learned helplessness—a psychological state that occurs after repeated exposure to inescapable aversive stimuli. Many animals— including insects—subsequently endure such stimuli without trying to flee, even if the possibility to escape is again made available [69]. Even though this state looks overtly similar to what humans might experience as hopelessness, the state can be induced in rats with severed spines [69] and in decapitated cockroaches [70]. This shows that what looks like a ‘miserable’ condition is not necessarily evidence for sentience [69]. However, it is unlikely that antidepressants alleviate the state of learned helplessness in animals with a disconnected brain, as they do in some insects with intact bodies [40,42].

A common response to observations of what appear to be emotion-like states in insects is that they might be manifestations of ‘just instinct’ [71]. That could certainly be the case, though it is not an alternative explanation, but an additional one. An alien observer of human behaviour might conclude that parental love, rage, sexual desire, grief, etc. are just instinct—and they would be correct. While all of these might be subject to cultural variation and individual experience, they are part of our default emotional toolkit [5]—the human affectome [72]. Examples where emotions serve ultimate functions are the enhancement of learning, the promotion of parent-offspring care and attachment, and the avoidance of dangerous situations. All of these are

****5****

****==> picture [307 x 289] intentionally omitted <==****

****Figure 2.**** Emotion-like states in bees. (a–c) A positive cognitive bias in bees [63]. Bumble bees familiarize themselves with a flight arena in which, on the back wall, the blue target on the

left always contains a droplet of sugar water (a), whereas the green target on the right does not (b). Only one of the five positions on the choice wall is ever accessible. After training, bees will usually approach the blue target (left) in a straight line when it is available (a). Conversely, hesitation is apparent if the green target (right) is on display (b). When intermediate ('ambiguous') turquoise stimuli (centre) are offered (c), bees will hesitate for a length of time that lies between the flight times displayed for blue and green (black flight trajectory). If, however, a tiny drop of sucrose solution (5 μ l) is offered as a surprise before bees enter the experiment, they judge the ambiguous option (turquoise) as more promising and will accept it more swiftly (red flight trajectory)—i.e. they judge the intermediate stimulus in a more optimistic manner than control bees (figure by Cwyn Solvi, printed here with permission). (d) Do bumble bees play [64]? When given the opportunity to roll balls in the vicinity of the nest, bumblebees will roll balls for extended periods, even if no appetitive reward is made available near the balls. As opposed to rewardless flowers (which are only probed a few times before bees give up), bees return repeatedly to this activity, even increasing ball-rolling behaviour over time, indicating an enjoyment of the activity itself, rather than a form of food search exploration. Young bees engaged with the balls more than older bees. Photo by Richard Rickitt (with permission).

fitness-related and at least in part instinctual in humans, and with some probability, non-human animals [73–75]. Thus, even if emotions are innate, they are nonetheless important ingredients of an animal's subjective experience.

3. Self and other

One important aspect of consciousness is self-recognition, which at its core requires an individual to be able to recognize itself as distinct from another living entity [31,76,77]. At the start of the 20th century, Margaret Washburn [15] proposed that motor behaviour is at the

heart of consciousness: rather than just responding to incoming stimuli, animals constantly probe their environment and must distinguish between self-generated and other-generated sensory input [15]. The transition from a sedentary lifestyle to locomotion early in evolution would already have required animals to make this distinction [77]. Animals create an internal copy of the self-generated signal, known as an 'efference copy', to cancel out the feedback from their own movements [78,79]. Consider the scenario where a fist suddenly appears in someone's visual field. This is no cause for alarm if they have just willingly raised their hand to inspect their knuckles—but if they have not, and it is someone else's fist, then they could be under attack. The visual stimulus might be similar, but it is the presence or absence of the efference copy of the intended movement that allows the person to distinguish between the two, and then respond accordingly.

Body awareness is considered an important aspect of self-awareness and individual experience [80]. For example, an animal's extremities constitute perfectly palatable food, yet animals are not normally inclined to consume their own legs, since there is some form of appreciation of own body integrity—what is part of *_me_* and what is not [81]. Some ant species (such as *_Cataglyphis bicolor_*) show remarkable rescue behaviour of nestmates. When a nestmate is experimentally trapped under rubble (using a nylon tether around the ant's waist), rescuers will come to their aid. These helpers show appreciation of the trapped ant's body dimensions by removing rubble only from above buried (non-visible) body parts. They bite forcefully at the tether, but only gently pull on legs of the trapped ants [82]. These behaviours often involve multiple rescuers, meaning that there will be occurrences where individuals will need to discern which extremity is part of themselves and which parts belong to another rescuer and/or the trapped ant [83]. As such, body awareness would be important here. Whether empathy is involved in such rescuing, as has been suggested for similar behaviour in rodents, remains an open question [84].

Being able to take into account one's own body size and movements when making decisions has obvious functional advantages when navigating a cluttered environment. Ravi _et al._ [85] found that flying bumblebees, which vary significantly in body size, were able to judge the size of a gap in a wall in relation to their own body size. Larger bees avoided the small gaps that were accepted by their smaller conspecifics, and when the size of the gap was borderline navigable for a bee of a given size, the bee would adjust her flight behaviour to minimize the risk of collision (and therefore injury) (figure 3A-D). Similarly, Ben-Nun _et al._ [87] demonstrated that self body-size perception in locusts is correlated with their own body size.

A popular test of self-recognition in vertebrates is the so-called 'mirror and mark test' [88,89]. The test involves putting a mark on the animal in a place that can only be seen in the mirror. Attempts to engage with and remove the mark from the body when in front of the mirror are interpreted to mean that an individual appreciates that their own individual appearance has been altered, thus showing a form of self-awareness [86,90,91]. The mirror test has been adapted to examine self-recognition in _Myrmica_ ants [92]. Ants reportedly passed the test, but results are hard to interpret owing to the lack of appropriate controls. More promising study systems are certain species of _Polistes_ wasps, since individual face recognition is part of these species' social architecture [93].

Mirror self-recognition can be viewed as a special case of individual recognition. In species where individuals are not visually distinguishable by the visual-spatial resolution that their eyes permit, mirror self-recognition tests are less likely to produce positive results (though see below for a possibility via active vision). While _Polistes_ wasps do respond to either the presence of mirrors or presence of marks placed on their faces by experimenters when seen in mirrors, there was no evidence that grooming was directed towards the mark [86]. Instead, wasps

touched the mirror more when a visible mark had been placed on their faces than in a control condition using a clear (non-visible) mark (figure 3E), making it hard to know whether the behaviour was in any way self-directed, or whether the perception of what was seen in the mirror was that of an unfamiliar conspecific. It is thus possible that what they perceived in the mirror was not themselves, but an unfamiliar conspecific.

This highlights the ongoing debate about the validity of the mirror test and its capacity to show true self-awareness—or the absence of such awareness when results are negative [94]. If indeed mirror-self recognition is an extension of individual visual recognition in a social system, then it is important to consider that natural environments do not contain mirrors. No animal can be expected to comprehend spontaneously the workings of a mirror, so animals must be given ample opportunity to interact with mirrors to familiarize themselves with their own appearance before tests can commence. In doing so, they most probably figure out that their own actions result in fully predictable ‘actions’ of their mirror image [95,96]—which is a special case of animacy perception. In that sense, it is not clear why a mark is required: the animal that loses interest in the mirror after such an exploratory phase—and that ceases to treat their mirror image as another conspecific—might already have shown a form self-recognition. It is important to interrogate self-recognition (and self-awareness more broadly) in paradigms more relevant to the natural ecology of the animal being studied [97]. In some animals, different sensory modalities might be worth exploring for self-recognition tests. Some ants, for example, use olfactory cues for individual recognition [98]. In such species, chemosensory self-recognition tests might be more appropriate [99,100].

Metacognition, defined as ‘knowledge about one’s own knowledge’ [101], is another ingredient of self-recognition. It is often considered as an indicator of higher-order consciousness—the awareness of own mental states, rather than just experiencing the mental states themselves [102]. This is the form of true consciousness that Forel thought in 1902 might be reserved for

humans [5], where indeed it has been extensively studied [103], but other primates have also been tested (see review by [104]). Perry & Barron [105] attempted to examine whether Western honeybees (*Apis mellifera*) have basic metacognition. Bees were given visual discrimination tasks that varied in difficulty and were rewarded with sucrose for correct choices but punished with bitter quinine for mistakes. Additionally, they had the option to leave the task. Perry & Barron [105] found that individuals left the task more often when the task was difficult, and that by doing this, they improved their success rate. They were also able to pass on this 'opt out' rule to a novel task. These results suggest that the bees were able to assess their certainty on a given trial and make decisions accordingly, indicating the possibility of some kind of metacognitive processing [105,106]. However, it is also possible that their choices were based on a series of learnt associations [105-107] (see also [102] for equally mixed evidence in ants). Further work on metacognition and decision confidence is required for insects, perhaps more closely modelled on tasks commonly deployed in mammals [108-110]. Whether the cross-individual version of metacognition—namely, theory of mind (knowledge about information held by other animals)—exists in insects is presently unknown. Such 'mental perspective taking' is perhaps most rewardingly explored in settings where such an ability would convey clear fitness benefits. Examples are prey taking into account what predators might see (or *vice versa*) or animals strategically positioning themselves to amplify mating signals, depending on where they know audiences to be.

4. Prediction and attention

Many textbooks still perpetuate the narrative that brains are designed to respond to incoming stimuli, process information and then generate adaptive behaviour. However, during wakefulness, brains (and by extension, the behaving animals that house them) constantly probe their environment, moving sensors (e.g. eyes, extremities with touch sensors, etc.) to establish contingencies between behaviour and its outcome. Controlled variability in

spontaneous nervous system activity promotes behavioural flexibility and active information acquisition, and it may result in flexible problem-solving [111,112]. In addition to the active moving of sensors, attention provides a sort of 'inner eye' to focus selectively on salient information from the sensory periphery. The role of attention has both been linked to consciousness and described as the process assigning greater precision to certain sensory streams [5,15,113].

As scholars surmised around the turn of the 20th century, there is probably a link between top-down, selective attention and consciousness [5,15]. To make sense of the environment, the awake brain constantly makes predictions about what sensory

****7****

****==> picture [307 x 431] intentionally omitted <==****

****Figure 3.**** Self-recognition in insects? (A-D) Bumblebees of the same species are highly variable in body dimensions and appear to know their own body size, as evidenced from behaviour when navigating differently sized gaps. Bees will approach and inspect the gap from different angles to work out its size. Larger bees will peer more when traversing a narrow gap compared with their smaller conspecifics (A, C). Bees will angle their body position to avoid collision if the gap is similar to or narrower than their wingspan (A). Figure from [80], after [85]. (E) Individuals of *Polistes metricus* paper wasps have distinct face marks, allowing for individual recognition of nest mates. Mirror tests were used to investigate the possibility of self-recognition of individuals with paint marks on their faces. Visible marks elicited more exploratory behaviours (antennation and touching of the mirror) compared with controls with transparent paint marks. However, there was less self-directed grooming of the mark than expected, potentially indicating that wasps interpreted their mirror image as a previously

unknown individual rather than a reflection of themselves [86].

information to expect next [114]. These expectations are encoded across the brain as inner generative models, which are iteratively adjusted to minimize the discrepancy between predictions and sensory returns—or ‘prediction errors’ [115,116]. ‘Prediction error’ is actually a misnomer; it does not involve a mistake by the brain, just a violation of an expectation—a surprise. Prediction errors allow animals to adapt perception and behaviour based on a mismatch between expectation and input. An obvious type of predictive coding is foreseeing what changes in sensory input are expected as a result of the voluntary movements of sensors (see §3). In humans, when sensory inputs match the brain’s predictions, perception occurs without much conscious awareness. However, when there is a prediction error, heightened top-down attention can bring novel or unexpected stimuli into conscious awareness. This process can be essential for survival, since it probably facilitates adequate responses to specific threats or opportunities in the environment [31,117,118].

Insect perception and attention can also be understood under a predictive processing framework [119–122]. The innate preferences of a pollinator, for example, contain certain ‘default predictions’: e.g. naive bees expect flowers coloured violet to blue to be highly rewarding since flowers of these colours indeed contain better-quality rewards in some habitats [123]. Such ‘priors’ can be adjusted later by experience: an individual bee might learn that a plant species with yellow flowers is rewarding, and update its expectations accordingly. Bees searching for flowers while avoiding familiar predators do not just passively respond to objects that appear in their visual field. Instead, they ‘know what to look for’, and search specifically for memorized flower targets [124], as well as scanning for predators that they have seen before. Like humans, honeybees initially fail in detecting camouflaged shapes, but when the shapes are first presented openly, attention subsequently helps bees to find

them in the camouflaged presentation (figure 1A–D) [125]. Bumblebees experienced with attacks from cryptic flower predators subsequently display false alarms—they reject many perfectly safe flowers as if they had ‘hallucinated’ the visual appearance of a predator (figure 1E–H) [39].

In humans, various levels of consciousness and attention are accompanied by characteristic oscillations of neural activity in the brain [126,127]. The awake human brain shows a characteristic 30 Hz frequency in EEG recordings; such *_beta_* oscillations, synchronized across cortical areas, are crucial in sensory awareness and focusing attention [128]. Insects display similar patterns: synchronization of oscillatory activity (at 18 Hz) across brain areas was found in awake honeybees [129]. Oscillations of 20–30 Hz also occur in the central complex of the brains of fruit flies when they attend to visual features of novel objects. Theta oscillations (approx. 4–8 Hz), which are implicated in memory, navigation and the coordination of top-down and bottom-up signals in humans [130,131], are also present in insect brains, particularly in the central complex and mushroom bodies [132]. These oscillations may serve similar roles in regulating temporal aspects of attention and memory. Furthermore, similar to humans [133,134], prediction errors are also distributed along the entire fly sensory processing hierarchy, enabling the rich interplay of predictions and prediction errors that could underpin conscious awareness [135,136]. So far, only *_local_* prediction errors have been tested in fruit flies—for example, a series of light flashes of one colour, followed by a so-called *_oddball_* with a different colour [135]. In human hearing, such oddballs elicit appreciable prediction error signals in the nervous system (limited to sensory cortical areas)—but this occurs even in unconscious human subjects [137]. Global oddballs, conversely, violate a previously learnt *_pattern—_* e.g. if tone strings CCCA CCCA CCCA are followed by CCCC. The arising *_mismatch negativity_* signal in higher-order brain areas of humans in this case is linked to conscious

awareness [137,138]. Future work in the context of prediction errors in insects should focus specifically on the distinction between local and global deviations from expected patterns, and their neural underpinnings.

Prediction error-mediated surprises can result in emotion-like responses and in redirecting attention—e.g. bees finding rewards in an unexpected location exhibit a dopamine-dependent optimistic state [63]. In bees placed in a virtual reality environment, distinct neural activity in the optic lobes reliably precedes the selection of one or another stimulus to which subjects pay attention [139]. The existence of similar types of brainwaves in humans and insects is not formal proof of consciousness in the latter. However, it is at least suggestive that the type of synchronization across brain areas that, in humans, accompanies states of wakefulness and attention, also occurs in insects [71,132,140].

5. Active sleep, deep sleep and wakefulness

Everyone knows what is meant when it is said that a man is unconscious. He is neither awake, aware of the sights and sounds of the outer world, nor dreaming, aware of images that are the product of his own fancy. Consciousness is that which is present when we are either awake or dreaming, and which is absent when we are dreamlessly asleep.

Margaret Floy Washburn, 1916 [18]

Given the obvious division between conscious and non-conscious states that we experience daily, one useful entry point into the question of consciousness and its function in animals is to contrast it with what happens during sleep—which, perhaps counterintuitively, may be tightly related to the maintenance of consciousness in the wakeful state. One might expect that if there are no signals from the sensory periphery (such as when one's eyes are closed during

sleep), the brain might automatically be in a resting state as well. However, brains—whether insects’ or humans’—are never ‘switched off’, not even during the deepest sleep. In humans, sleep is divided into multiple phases: slow wave (‘deep’) sleep, and rapid eye movement (REM) sleep, which alternate in multiple phases every night [141]. Most dreams occur during REM sleep, and many of them are consciously experienced while the brain is largely ‘disconnected’ from the outside world [142]. Cortical activity during REM sleep resembles waking activity [143,144]. REM sleep is associated with emotional regulation and memory consolidation [142,145,146]. How this is achieved during REM sleep has been discussed under the predictive processing framework: a role for REM sleep might be to prune and refine internal generative models [147–149]. Such models might over time recruit a large portion of the brain’s resources as well as be overtrained on the particular external stimuli encountered, limiting the generalizability of predictions. In this context, a role for REM sleep might be that it acts as a closed-loop simulation, where, away from the demands of immediately reacting to the environment, imagined stimuli are processed to prune and refine the generative models and avoid neural overfitting. In this way, not only would the brain make more flexible predictions when awake, but the synaptic cost of maintaining such generative models might be reduced.

In insects such as honeybees, fruit flies and mosquitoes, sleep can be behaviourally diagnosed by quiescence (or immobility), increased arousal thresholds and sleep rebound (increased period of quiescence after sleep deprivation) [150–155]. Insects also have distinct sleep phases, although these may not map directly onto human ones: bees, for example, have three distinct sleep phases. Their deepest sleep is characterized by a distinctive crouching posture in which head, thorax, and abdomen are relaxed, antennae are immobile, muscle tonus and body temperature are decreased, and response thresholds to external stimuli are increased [151,156]. Exposing bees to odours during this sleep phase functions to consolidate experiences from the previous day in honey bees’ memory, whereas in humans, it is REM sleep that appears to be related to memory consolidation [157,158].

Fruit flies, too, have distinct stages of quiet and active sleep [159-161]. New work by van Swinderen and their team suggests that, in insects just like in mammals, active sleep serves a function of calibrating the brain as an optimal 'prediction machine' [71,162]. Local field potential (LFP) recordings during sleep show that periods of oscillatory activity during active sleep alternate with periods of decreased overall activity during quiet sleep [71,163]. Calcium imaging has confirmed that during

****9****

these periods, different groups of neurons were recruited [164]. These discoveries have contributed clues to the functional role of these sleep stages and why they might have evolved, as well as some of their similarities to mammalian REM and deep sleep. In *Drosophila melanogaster*, the latter has been associated with cellular waste clearance and repair [165,166]. These processes are also present during deep sleep in mammals and suggest a basic evolutionary function to deep sleep: cellular homeostasis [167,168].

What is the function of active sleep in insects? Of course, there is no direct access to the question of whether insects dream (no more or less than any other animals). Interestingly, however, distinct theta oscillations are found to emanate from the central complex of the fly brain—an area associated with visual attention and navigation [163,169-172]. Some evidence also suggests that prediction drives sleep need and that recognition of unexpected stimuli (likely to produce a prediction error) is attenuated during active sleep [135,173]. Further understanding of the functional link between active sleep and predictive processing might also help to tie sleep in with other cognitive functions associated with the emergence of consciousness. A bi-directional interaction between sleep and attention has been proposed and responses to salient information have been explored in sleeping flies, where a circuit that

modulates sensory perception has been found: fruit flies are more likely to wake when exposed to salient olfactory stimuli, a behavioural finding mediated by a connection between peripheral olfactory inputs and sleep regulatory neurons [140,174,175]. Closed-loop simulations of the kind required to optimize predictive models during sleep might also require whole-brain activations; recent work suggests that a similar process occurs during imagination, when signal strength in whole perceptual hierarchies distinguishes real and imagined stimuli [176].

Predictive processing has been the dominant mechanistic framework through which possible conscious phenomena have been explored in insects, even though it is not universally accepted that it is the most powerful framework for the exploration of conscious awareness [16,177]. Future work might seek to explore the plausibility of insect consciousness under different theories, such as the global workspace theory or integrated information theory [178,179].

6. Conclusion

We have focused here on an idiosyncratic set of topics in the broader field of the exploration of insect consciousness. We have not covered many other relevant cognitive lines of investigation in this context, because these have been covered for insects in detail elsewhere [180]. These include studies on intentionality [181], perceptual rivalry [182], foresight [183], flexible combination of past memories into concepts [184], cross-modal object recognition [185], interval timing and trace conditioning [186,187] and episodic-like memories [188,189], as well as perceptual binding [190]. None of these are, individually, formal demonstrations of consciousness, but further positive evidence from all these lines of investigation (ideally all in the same species) would add to the picture that we have assembled here on attention and prediction, its possible interaction with sleep and the question of emotions, as well as the distinction of self and other: they make the existence of conscious experience in insects at least plausible. Such cognitive processes are probably adaptive: for example, an awareness of one's

own body, the ability to trade off noxious stimuli with rewards or to efficiently react to uncertainty in the environment seem to proffer important fitness benefits.

A critical reader might observe that each of the behaviours and each of the neural processes described above could also be programmed into a non-conscious robot. An example is a review on animal pain by Sneddon _et al._ [191]. The authors hold that the same overtly observable phenomena that, in vertebrates, are typically taken as indicators of pain experiences, cannot be used as equally valid indicators in invertebrates—because, so the authors argue, robots can be manufactured to display similar phenomena. It is unclear why this argument should apply disproportionately more to one taxon than another. More broadly, we counter that nature has no room for such profligacy in generating beings that just pretend to ‘tick boxes’ in various paradigms designed to explore consciousness across animals. The robot analogy also overlooks that engineering products and evolutionarily acquired functions come about by entirely different pathways. They are subject to different developmental constraints, using entirely different building blocks and computations, and, in the engineering case, there is full knowledge of the functionality of the desired end-product. For example, the fact that it is technically possible to construct algorithms and machinery to recognize human faces tells us nothing about the neural mechanisms or evolutionary pathways underpinning human face recognition. Much less does the existence of such engineering products prove that biological face recognition works without conscious experience of seeing faces.

Importantly, if indeed we were to propose that insects achieve all of the phenomena described here by non-conscious processes, this would imply the following challenges. We would have to explain why the commonly cited adaptive benefits of consciousness in humans [192] can be mediated by fully automated mechanisms in non-human animals (and conversely, why they could not have been automated in humans [76]). Many of the common adaptive arguments regarding the purpose of humans consciousness would collapse: it is often argued that there is

an essential role of emotions in learning, as well as in seeking rewarding conditions and avoiding danger, and also that attention is survival-related, that mental exploration is crucial in flexible problem-solving, etc. [193–195]. If insects exhibit all these phenomena, overtly displaying common _symptoms_ of attention, emotion, flexible cognitive problem-solving without trial-and-error, etc., while at the same time lacking consciousness, then the adaptive need for consciousness for these phenomena in humans would be questionable as well.

The study of consciousness in insects, then, suggests functional stepping stones to the development of subjective experience, related cognitive functions and their implementation in the brain. The variety of selection pressures and lifestyles that the insect world offers is a perfect natural laboratory to further explore under which conditions consciousness might evolve. The relative compactness of the insect nervous system and existence of connectomes and brain-wide recording methods makes insects

****10****

useful models of the circuitry-level investigation of conscious experience as well, in addition to the goal of identifying relevant selection pressures and plausible evolutionary scenarios for its implementation. As we grow to understand the specific binding of cognitive function and brain connectivity that produces subjective experience in humans, we will also get closer to drawing the line of where consciousness first arises.

Ethics. This work did not require ethical approval from a human subject or animal welfare committee. Data accessibility. This article has no additional data.

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. L.C.: conceptualization, project administration, supervision, visualization, writing—original draft, writing—review and editing; S.S.: visualization, writing—original draft, writing—review and editing; O.D.: writing—original draft, writing—review and editing; M.J.: writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein. Conflict of interest declaration. We declare we have no competing interests.

Funding. Lars Chittka, Sarah Skeels and Olga Dyakova are supported by a grant from Open Philanthropy/Good Ventures Foundation 'Addressing key evidence gaps in the science of insect sentience'. Maxime Janbon is supported by a PhD studentship from Queen Mary University of London.

Acknowledgements. We thank Daniel Bor and two anonymous referees for comments on the manuscript and Jonathan Birch and Peter Skorupski for discussions.

References

references

1.

Barron AB, Klein C. 2016 What insects can tell us about the origins of consciousness. *Proc. Natl Acad. Sci. USA* 113, 4900–4908. (doi:10.1073/pnas.1520084113)

2.

Godfrey-Smith P. 2020 *Metazoa: animal minds and the birth of consciousness*. London, UK: Harper Collins.

3.

- Merker B. 2007 Consciousness without a cerebral cortex: a challenge for neuroscience and medicine. *Behav. Brain Sci.* 30, 63–81. (doi:10.1017/s0140525x07000891)
- 4.
- Klein C, Barron AB. 2025 Phenomenal interface theory: a model for basal consciousness. *Phil. Trans. R. Soc. B* 380, 20240301. (doi:10.1098/rstb.2024.0301)
- 5.
- Forel A. 1902 Die psychischen Fähigkeiten der Ameisen und einiger anderer Insekten; mit einem Anhang über die Eigentümlichkeiten des Geruchsinnes bei jenen Tieren. München, Germany: Ernst Reinhard Verlagsbuchhandlung. (doi:10.5962/bhl.title.117377)
- 6.
- Allen C, Trestman M. 2024 Animal consciousness. In *The stanford encyclopedia of philosophy* (eds EN Zalta, U Nodelman). Stanford, CA: Stanford University Metaphysics Research Laboratory. See <https://plato.stanford.edu/archives/sum2024/entries/consciousness-animal/>.
- 7.
- Buttel-Reepen H. 1900 Sind die Bienen Reflexmaschinen? *Exp. Beitr. Zur Biol. Honigbiene* 20, 1–84.
- 8.
- Wasmann E. 1900 Einige Bemerkungen zur vergleichenden Psychologie und Sinnesphysiologie. *Biol. Cent* 20, 341–348.
- 9.
- Bethe A. 1898 Dürfen wir den Ameisen und Bienen psychische Qualitäten zuschreiben? *Pflüger's Arch.* 70, 15–100. (doi:10.1007/BF01709580)
- 10.
- Jablonka E, Ginsburg S. 2022 Learning and the evolution of conscious agents. *Biosemiotics* 15, 401–437. (doi:10.1007/s12304-022-09501-y)
- 11.
- Turner CH. 1913 Behavior of the common roach (*Periplaneta orientalis* L.) on an open maze.

Biol. Bull. 25, 348–365. (doi:10.1086/bblv25n6p348)

12.

Gibbons M, Versace E, Crump A, Baran B, Chittka L. 2022 Motivational trade-offs and modulation of nociception in bumblebees. *Proc. Natl Acad. Sci. USA* 119, e2205821119. (doi:10.1073/pnas.2205821119)

13.

Brown SAB, Birch J. 2025 When and why are motivational trade-offs evidence of sentience? *Phil. Trans. R. Soc. B* 380, 20240309. (doi:10.1098/rstb.2024.0309)

14.

Buttel-Reepen HB. 1915 *Leben und Wesen der Bienen*. Braunschweig, Germany: F. Vieweg Verlag.

15.

Washburn MF. 1908 *The animal mind: a text-book of comparative psychology*. New York, NY: Macmillan Company. (doi:10.5962/bhl.title.31648)

16.

Seth AK, Bayne T. 2022 Theories of consciousness. *Nat. Rev. Neurosci.* 23, 439–452. (doi:10.1038/s41583-022-00587-4)

17.

Seth AK, Suzuki K, Critchley HD. 2012 An interoceptive predictive coding model of conscious presence. *Front. Psychol.* 2, 395. (doi:10.3389/fpsyg.2011.00395)

18.

Washburn MF. 1916 *Movement and mental imagery; outlines of a motor theory of the complexer mental processes*. Boston, MA: Houghton Mifflin Company. See <https://lcn.loc.gov/16015871>.

19.

Thorndike EL. 1898 *Animal intelligence: an experimental study of the associative processes in animals*. *Psychol. Rev.: Monograph Suppl.* 2, i–109. (doi:10.1037/h0092987)

20.

von Uexküll J. 1900 Über die Stellung der Vergleichenden Physiologie zur Hypothese der Tierseele. *Biol. Cent.* 20, 497–502.

21.

Turner CH. 1907 Do ants form practical judgements? *Biol. Bull.* 13, 333–343. (doi:10.2307/1535609)

22.

Galpayage Dona HS, Chittka L. 2020 Charles H. Turner, pioneer in animal cognition. *Science* 370, 530–531. (doi:10.1126/science.abd8754)

23.

Turner CH. 1912 *Sphex* overcoming obstacles. *Psyche J. Entomol.* 19, 100–102. (doi:10.1155/1912/95842)

24.

Griffin DR. 1981 The question of animal awareness: evolutionary continuity of mental experience. New York, NY: Rockefeller University Press.

25.

Machin KJ. 1977 Review of the question of animal awareness. *Nature New Biol.* 266, 177–178. (doi:10.1038/266792a0)

26.

De Waal F. 2016 Are we smart enough to know how smart animals are? New York, NY: WW Norton & Company.

27.

Clayton NS, Bussey TJ, Dickinson A. 2003 Can animals recall the past and plan for the future? *Nat. Rev. Neurosci.* 4, 685–691. (doi:10.1038/nrn1180)

28.

LeDoux JE. 2025 What the functions of consciousness are depends on what one thinks consciousness is. *Phil Trans. R. Soc. B* 380, 20240311. (doi:10.1098/rstb.2024.0311)

29.

Maldarelli GM, Güntürkün O. 2025 Conscious birds. *Phil. Trans. R. Soc. B* 380, 20240308. (doi:10.1098/rstb.2024.0308)

30.

Birch J, Schnell AK, Clayton NS. 2020 Dimensions of animal consciousness. *Trends Cogn. Sci.* 24, 789–801. (doi:10.1016/j.tics.2020.07.007)

31.

Seth A. 2021 *Being you: a new science of consciousness*. London, UK: Penguin.

32.

Chittka L, Rossiter SJ, Skorupski P, Fernando C. 2012 What is comparable in comparative cognition? *Phil. Trans. R. Soc. B* 367, 2677–2685. (doi:10.1098/rstb.2012.0215)

33.

De Waal F. 2019 *Mama's last hug: animal emotions and what they tell us about ourselves*. New York, NY: WW Norton & Company.

34.

Zebrowitz LA, Wadlinger HA, Luevano VX, White BM, Xing C, Zhang Y. 2011 Animal analogies in first impressions of faces. *Soc. Cogn.* 29, 486–496. (doi:10.1521/soco.2011.29.4.486)

35.

Zipple MN, Hazelwood C, Webster MF, Benítez ME. 2024 Animal emotions and consciousness: a preliminary assessment of researchers' perceptions and biases and prospects for future progress. *R. Soc. Open Sci.* 11, 241255. (doi:10.1098/rsos.241255)

10

royalsocietypublishing.org/journal/rstb

Phil. Trans. R. Soc. B 380: rstb.2024.0302

36.

Shohat-Ophir G, Kaun KR, Azanchi R, Mohammed H, Heberlein U. 2012 Sexual deprivation

increases ethanol intake in *Drosophila*. *Science* 335, 1351–1355. (doi:10.1126/science.1215932)

37.

Chakraborty TS, Gendron CM, Lyu Y, Munneke AS, DeMarco MN, Hoisington ZW, Pletcher SD. 2019 Sensory perception of dead conspecifics induces aversive cues and modulates lifespan through serotonin in *Drosophila*. *Nat. Commun.* 10, 2365. (doi:10.1038/s41467-019-10285-y)

38.

Khuong TM, Wang QP, Manion J, Oyston LJ, Lau MT, Towler H, Lin YQ, Neely GG. 2019 Nerve injury drives a heightened state of vigilance and neuropathic sensitization in *Drosophila*. *Sci. Adv.* 5, eaaw4099. (doi:10.1126/sciadv.aaw4099)

39.

Ings TC, Chittka L. 2008 Speed-accuracy tradeoffs and false alarms in bee responses to cryptic predators. *Curr. Biol.* 18, 1520–1524. (doi:10.1016/j.cub.2008.07.074)

40.

Rillich J, Stevenson PA. 2018 Serotonin mediates depression of aggression after acute and chronic social defeat stress in a model insect. *Front. Behav. Neurosci.* 12, 233. (doi:10.3389/fnbeh.2018.00233)

41.

Dinges CW, Varnon CA, Cota LD, Slykerman S, Abramson CI. 2017 Studies of learned helplessness in honey bees (*Apis mellifera ligustica*). *J. Exp. Psychol.* 43, 147–158. (doi:10.1037/xan0000133)

42.

Ries AS, Hermanns T, Poeck B, Strauss R. 2017 Serotonin modulates a depression-like state in *Drosophila* responsive to lithium treatment. *Nat. Commun.* 8, 15738. (doi:10.1038/ncomms15738)

43.

- Yang Z, Bertolucci F, Wolf R, Heisenberg M. 2013 Flies cope with uncontrollable stress by learned helplessness. *Curr. Biol.* 23, 799–803. (doi:10.1016/j.cub.2013.03.054)
- 44.
- Cooper ML, Frone MR, Russell M, Mudar P. 1995 Drinking to regulate positive and negative emotions: a motivational model of alcohol use. *J. Pers. Soc. Psychol.* 69, 990–1005. (doi:10.1037/0022-3514.69.5.990)
- 45.
- Schrieks IC, Stafleu A, Kallen VL, Grootjen M, Witkamp RF, Hendriks HFJ. 2014 The biphasic effects of moderate alcohol consumption with a meal on ambiance-induced mood and autonomic nervous system balance: a randomized crossover trial. *PLoS One* 9, e86199. (doi:10.1371/journal.pone.0086199)
- 46.
- Quirk GJ, Armony JL, LeDoux JE. 1997 Fear conditioning enhances different temporal components of tone-evoked spike trains in auditory cortex and lateral amygdala. *Neuron* 19, 613–624. (doi:10.1016/s0896-6273(00)80375-x)
- 47.
- Zhang Y, Ouyang K, Lipina TV, Wang H, Zhou Q. 2019 Conditioned stimulus presentations alter anxiety level in fear-conditioned mice. *Mol. Brain* 12, 28. (doi:10.1186/s13041-019-0445-4)
- 48.
- Le QSE, Hereford D, Borkar CD, Aldaco Z, Klar J, Resendez A, Fadok JP. 2024 Contributions of associative and non-associative learning to the dynamics of defensive ethograms. *eLife* 12, RP90414. (doi:10.7554/eLife.90414)
- 49.
- Aceves-Piña EO, Quinn WG. 1979 Learning in normal and mutant *Drosophila* larvae. *Science* 206, 93–96. (doi:10.1126/science.206.4414.93)
- 50.

- Eschbach C, Cano C, Haberkern H, Schraut K, Guan C, Triphan T, Gerber B. 2011 Associative learning between odorants and mechanosensory punishment in larval *Drosophila*. *J. Exp. Biol.* 214, 3897–3905. (doi:10.1242/jeb.060533)
- 51.
- Khurana S, Abubaker MB, Siddiqi O. 2009 Odour avoidance learning in the larva of *Drosophila melanogaster*. *J. Biosci.* 34, 621–631. (doi:10.1007/s12038-009-0080-9)
- 52.
- Quinn WG, Harris WA, Benzer S. 1974 Conditioned behavior in *Drosophila melanogaster*. *Proc. Natl Acad. Sci. USA* 71, 708–712. (doi:10.1073/pnas.71.3.708)
- 53.
- Tully T, Quinn WG. 1985 Classical conditioning and retention in normal and mutant *Drosophila melanogaster*. *J. Comp. Physiol. Sens. Neural Behav. Physiol* 157, 263–277. (doi:10.1007/BF01350033)
- 54.
- Vergoz V, Roussel E, Sandoz JC, Giurfa M. 2007 Aversive learning in honeybees revealed by the olfactory conditioning of the sting extension reflex. *PLoS One* 2, e288. (doi:10.1371/journal.pone.0000288)
- 55.
- Dawson EH, Chittka L, Leadbeater E. 2016 Alarm substances induce associative social learning in honeybees, *Apis mellifera*. *Anim. Behav.* 122, 17–22. (doi:10.1016/j.anbehav.2016.08.006)
- 56.
- Black TE, Fofah O, Dinges CW, Ortiz-Alvarado CA, Avalos A, Ortiz-Alvarado Y, Abramson CI. 2021 Effects of aversive conditioning on expression of physiological stress in honey bees (*Apis mellifera*). *Neurobiol. Learn. Mem.* 178, 107363. (doi:10.1016/j.nlm.2020.107363)
- 57.
- Barrios N, Farias M, Moita MA. 2021 Threat induces cardiac and metabolic changes that

negatively impact survival in flies. *Curr. Biol.* 31, 5462–5472.(doi:10.1016/j.cub.2021.10.

013)

58.

Pasquini E, Brouwer J, Di Rollo V, Baracchi D, Messina A, Frasnelli E. 2025 Central GABAergic neuromodulation of nocifensive behaviors in bumble bees. *iScience* 28, 112024. (doi:10.1016/j.isci.2025.112024)

59.

Jang W, Oh M, Cho EH, Baek M, Kim C. 2023 *Drosophila* pain sensitization and modulation unveiled by a novel pain model and analgesic drugs. *PLoS One* 18, e0281874. (doi:10.1371/journal.pone.0281874)

60.

Gibbons M, Crump A, Barrett M, Sarlak S, Birch J, Chittka L. 2022 Chapter three - Can insects feel pain? A review of the neural and behavioural evidence. In *Advances in insect physiology* (ed. R Jurenka), pp. 155–229. Cambridge, MA: Academic Press. (doi:10.1016/bs.aiip.2022.10.001)

61.

Wu MS, Liao TW, Wu CY, Hsieh TH, Kuo PC, Li YC, Cheng KC, Chiang HC. 2023 Aversive conditioning information transmission in *Drosophila*. *Cell Rep.* 42, 113207. (doi:10.1016/j.celrep.2023.113207)

62.

Mendl M, Burman OHP, Parker RMA, Paul ES. 2009 Cognitive bias as an indicator of animal emotion and welfare: emerging evidence and underlying mechanisms. *Appl. Anim. Behav. Sci.* 118, 161–181. (doi:10.1016/j.applanim.2009.02.023)

63.

Solvi C, Baciadonna L, Chittka L. 2016 Unexpected rewards induce dopamine-dependent positive emotion-like state changes in bumblebees. *Science* 353, 1529–1531. (doi:10.1126/science.aaf4454)

64.

Galpayage Dona HS, Solvi C, Kowalewska A, Mäkelä K, MaBouDi H, Chittka L. 2022 Do bumble bees play? *Anim. Behav.* 194, 239–251. (doi:10.1016/j.anbehav.2022.08.013)

65.

Bateson M, Desire S, Gartside SE, Wright GA. 2011 Agitated honeybees exhibit pessimistic cognitive biases. *Curr. Biol.* 21, 1070–1073. (doi:10.1016/j.cub.2011.05.017)

66.

Baracchi D, Lihoreau M, Giurfa M. 2017 Do insects have emotions? Some insights from bumble bees. *Front. Behav. Neurosci.* 11, 157. (doi:10.3389/fnbeh.2017.00157)

67.

Procenko O, Read JCA, Nityananda V. 2024 Physically stressed bees expect less reward in an active choice judgement bias test. *Proc. R. Soc. B* 291, 20240512. (doi:10.1098/rspb.2024.0512)

68.

Triphan T, Ferreira CH, Huetteroth W. 2025 Play-like behavior exhibited by the vinegar fly *Drosophila melanogaster*. *Curr. Biol* 35, 1145–1155. (doi:10.1016/j.cub.2025.01.025)

69.

Eisenstein EM, Carlson AD. 1997 A comparative approach to the behavior called ‘learned helplessness’. *Behav. Brain Res.* 86, 149–160. (doi:10.1016/s0166-4328(96)02260-7)

70.

Eisenstein EM, Carlson AD, Harris JT. 1997 A ganglionic model of ‘learned helplessness’. *Integr. Physiol. Behav. Sci.* 32, 265–271. (doi:10.1007/BF02688624)

71.

Van De Poll MN, van Swinderen B. 2021 Balancing prediction and surprise: a role for active sleep at the dawn of consciousness? *Front. Syst. Neurosci.* 15, 768762. (doi:10.3389/fnsys.2021.768762)

72.

Eibl-Eibesfeldt I. 1973 Der vorprogrammierte Mensch: Das Ererbte als bestimmender Faktor im menschlichen Verhalten. Vienna, Austria: Molden.

73.

LeDoux JE. 1998 The emotional brain: the mysterious underpinnings of emotional life. New York, NY: Simon and Schuster. (doi:10.1176/ajp.155.4.57)

11

royalsocietypublishing.org/journal/rstb

Phil. Trans. R. Soc. B 380: rstb.2024.0302

74.

Panksepp J. 2004 Affective neuroscience: the foundations of human and animal emotions. Oxford, UK: Oxford University Press.

75.

Darwin CR. 1998 The expression of the emotions in man and animals, 1st edn. London, UK: John Murray. (doi:10.1093/oso/9780195112719.001.0001)

76.

Chittka L, Wilson C. 2019 Expanding consciousness. Am. Sci. 107, 364. (doi:10.1511/2019.107.6.364)

77.

Godfrey-Smith P. 2016 Other minds: the octopus and the evolution of intelligent life. London, UK: William Collins.

78.

Sperry RW. 1950 Neural basis of the spontaneous optokinetic response produced by visual inversion. J. Comp. Physiol. Psychol. 43, 482–489. (doi:10.1037/h0055479)

79.

Poulet JFA, Hedwig B. 2006 The cellular basis of a corollary discharge. Science 311, 518–522. (doi:10.1126/science.1120847)

80.

- Brebner J, Chittka L. 2021 Animal cognition: the self-image of a bumblebee. *Curr. Biol.* 31, R207–R209. (doi:10.1016/j.cub.2020.12.027)
- 81.
- De Agrò M, Galpayage Dona HS, Vallortigara G. 2024 Seeing life in the teeming world: animacy perception in arthropods. *Front. Psychol.* 15, 1492239. (doi:10.3389/fpsyg.2024.1492239)
- 82.
- Nowbahari E, Scohier A, Durand JL, Hollis KL. 2009 Ants, *Cataglyphis cursor*, use precisely directed rescue behavior to free entrapped relatives. *PLoS One* 4, e6573. (doi:10.1371/journal.pone.0006573)
- 83.
- Chittka L, Rossi N. 2022 Social cognition in insects. *Trends Cogn. Sci.* 26, 578–592. (doi:10.1016/j.tics.2022.04.001)
- 84.
- Vasconcelos M, Hollis K, Nowbahari E, Kacelnik A. 2012 Pro-sociality without empathy. *Biol. Lett.* 8, 910–912. (doi:10.1098/rsbl.2012.0554)
- 85.
- Ravi S, Siesenop T, Bertrand O, Li L, Doussot C, Warren WH, Combes SA, Egelhaaf M. 2020 Bumblebees perceive the spatial layout of their environment in relation to their body size and form to minimize inflight collisions. *Proc. Natl Acad. Sci. USA* 117, 31494–31499. (doi:10.1073/pnas.2016872117)
- 86.
- Riojas-Schnier JL, Toth AL. 2022 Insights into insect cognition through mirror-induced behaviour in paper wasps. *Behaviour* 159, 735–758. (doi:10.1163/1568539x-bja10148)
- 87.
- Ben-Nun A, Guershon M, Ayali A. 2013 Self body-size perception in an insect. *Naturwissenschaften* 100, 479–484. (doi:10.1007/s00114-013-1042-5)

88.

Gallup G, Anderson JR, Shillito DJ. 2002 The mirror test. In *The cognitive animal: empirical and theoretical perspectives on animal cognition* (eds M Bekoff, C Allen, GM Burghardt), pp. 325–333. Cambridge, MA: MIT Press. (doi:10.7551/mitpress/1885.003.0046)

89.

Gallup GG, Anderson JR. 2020 Self-recognition in animals: where do we stand 50 years later? Lessons from cleaner wrasse and other species. *Psychol. Conscious.* 7, 46–58. (doi:10.1037/cns0000206)

90.

Clary D, Kelly DM. 2016 Graded mirror self-recognition by Clark's nutcrackers. *Sci. Rep.* 6, 36459. (doi:10.1038/srep36459)

91.

Plotnik JM, De Waal FBM, Reiss D. 2006 Self-recognition in an Asian elephant. *Proc. Natl Acad. Sci. USA* 103, 17053–17057. (doi:10.1073/pnas.0608062103)

92.

Cammaerts MC, Cammaerts R. 2015 Are ants (Hymenoptera, Formicidae) capable of self recognition? *J. Sci.* 5, 521–532.

93.

Sheehan MJ, Tibbetts EA. 2011 Specialized face learning is associated with individual recognition in paper wasps. *Science* 334, 1272–1275. (doi:10.1126/science.1211334)

94.

Swartz KB. 1997 What is mirror self-recognition in nonhuman primates, and what is it not? *Ann. N. Y. Acad. Sci.* 818, 65–71. (doi:10.1111/j.1749-6632.1997.tb48246.x)

95.

McFarland D. 2009 *Guilty robots, happy dogs: the question of alien minds*. Oxford, UK: Oxford University Press. (doi:10.1093/oso/9780199219292.001.0001)

96.

- Mason GJ, Lavery JM. 2022 What is it like to be a bass? Red herrings, fish pain and the study of animal sentience. *Front. Vet. Sci.* 9, 788289. (doi:10.3389/fvets.2022.788289)
- 97.
- Kohda M, Sogawa S, Jordan AL, Kubo N, Awata S, Satoh S, Kobayashi T, Fujita A, Bshary R. 2022 Further evidence for the capacity of mirror self-recognition in cleaner fish and the significance of ecologically relevant marks. *PLoS Biol.* 20, e3001529. (doi:10.1371/journal.pbio.3001529)
- 98.
- D'Ettorre P, Heinze J. 2005 Individual recognition in ant queens. *Curr. Biol.* 15, 2170–2174. (doi:10.1016/j.cub.2005.10.067)
- 99.
- Tsutsui ND. 2004 Scents of self: the expression component of self/non-self recognition systems. *Ann. Zool. Fenn* 41, 713–727. <https://www.jstor.org/stable/23736139>
100. Capodeanu-Nägler A, Rapkin J, Sakaluk SK, Hunt J, Steiger S. 2014 Self-recognition in crickets via on-line processing. *Curr. Biol.* 24, R1117–R1118. (doi:10.1016/j.cub.2014.10.050)
101. Terrace HS, Son LK. 2009 Comparative metacognition. *Curr. Opin. Neurobiol.* 19, 67–74. (doi:10.1016/j.conb.2009.06.004)
102. Czaczkes TJ. 2022 Advanced cognition in ants. *Myrmecol. News* 32, 51–64. (doi:10.25849/myrmecol.news_032:051)
103. Yeung N, Summerfield C. 2012 Metacognition in human decision-making: confidence and error monitoring. *Phil. Trans. R. Soc. B* 367, 1310–1321. (doi:10.1098/rstb.2011.0416)
104. Tomasello M. 2023 Social cognition and metacognition in great apes: a theory. *Anim. Cogn.* 26, 25–35. (doi:10.1007/s10071-022-01662-0)
105. Perry CJ, Barron AB. 2013 Honey bees selectively avoid difficult choices. *Proc. Natl Acad. Sci. USA* 110, 19155–19159. (doi:10.1073/pnas.1314571110)
106. Perry CJ, Barron AB, Chittka L. 2017 The frontiers of insect cognition. *Curr. Opin. Behav. Sci.* 16, 111–118. (doi:10.1016/j.cobeha.2017.05.011)

107. Giurfa M. 2015 Learning and cognition in insects. *WIREs Cogn. Sci.* 6, 383–395. (doi:10.1002/wcs.1348)
108. Fleming SM, Lau HC. 2014 How to measure metacognition. *Front. Hum. Neurosci.* 8, 443. (doi:10.3389/fnhum.2014.00443)
109. Menzel R. 2019 Search Strategies for Intentionality in the Honeybee Brain. In *The Oxford Handbook of Invertebrate Neurobiology* (ed. J Byrne), pp. 663–684. Oxford, UK: Oxford University Press. (doi:10.1093/oxfordhb/9780190456757.013.27)
110. Smith JD. 2009 The study of animal metacognition. *Trends Cogn. Sci.* 13, 389–396. (doi:10.1016/j.tics.2009.06.009)
111. Brembs B. 2011 Towards a scientific concept of free will as a biological trait: spontaneous actions and decision-making in invertebrates. *Proc. R. Soc. B* 278, 930–939. (doi:10.1098/rspb.2010.2325)
112. Juusola M, Takalo J, Kemppainen J, Haghighi KR, Scales B, McManus J, Bridges A, MaBouDi H, Chittka L. 2025 Theory of morphodynamic information processing: linking sensing to behaviour. *Vision Res.* 227, 108537. (doi:10.1016/j.visres.2024.108537)
113. Feldman H, Friston KJ. 2010 Attention, uncertainty, and free-energy. *Front. Hum. Neurosci.* 4, 215. (doi:10.3389/fnhum.2010.00215)
114. Von Helmholtz H. 1925 Helmholtz's treatise on physiological optics. (ed. JPC Southall). Washington, DC: Optica Publishing Group. (doi:10.1037/13536-000)
115. Friston K. 2005 A theory of cortical responses. *Phil. Trans. R. Soc. B* 360, 815–836. (doi:10.1098/rstb.2005.1622)
116. Rao RPN, Ballard DH. 1999 Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. *Nat. Neurosci.* 2, 79–87. (doi:10.1038/4580)
117. Clark A. 2013 Whatever next? Predictive brains, situated agents, and the future of cognitive science. *Behav. Brain Sci.* 36, 181–204. (doi:10.1017/s0140525x12000477)

118. Hohwy J. 2013 The predictive mind. Oxford, UK: Academic. (doi:10.1093/acprof:oso/9780199682737.001.0001)
119. Jürgensen AM, Sakagiannis P, Schleyer M, Gerber B, Nawrot MP. 2024 Prediction error drives associative learning and conditioned behavior in a spiking model of *Drosophila* larva. *iScience* 27, 108640. (doi:10.1016/j.isci.2023.108640)
120. Srinivasan MV, Laughlin SB, Dubs A, Horridge GA. 1982 Predictive coding: a fresh view of inhibition in the retina. *Proc. R. Soc. Lond. B* 216, 427–459. (doi:10.1098/rspb.1982.0085)
121. Terao K, Mizunami M. 2017 Roles of dopamine neurons in mediating the prediction error in aversive learning in insects. *Sci. Rep.* 7, 14694. (doi:10.1038/s41598-017-14473-y)
122. Webb B. 2004 Neural mechanisms for prediction: do insects have forward models? *Trends Neurosci.* 27, 278–282. (doi:10.1016/j.tins.2004.03.004)
123. Chittka L, Ings TC, Raine NE. 2004 Chance and adaptation in the evolution of island bumblebee behaviour. *Popul. Ecol.* 46, 243–251. (doi:10.1007/s10144-004-0180-1)
124. Chittka L, Spaethe J. 2007 Visual search and the importance of time in complex decision making by bees. *Arthropod Plant Interact.* 1, 37–44. (doi:10.1007/s11829-007-9001-8)
125. Zhang SW, Srinivasan MV. 1994 Prior experience enhances pattern discrimination in insect vision. *Nature* 368, 330–332. (doi:10.1038/368330a0)

12

royalsocietypublishing.org/journal/rstb

Phil. Trans. R. Soc. B 380: rstb.2024.0302

126. Engel AK, Fries P. 2016 Neuronal oscillations, coherence, and consciousness. In *The neurology of consciousness* (eds S Laureys, O Gosseries, G Tononi), pp. 49–60. Cambridge, MA: Academic. (doi:10.1016/b978-0-12-800948-2.00003-0)
127. Gallot S, Sack AT, Schuhmann T, de Graaf TA. 2017 Oscillatory correlates of visual consciousness. *Front. Psychol.* 8, 1147. (doi:10.3389/fpsyg.2017.01147)
128. Engel AK, Singer W. 2001 Temporal binding and the neural correlates of sensory awareness. *Trends Cogn. Sci.* 5, 16–25. (doi:10.1016/s1364-6613(00)01568-0)

129. Popov T, Szyszka P. 2020 Alpha oscillations govern interhemispheric spike timing coordination in the honey bee brain. *Proc. R. Soc. B* 287, 20200115. (doi:10.1098/rspb.2020.0115)
130. Sauseng P, Klimesch W, Schabus M, Doppelmayr M. 2005 Fronto-parietal EEG coherence in theta and upper alpha reflect central executive functions of working memory. *Int. J. Psychophysiol.* 57, 97–103. (doi:10.1016/j.ijpsycho.2005.03.018)
131. Cavanagh JF, Frank MJ. 2014 Frontal theta as a mechanism for cognitive control. *Trends Cogn. Sci.* 18, 414–421. (doi:10.1016/j.tics.2014.04.012)
132. Grabowska MJ, Jeans R, Steeves J, van Swinderen B. 2020 Oscillations in the central brain of *Drosophila* are phase locked to attended visual features. *Proc. Natl Acad. Sci. USA* 117, 29925–29936. (doi:10.1073/pnas.2010749117)
133. Chao ZC, Takaura K, Wang L, Fujii N, Dehaene S. 2018 Large-scale cortical networks for hierarchical prediction and prediction error in the primate brain. *Neuron* 100, 1252–1266. (doi:10.1016/j.neuron.2018.10.004)
134. Johnston P, Robinson J, Kokkinakis A, Ridgeway S, Simpson M, Johnson S, Kaufman J, Young AW. 2017 Temporal and spatial localization of prediction-error signals in the visual brain. *Biol. Psychol.* 125, 45–57. (doi:10.1016/j.biopsycho.2017.02.004)
135. Van De Poll MN. 2024 Electrophysiological correlates of predictive processing in the fly brain and their modulation during sleep. PhD Dissertation, Queensland Brain Institute, The University of Queensland. (doi:10.14264/95792f5)
136. Zhao C, Widmer YF, Diegelmann S, Petrovici MA, Sprecher SG, Senn W. 2021 Predictive olfactory learning in *Drosophila*. *Sci. Rep.* 11, 6795. (doi:10.1038/s41598-021-85841-y)
137. Bekinschtein TA, Dehaene S, Rohaut B, Tadel F, Cohen L, Naccache L. 2009 Neural signature of the conscious processing of auditory regularities. *Proc. Natl Acad. Sci. USA* 106, 1672–1677. (doi:10.1073/pnas.0809667106)
138. Gabhart KM, Xiong YS, Bastos AM. 2025 Predictive coding: a more cognitive process than

we thought? Trends Cogn. Sci. 29, 627–640. (doi:10.1016/j.tics.2025.01.012)

139. Paulk AC, Stacey JA, Pearson TWJ, Taylor GJ, Moore RJD, Srinivasan MV, van Swinderen B. 2014 Selective attention in the honeybee optic lobes precedes behavioral choices. Proc. Natl Acad. Sci. USA 111, 5006–5011. (doi:10.1073/pnas.1323297111)

140. Kirszenblat L, van Swinderen B. 2015 The yin and yang of sleep and attention. Trends Neurosci. 38, 776–786. (doi:10.1016/j.tins.2015.10.001)

141. Dijk DJ. 2009 Regulation and functional correlates of slow wave sleep. J. Clin. Sleep Med. 5, S6–S15. (doi:10.5664/jcsm.5.2s.s6)

142. Blumberg MS, Lesku JA, Libourel PA, Schmidt MH, Rattenborg NC. 2020 What is REM sleep? Curr. Biol. 30, R38–R49. (doi:10.1016/j.cub.2019.11.045)

143. Andrillon T, Nir Y, Cirelli C, Tononi G, Fried I. 2015 Single-neuron activity and eye movements during human REM sleep and awake vision. Nat. Commun. 6, 7884. (doi:10.1038/ncomms8884)

144. Nielsen TA, Deslauriers D, Baylor GW. 1991 Emotions in dream and waking event reports. Dreaming 1, 287–300. (doi:10.1037/h0094340)

145. Galbiati A, Sforza M, Fasiello E, Casoni F, Marrella N, Leitner C, Zucconi M, Ferini-Strambi L. 2020 The association between emotional dysregulation and REM sleep features in insomnia disorder. Brain Cogn. 146, 105642. (doi:10.1016/j.bandc.2020.105642)

146. Boyce R, Glasgow SD, Williams S, Adamantidis A. 2016 Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation. Science 352, 812–816. (doi:10.1126/science.aad5252)

147. Hobson JA, Hong CCH, Friston KJ. 2014 Virtual reality and consciousness inference in dreaming. Front. Psychol. 5, 1133. (doi:10.3389/fpsyg.2014.01133)

148. Hoel E. 2021 The overfitted brain: dreams evolved to assist generalization. Patterns 2, 100244. (doi:10.1016/j.patter.2021.100244)

149. Llewellyn S. 2016 Dream to predict? REM dreaming as prospective coding. Front. Psychol. 6, 1961. (doi:10.3389/fpsyg.2015.01961)

150. Ajayi OM, Marlman JM, Gleitz LA, Smith ES, Piller BD, Krupa JA, Vinauger C, Benoit JB. 2022 Behavioral and postural analyses establish sleep-like states for mosquitoes that can impact host landing and blood feeding. *J. Exp. Biol.* 225, jeb244032. (doi:10.1242/jeb.244032)
151. Eban-Rothschild AD, Bloch G. 2008 Differences in the sleep architecture of forager and young honeybees (*Apis mellifera*). *J. Exp. Biol.* 211, 2408–2416. (doi:10.1242/jeb.016915)
152. Hendricks JC, Finn SM, Panckeri KA, Chavkin J, Williams JA, Sehgal A, Pack AI. 2000 Rest in *Drosophila* is a sleep-like state. *Neuron* 25, 129–138. (doi:10.1016/s0896-6273(00)80877-6)
153. Kaiser W, Steiner-Kaiser J. 1983 Neuronal correlates of sleep, wakefulness and arousal in a diurnal insect. *Nature* 301, 707–709. (doi:10.1038/301707a0)
154. Sauer S, Kinkelin M, Herrmann E, Kaiser W. 2003 The dynamics of sleep-like behaviour in honey bees. *J. Comp. Physiol.* 189, 599–607. (doi:10.1007/s00359-003-0436-9)
155. Shaw PJ, Cirelli C, Greenspan RJ, Tononi G. 2000 Correlates of sleep and waking in *Drosophila melanogaster*. *Science* 287, 1834–1837. (doi:10.1126/science.287.5459.1834)
156. Kaiser W. 1988 Busy bees need rest, too: behavioural and electromyographical sleep signs in honeybees. *J. Comp. Physiol. A* 163, 565–584. (doi:10.1007/BF00603841)
157. Frighetto G, Cellini N. 2017 Sleep and memory consolidation: a common mechanism across species? *J. Neurophysiol.* 117, 1–3. (doi:10.1152/jn.00213.2016)
158. Zwaka H, Bartels R, Gora J, Franck V, Culo A, Götsch M, Menzel R. 2015 Context odor presentation during sleep enhances memory in honeybees. *Curr. Biol.* 25, 2869–2874. (doi:10.1016/j.cub.2015.09.069)
159. van Alphen B, Yap MHW, Kirszenblat L, Kottler B, van Swinderen B. 2013 A dynamic deep sleep stage in *Drosophila*. *J. Neurosci.* 33, 6917–6927. (doi:10.1523/jneurosci.0061-13.2013)
160. Jagannathan SR, Jeans T, Van De Poll MN, van Swinderen B. 2024 Multivariate classification of multichannel long-term electrophysiology data identifies different sleep stages in fruit flies. *Sci. Adv.* 10, eadj4399. (doi:10.1126/sciadv.adj4399)

161. Wiggin TD, Goodwin PR, Donelson NC, Liu C, Trinh K, Sanyal S, Griffith LC. 2020 Covert sleep-related biological processes are revealed by probabilistic analysis in *Drosophila*. *Proc. Natl Acad. Sci. USA* 117, 10024–10034. (doi:10.1073/pnas.1917573117)
162. van Swinderen B, Van De Poll M. 2025 Long-term multichannel recordings in *Drosophila* flies reveal distinct responses to deviant visual stimuli during sleep compared to wake. *J. Exp. Biol.* (doi:10.1242/jeb.250165)
163. Yap MHW, Grabowska MJ, Rohrscheib C, Jeans R, Troup M, Paulk AC, van Alphen B, Shaw PJ, van Swinderen B. 2017 Oscillatory brain activity in spontaneous and induced sleep stages in flies. *Nat. Commun.* 8, 1815. (doi:10.1038/s41467-017-02024-y)
164. Tainton-Heap LAL, Kirszenblat LC, Notaras ET, Grabowska MJ, Jeans R, Feng K, Shaw PJ, van Swinderen B. 2021 A paradoxical kind of sleep in *Drosophila melanogaster*. *Curr. Biol.* 31, 578–590. (doi:10.1016/j.cub.2020.10.081)
165. Stanhope BA, Jaggard JB, Gratton M, Brown EB, Keene AC. 2020 Sleep regulates glial plasticity and expression of the engulfment receptor draper following neural injury. *Curr. Biol.* 30, 1092–1101. (doi:10.1016/j.cub.2020.02.057)
166. van Alphen B, Semenza ER, Yap M, van Swinderen B, Allada R. 2021 A deep sleep stage in *Drosophila* with a functional role in waste clearance. *Sci. Adv.* 7, eabc2999. (doi:10.1126/sciadv.abc2999)
- 13
- royalsocietypublishing.org/journal/rstb
- Phil. Trans. R. Soc. B 380: rstb.2024.0302
167. Fultz NE, Bonmassar G, Setsompop K, Stickgold RA, Rosen BR, Polimeni JR, Lewis LD. 2019 Coupled electrophysiological, hemodynamic, and cerebrospinal fluid oscillations in human sleep. *Science* 366, 628–631. (doi:10.1126/science.aax5440)
168. Xie L et al. 2013 Sleep drives metabolite clearance from the adult brain. *Science* 342, 373–377. (doi:10.1126/science.1241224)
169. de Bivort BL, van Swinderen B. 2016 Evidence for selective attention in the insect brain.

Curr. Opin. Insect Sci. 15, 9–15. (doi:10.1016/j.cois.2016.02.007)

170. Honkanen A, Adden A, da Silva Freitas J, Heinze S. 2019 The insect central complex and the neural basis of navigational strategies. *J. Exp. Biol.* 222, jeb188854. (doi:10.1242/jeb.188854)

171. Liu G, Seiler H, Wen A, Zars T, Ito K, Wolf R, Heisenberg M, Liu L. 2006 Distinct memory traces for two visual features in the *Drosophila* brain. *Nature* 439, 551–556. (doi:10.1038/nature04381)

172. Troup M et al. 2018 Acute control of the sleep switch in *Drosophila* reveals a role for gap junctions in regulating behavioral responsiveness. *eLife* 7, e37105. (doi:10.7554/elife.37105)

173. Kirszenblat L, Yaun R, van Swinderen B. 2019 Visual experience drives sleep need in *Drosophila*. *Sleep* 42, zsz102. (doi:10.1093/sleep/zsz102)

174. French AS, Geissmann Q, Beckwith EJ, Gilestro GF. 2021 Sensory processing during sleep in *Drosophila melanogaster*. *Nature* 598, 479–482. (doi:10.1038/s41586-021-03954-w)

175. Kirszenblat L, Ertekin D, Goodsell J, Zhou Y, Shaw PJ, van Swinderen B. 2018 Sleep regulates visual selective attention in *Drosophila*. *J. Exp. Biol.* 221, jeb191429. (doi:10.1242/jeb.191429)

176. Dijkstra N, Fleming SM. 2023 Subjective signal strength distinguishes reality from imagination. *Nat. Commun.* 14, 1627. (doi:10.1038/s41467-023-37322-1)

177. Dehaene S, Changeux JP, Naccache L, Sackur J, Sergent C. 2006 Conscious, preconscious, and subliminal processing: a testable taxonomy. *Trends Cogn. Sci.* 10, 204–211. (doi:10.1016/j.tics.2006.03.007)

178. Tononi G. 2004 An information integration theory of consciousness. *BMC Neurosci.* 5, 42. (doi:10.1186/1471-2202-5-42)

179. Dehaene S. 2001 Towards a cognitive neuroscience of consciousness: basic evidence and a workspace framework. *Cognition* 79, 1–37. (doi:10.1016/s0010-0277(00)00123-2)

180. Chittka L. 2022 The mind of a bee. Princeton, NJ: Princeton University Press.

(doi:10.1515/9780691236247)

181. Heisenberg M. 2015 Outcome learning, outcome expectations, and intentionality in *Drosophila*. *Learn. Mem.* 22, 294–298. (doi:10.1101/lm.037481.114)
182. Carter O, van Swinderen B, Leopold DA, Collin SP, Maier A. 2020 Perceptual rivalry across animal species. *J. Comp. Neurol.* 528, 3123–3133. (doi:10.1002/cne.24939)
183. Perry CJ, Chittka L. 2019 How foresight might support the behavioral flexibility of arthropods. *Curr. Opin. Neurobiol.* 54, 171–177. (doi:10.1016/j.conb.2018.10.014)
184. Avarguès-Weber A, Giurfa M. 2013 Conceptual learning by miniature brains. *Proc. R. Soc. B* 280, 20131907. (doi:10.1098/rspb.2013.1907)
185. Solvi C, Gutierrez Al-Khudhairy S, Chittka L. 2020 Bumble bees display cross-modal object recognition between visual and tactile senses. *Science* 367, 910–912. (doi:10.1126/science.aay8064)
186. Grover D, Chen JY, Xie J, Li J, Changeux JP, Greenspan RJ. 2022 Differential mechanisms underlie trace and delay conditioning in *Drosophila*. *Nature* 603, 302–308. (doi:10.1038/s41586-022-04433-6)
187. Skorupski P, Chittka L. 2006 Animal cognition: an insect's sense of time? *Curr. Biol.* 16, R851–R853. (doi:10.1016/j.cub.2006.08.069)
188. Field J, Savill C, Foster WA. 2025 Memory and the scheduling of parental care in an insect population in the wild. *Curr. Biol.* 35, 2740–2745. (doi:10.1016/j.cub.2025.04.045)
189. Crystal JD. 2025 Episodic memory in nonhumans: an approach to understand an evolutionary function of consciousness. *Phil. Trans. R. Soc. B* 380, 20240304. (doi:10.1098/rstb.2024.0304)
190. Fauria K, Colborn M, Collett TS. 2000 The binding of visual patterns in bumblebees. *Curr. Biol.* 10, 935–938. (doi:10.1016/s0960-9822(00)00623-0)
191. Sneddon LU, Elwood RW, Adamo SA, Leach MC. 2014 Defining and assessing animal pain. *Anim. Behav.* 97, 201–212. (doi:10.1016/j.anbehav.2014.09.007)
192. Moncoucy L, Dołęga K, Tallon-Baudry C, Cleeremans A. 2025 The value of consciousness:

experiences worth having. *Phil. Trans. R. Soc. B* 380, 20240303. (doi:10.1098/rstb.2024.

0303)

193. Baars BJ. 1997 *In the theater of consciousness: the workspace of the mind*. Oxford, UK:

Oxford University Press. (doi:10.1093/acprof:oso/9780195102659.001.1)

194. Grossberg S. 2021 Attention: Multiple types, brain resonances, psychological functions, and conscious states. *J. Integr. Neurosci* 20, 197–232. (doi:10.31083/j.jin.2021.01.406)

195. Tyng CM, Amin HU, Saad MNM, Malik AS. 2017 The influences of emotion on learning and memory. *Front. Psychol.* 8, 235933. (doi:10.3389/fpsyg.2017.01454)

14

royalsocietypublishing.org/journal/rstb

Phil. Trans. R. Soc. B 380: rstb.2024.0302