

# Tuples and Permutations

---

Vladimir Podolskii

Computer Science Department, Higher School of Economics

# Outline

Number of Tuples

Licence Plates

Tuples with Restrictions

Permutations

# Number of Passwords

## Problem

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

# Number of Passwords

## Problem

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

- It turns out that the rule of product is all we need to solve this problem

# Number of Passwords

## Problem

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

- It turns out that the rule of product is all we need to solve this problem
- But we need to do it step by step

# Number of Passwords

- Let's start with 1 letter password

\*

# Number of Passwords

- Let's start with 1 letter password
- Clearly, then the answer is 26

\*

# Number of Passwords

- Let's start with 1 letter password
- Clearly, then the answer is 26
- What about two letters?

\* \*



# Number of Passwords

- Let's start with 1 letter password
- Clearly, then the answer is 26
- What about two letters?
- Then we can choose both letters in 26 ways

26    26

\*    \*

# Number of Passwords

- Let's start with 1 letter password
- Clearly, then the answer is 26
- What about two letters?
- Then we can choose both letters in 26 ways
- Use the rule of product: the answer is 676

$$26 \times 26 = 676$$

\*     \*

# Number of Passwords

- Let's move on to the case of 3 letters

\* \* \*

# Number of Passwords

- Let's move on to the case of 3 letters
- We already know that we can choose the first two letters in 676 ways

676

\* \* \*

# Number of Passwords

- Let's move on to the case of 3 letters
- We already know that we can choose the first two letters in 676 ways
- We apply the rule of product again!

$$676 \times 26$$

$$\begin{array}{|c|c|c|} \hline * & * & * \\ \hline \end{array}$$

# Number of Passwords

- Let's move on to the case of 3 letters
- We already know that we can choose the first two letters in 676 ways
- We apply the rule of product again!
- And the answer is 17 576

$$676 \times 26 = 17\,576$$

\* \* \*

# Number of Passwords

- Now we can proceed in the same way

$$26 \times 26 \times 26$$

\* \* \* \*

# Number of Passwords

- Now we can proceed in the same way

$$26 \times 26 \times 26 \times 26$$

$$* \quad * \quad * \quad * \quad *$$



# Number of Passwords

- Now we can proceed in the same way

$$\begin{array}{ccccccccc} 26 & \times & 26 & \times & 26 & \times & 26 & \times & 26 \\ * & & * & & * & & * & & * \end{array}$$

# Number of Passwords

- Now we can proceed in the same way
- And the answer is 11 881 376

$$\begin{array}{ccccccccc} 26 & \times & 26 & \times & 26 & \times & 26 & \times & 26 & = & 11\,881\,376 \\ * & & * & & * & & * & & * & & \end{array}$$

# Number of Tuples

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

These sequences are usually called **tuples**

# Number of Tuples

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

These sequences are usually called **tuples**

- Can apply the same argument

# Number of Tuples

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

These sequences are usually called **tuples**

- Can apply the same argument
- There are  $n$  possibilities to pick the first letter

# Number of Tuples

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

These sequences are usually called **tuples**

- Can apply the same argument
- There are  $n$  possibilities to pick the first letter
- Each next letter multiplies the number of sequences by  $n$

# Number of Tuples

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

These sequences are usually called **tuples**

- Can apply the same argument
- There are  $n$  possibilities to pick the first letter
- Each next letter multiplies the number of sequences by  $n$
- Thus the answer is a product of  $n$  by itself  $k$  times, that is  $n^k$

# Outline

Number of Tuples

**Licence Plates**

Tuples with Restrictions

Permutations



# Licence Plates



[wikimedia.org](https://commons.wikimedia.org/wiki/File:Russian_license_plate_C_065_MK_78.jpg)

- Now we are ready to get back to our motivating example

# Licence Plates



[wikimedia.org](https://commons.wikimedia.org/wiki/File:Russian_license_plate_78_C065MK.jpg)

- Now we are ready to get back to our motivating example
- Russian license plate: 3 digits, 3 letters; 78 is a regional code

# Licence Plates



wikimedia.org

- Now we are ready to get back to our motivating example
- Russian license plate: 3 digits, 3 letters; 78 is a regional code
- We have: 10 options for digits, 12 options for letters (only those Cyrillic letters that are similar to Latin ones are used)

# Licence Plates



wikimedia.org

- Now we are ready to get back to our motivating example
- Russian license plate: 3 digits, 3 letters; 78 is a regional code
- We have: 10 options for digits, 12 options for letters (only those Cyrillic letters that are similar to Latin ones are used)
- How many plates are there for one region?

# Licence Plates



[wikimedia.org](https://commons.wikimedia.org/wiki/File:Russian_license_plate_78_C065MK.jpg)

- Each digit can be chosen in 10 ways

# Licence Plates



wikimedia.org

- Each digit can be chosen in 10 ways
- Thus a sequence of digits can be chosen in  $10 \times 10 \times 10 = 1\,000$  ways

# Licence Plates



wikimedia.org

- Each digit can be chosen in 10 ways
- Thus a sequence of digits can be chosen in  $10 \times 10 \times 10 = 1\,000$  ways
- Each letter can be chosen in 12 ways

# Licence Plates



wikimedia.org

- Each digit can be chosen in 10 ways
- Thus a sequence of digits can be chosen in  $10 \times 10 \times 10 = 1\,000$  ways
- Each letter can be chosen in 12 ways
- Thus a sequence of three letters can be chosen in  $12 \times 12 \times 12 = 1\,728$  ways



# Licence Plates



wikimedia.org

- Each digit can be chosen in 10 ways
- Thus a sequence of digits can be chosen in  $10 \times 10 \times 10 = 1\,000$  ways
- Each letter can be chosen in 12 ways
- Thus a sequence of three letters can be chosen in  $12 \times 12 \times 12 = 1\,728$  ways
- Overall, there are  $1\,728\,000$  license plates for a region

# Licence Plates



wikimedia.org

- There are 1 728 000 license plates for a region
- Is it enough?

# Licence Plates



wikimedia.org

- There are 1 728 000 license plates for a region
- Is it enough?
- No, it's not: for example, there are about 5 600 000 vehicles in Moscow (as of 2016)

# Licence Plates



wikimedia.org

- There are 1 728 000 license plates for a region
- Is it enough?
- No, it's not: for example, there are about 5 600 000 vehicles in Moscow (as of 2016)
- Does it mean that by the Pigeonhole principle there are vehicles with identical license plates?

# Licence Plates



[wikimedia.org](https://commons.wikimedia.org/wiki/File:Russian_license_plate_C_065_MK_78_RUS.jpg)

- No, several regional codes were introduced for the same region

# Licence Plates



[wikimedia.org](https://commons.wikimedia.org/wiki/File:Russian_license_plate_78_C065MK.jpg)

- No, several regional codes were introduced for the same region
- But this required introduction of three-digit regional codes

# Outline

Number of Tuples

Licence Plates

**Tuples with Restrictions**

Permutations

# Tuples with Restrictions

- We have shown how using the rule of product we can compute the number of tuples or a certain length of a fixed set of symbols



# Tuples with Restrictions

- We have shown how using the rule of product we can compute the number of tuples or a certain length of a fixed set of symbols
- But the rule of product can also give us other things too

# Numbers with exactly one digit 7

## Problem

How many integer numbers between 0 and 9999 are there that have exactly one 7 digit?

# Numbers with exactly one digit 7

## Problem

How many integer numbers between 0 and 9999 are there that have exactly one 7 digit?

- Numbers between 0 and 9999 are sequences of digits of length 4

# Numbers with exactly one digit 7

## Problem

How many integer numbers between 0 and 9999 are there that have exactly one 7 digit?

- Numbers between 0 and 9999 are sequences of digits of length 4
- Three digital numbers correspond to sequences starting with 0

# Numbers with exactly one 7 digit

\* \* \* \*

- We can place the unique 7 at any of four positions

# Numbers with exactly one 7 digit

\* \* \* \*

- We can place the unique 7 at any of four positions
- This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum

# Numbers with exactly one 7 digit

\* 7 \* \*

- We can place the unique 7 at any of four positions
- This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum
- Consider one of the cases

# Numbers with exactly one 7 digit

\* 7 \* \*

- We can place the unique 7 at any of four positions
- This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum
- Consider one of the cases
- Each of other three digits can be picked out of 9 options! (digit 7 is forbidden)



# Numbers with exactly one 7 digit

$$* \quad 7 \quad * \quad *$$

- Thus there are  $9 \times 9 \times 9 = 729$  sequences in this case

# Numbers with exactly one 7 digit

\* \* \* \*

- Thus there are  $9 \times 9 \times 9 = 729$  sequences in this case
- And in all other cases as well!

# Numbers with exactly one 7 digit

\* \* \* \*

- Thus there are  $9 \times 9 \times 9 = 729$  sequences in this case
- And in all other cases as well!
- There are 4 cases, so there are  $4 \times 729 = 2916$  four numbers below 10 000 with exactly one digit 7

# Numbers with exactly one 7 digit

\* \* \* \*

- Thus there are  $9 \times 9 \times 9 = 729$  sequences in this case
- And in all other cases as well!
- There are 4 cases, so there are  $4 \times 729 = 2916$  four numbers below 10 000 with exactly one digit 7
- This is below  $1/3$ , but above  $1/4$  of all four digit numbers

# Numbers with exactly one 7 digit

\* \* \* \*

- Thus there are  $9 \times 9 \times 9 = 729$  sequences in this case
- And in all other cases as well!
- There are 4 cases, so there are  $4 \times 729 = 2916$  four numbers below 10 000 with exactly one digit 7
- This is below  $1/3$ , but above  $1/4$  of all four digit numbers
- This is an estimation of the probability to get exactly one digit 7 if we pick a number below 10 000 “randomly”

# Outline

Number of Tuples

Licence Plates

Tuples with Restrictions

**Permutations**

# Permutations

- We have discussed how to count the number of tuples

# Permutations

- We have discussed how to count the number of tuples
- Now we are ready to proceed to the second standard combinatorial setting: **permutations**



# Permutations

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- Tuples of length  $k$  without repetitions are called  $k$ -permutations

# Permutations

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- Tuples of length  $k$  without repetitions are called  $k$ -permutations
- Observe that if  $n < k$ , then there are no  $k$ -permutations: there are simply not enough different letters

# Permutations

## Problem

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- Tuples of length  $k$  without repetitions are called  **$k$ -permutations**
- Observe that if  $n < k$ , then there are no  $k$ -permutations: there are simply not enough different letters
- So it is enough to solve the problem for the case  $k \leq n$

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \end{array}$$

- Let us apply rule of product

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n \end{array}$$

- Let us apply rule of product
- The first symbol can be picked in  $n$  ways

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n \end{array}$$

- Let us apply rule of product
- The first symbol can be picked in  $n$  ways
- How many choices there are for the second symbol?

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n \end{array}$$

- Let us apply rule of product
- The first symbol can be picked in  $n$  ways
- How many choices there are for the second symbol?
- We can place anything there, except for the symbol on the first place

# Permutations

$$\begin{array}{cccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & & & \end{array}$$

- Let us apply rule of product
- The first symbol can be picked in  $n$  ways
- How many choices there are for the second symbol?
- We can place anything there, except for the symbol on the first place
- Symbol on the first place might be arbitrary, but whatever it is there are  $n-1$  choices for the second symbol!



# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & & & \end{array}$$

- So we can pick the first and the second symbol in  $n \times (n-1)$  ways

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & & \end{array}$$

- So we can pick the first and the second symbol in  $n \times (n-1)$  ways
- Now, the third symbol can be picked among  $n-2$  options: all except the symbols in the first and the second position

# Permutations

$$\begin{array}{cccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & \end{array}$$

- So we can pick the first and the second symbol in  $n \times (n-1)$  ways
- Now, the third symbol can be picked among  $n-2$  options: all except the symbols in the first and the second position
- And so on; for each next symbol we have one less option

# Permutations

$$\begin{array}{ccccccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & n-k+1 \end{array}$$

- So we can pick the first and the second symbol in  $n \times (n-1)$  ways
- Now, the third symbol can be picked among  $n-2$  options: all except the symbols in the first and the second position
- And so on; for each next symbol we have one less option
- In the end for the last object we have  $n-k+1$  options

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & n-k+1 \end{array}$$

- Overall we have  $n \times (n-1) \times \dots \times (n-k+1)$   
 $k$ -permutations

# Permutations

$$\begin{array}{ccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & n-k+1 \end{array}$$

- Overall we have  $n \times (n-1) \times \dots \times (n-k+1)$   
 $k$ -permutations
- Convenient notation:  $n! = 1 \times 2 \times \dots \times n$ ; this number is called **factorial** of  $n$

# Permutations

$$\begin{array}{cccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & n-k+1 \end{array}$$

- Overall we have  $n \times (n-1) \times \dots \times (n-k+1)$   $k$ -permutations
- Convenient notation:  $n! = 1 \times 2 \times \dots \times n$ ; this number is called **factorial** of  $n$
- In this notation the number of  $k$ -permutations of  $n$  symbols of length  $k$  looks nicer: it is  $n! / (n-k)!$

# Permutations

$$\begin{array}{cccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & n-k+1 \end{array}$$

- Overall we have  $n \times (n-1) \times \dots \times (n-k+1)$   $k$ -permutations
- Convenient notation:  $n! = 1 \times 2 \times \dots \times n$ ; this number is called **factorial** of  $n$
- In this notation the number of  $k$ -permutations of  $n$  symbols of length  $k$  looks nicer: it is  $n! / (n-k)!$
- What if  $n-k=0$ ?



# Permutations

$$\begin{array}{cccccc} 1 & 2 & 3 & \dots & k \\ * & * & * & \dots & * \\ n & n-1 & n-2 & \dots & n-k+1 \end{array}$$

- Overall we have  $n \times (n-1) \times \dots \times (n-k+1)$   $k$ -permutations
- Convenient notation:  $n! = 1 \times 2 \times \dots \times n$ ; this number is called **factorial** of  $n$
- In this notation the number of  $k$ -permutations of  $n$  symbols of length  $k$  looks nicer: it is  $n! / (n-k)!$
- What if  $n-k=0$ ? Convention:  $0! = 1$

# Permutations

## Problem

In how many orders can we place  $n$  books on the shelf?

# Permutations

## Problem

In how many orders can we place  $n$  books on the shelf?

- Each book is a symbol

# Permutations

## Problem

In how many orders can we place  $n$  books on the shelf?

- Each book is a symbol
- We need to count  $n$ -permutations of  $n$  symbols; these are called **permutations**

# Permutations

## Problem

In how many orders can we place  $n$  books on the shelf?

- Each book is a symbol
- We need to count  $n$ -permutations of  $n$  symbols; these are called **permutations**
- By the previous result there are  $n!$  of them

# Permutations

## Problem

In how many orders can we place  $n$  books on the shelf?

- Each book is a symbol
- We need to count  $n$ -permutations of  $n$  symbols; these are called **permutations**
- By the previous result there are  $n!$  of them
- This is the formula that was used in the discussion of magic square in course “What is a Proof?”

# Conclusion

- We have started the discussion of **Combinatorics**

# Conclusion

- We have started the discussion of **Combinatorics**
- Important for Probability Theory, estimations on running time of algorithms, mathematics in general



# Conclusion

- We have started the discussion of **Combinatorics**
- Important for Probability Theory, estimations on running time of algorithms, mathematics in general
- We have discussed two standard settings: **tuples** and **permutations**; they help in many cases

# Conclusion

- We have started the discussion of **Combinatorics**
- Important for Probability Theory, estimations on running time of algorithms, mathematics in general
- We have discussed two standard settings: **tuples** and **permutations**; they help in many cases
- **Recursive counting** is also useful, especially for computational applications

# Conclusion

- We have started the discussion of **Combinatorics**
- Important for Probability Theory, estimations on running time of algorithms, mathematics in general
- We have discussed two standard settings: **tuples** and **permutations**; they help in many cases
- **Recursive counting** is also useful, especially for computational applications
- Still are not ready to count, say, what are the chances to get two aces in 6 card hand

# Conclusion

- We have started the discussion of **Combinatorics**
- Important for Probability Theory, estimations on running time of algorithms, mathematics in general
- We have discussed two standard settings: **tuples** and **permutations**; they help in many cases
- **Recursive counting** is also useful, especially for computational applications
- Still are not ready to count, say, what are the chances to get two aces in 6 card hand
- More to come next week!