

# Pascal's Triangle

---

Alexander S. Kulikov

Steklov Mathematical Institute at St. Petersburg, Russian Academy of Sciences  
and  
University of California, San Diego

# Outline

Pascal's Triangle

Symmetries

Row Sums

Binomial Theorem

# Combinations

## Question

There are  $n$  students. What is the number of ways of forming a team of  $k$  students out of them?

# Combinations

## Question

There are  $n$  students. What is the number of ways of forming a team of  $k$  students out of them?

## Answer

$$\binom{n}{k}$$

# Forming a Team

- Fix one of the students, call her Alice

# Forming a Team

- Fix one of the students, call her Alice
- There are two types of teams:

# Forming a Team

- Fix one of the students, call her Alice
- There are two types of teams:
  1. Teams with Alice:  $\binom{n-1}{k-1}$

# Forming a Team

- Fix one of the students, call her Alice
- There are two types of teams:
  1. Teams with Alice:  $\binom{n-1}{k-1}$
  2. Teams without Alice:  $\binom{n-1}{k}$



# Forming a Team

- Fix one of the students, call her Alice
- There are two types of teams:
  1. Teams with Alice:  $\binom{n-1}{k-1}$
  2. Teams without Alice:  $\binom{n-1}{k}$
- Hence,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

# Pascal's Triangle


$$n = 0$$

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

# Pascal's Triangle

$$n = 0$$

$$n = 1$$




# Pascal's Triangle

$$n = 0$$

$$n = 1$$

$$n = 2$$




# Pascal's Triangle

$$n = 0$$

$$n = 1$$

$$n = 2$$

$$n = 3$$



# Pascal's Triangle


$$n = 0$$

$$n = 1$$

$$n = 2$$

$$n = 3$$

$$n = 4$$



# Pascal's Triangle

$$n = 0$$


$$n = 1$$

$$n = 2$$

$$n = 3$$

$$n = 4$$

$$n = 5$$



# Pascal's Triangle

$$n = 0$$


$$n = 1$$

$$n = 2$$

$$n = 3$$

$$n = 4$$

$$n = 5$$



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$



# Pascal's Triangle

$$n = 0$$


$$n = 1$$

$$n = 2$$

$$n = 3$$

$$n = 4$$

$$n = 5$$



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

# Pascal's Triangle

$$n = 0$$


$$n = 1$$

$$n = 2$$

$$n = 3$$

$$n = 4$$

$$n = 5$$



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

# Pascal's Triangle

$n = 0$


$n = 1$

$n = 2$

$n = 3$

$n = 4$

$n = 5$



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

# Code

```
C = dict() #  $C([n, k])$  is equal to  $n$  choose  $k$ 
```

```
for n in range(8):
```

```
    C[n, 0] = 1
```

```
    C[n, n] = 1
```

```
    for k in range(1, n):
```

```
        C[n, k] = C[n - 1, k - 1] + C[n - 1, k]
```

```
print(C[7, 4])
```

# Code

```
C = dict() # C([n,k]) is equal to n choose k

for n in range(8):
    C[n, 0] = 1
    C[n, n] = 1

    for k in range(1, n):
        C[n, k] = C[n - 1, k - 1] + C[n - 1, k]

print(C[7, 4])
```

# Outline

Pascal's Triangle

**Symmetries**

Row Sums


Binomial Theorem







# Pascal's Triangle is Symmetric



$$\binom{n}{k} = \binom{n}{n-k}$$

## Theorem

$$\binom{n}{k} = \binom{n}{n-k}$$

## Theorem

$$\binom{n}{k} = \binom{n}{n-k}$$

## Proof

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n!}{(n-k)!k!} = \binom{n}{n-k}$$

# Combinatorial Proof

- $\binom{n}{k}$  is the number of ways of selecting a team of size  $k$  out of  $n$  students

# Combinatorial Proof

- $\binom{n}{k}$  is the number of ways of selecting a team of size  $k$  out of  $n$  students
- $\binom{n}{n-k}$  is the number of ways of selecting a team of size  $n - k$  out of  $n$  students

# Combinatorial Proof

- $\binom{n}{k}$  is the number of ways of selecting a team of size  $k$  out of  $n$  students
- $\binom{n}{n-k}$  is the number of ways of selecting a team of size  $n - k$  out of  $n$  students
- this is just the number of ways of partitioning  $n$  students into two teams of size  $k$  and  $n - k$

# Outline


Pascal's Triangle

Symmetries

**Row Sums**

Binomial Theorem

## Row Sums





## Row Sums

$$\begin{array}{cccccccccc}
 & & & & 1 & & & & & \\
 & & & & 1 & + & 1 & & & \\
 & & & 1 & + & 2 & + & 1 & & \\
 & & 1 & + & 3 & + & 3 & + & 1 & \\
 & 1 & + & 4 & + & 6 & + & 4 & + & 1 \\
 1 & + & 5 & + & 10 & + & 10 & + & 5 & + & 1
 \end{array}$$

# Row Sums

$$\begin{array}{cccccccccccl} & & & & 1 & & & & & & = 1 \\ & & & 1 & + & 1 & & & & & = 2 \\ & & 1 & + & 2 & + & 1 & & & & = 4 \\ & 1 & + & 3 & + & 3 & + & 1 & & & = 8 \\ 1 & + & 4 & + & 6 & + & 4 & + & 1 & & = 16 \\ 1 & + & 5 & + & 10 & + & 10 & + & 5 & + & 1 & = 32 \end{array}$$

## Theorem

The sum of all the numbers in the  $n$ -th row of Pascal's triangle is equal to  $2^n$ :


$$\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n-1} + \binom{n}{n} = 2^n$$

# Proof by Induction

- The base case (0-th row) holds

# Proof by Induction

- The base case (0-th row) holds
- We'll show that the sum of each row is twice the sum of the previous row:




$$\begin{array}{ccccccccc} 1 & & 4 & & 6 & & 4 & & 1 \\ 1 & (1+3) & (3+3) & (3+1) & 1 \\ (1+1) & (3+3) & (3+3) & (1+1) \end{array}$$

# Combinatorial Proof

- $\binom{n}{k}$  is the number of  $k$ -subsets of a set of size  $n$
- the sum of  $\binom{n}{k}$  for all  $k$  (from 0 to  $n$ ) is the number of all subsets of an  $n$  element set
- this is  $2^n$  by the product rule: each of the  $n$  elements is either included or not

# Alternating Row Sums



# Alternating Row Sums

$$\begin{array}{cccccccccc} & & & & 1 & & & & & & \\ & & & 1 & - & 1 & & & & & \\ & & 1 & - & 2 & + & 1 & & & & \\ & 1 & - & 3 & + & 3 & - & 1 & & & \\ 1 & - & 4 & + & 6 & - & 4 & + & 1 & & \\ 1 & - & 5 & + & 10 & - & 10 & + & 5 & - & 1 \end{array}$$



## Alternating Row Sums

$$\begin{array}{cccccccccccl}
 & & & & & & & & & & 1 & & & & & & & & & & \\
 & & & & & & & & & & 1 & - & 1 & & & & & & & & = 0 \\
 & & & & & & & & & & 1 & - & 2 & + & 1 & & & & & & & = 0 \\
 & & & & & & & & & & 1 & - & 3 & + & 3 & - & 1 & & & & & & = 0 \\
 & & & & & & & & & & 1 & - & 4 & + & 6 & - & 4 & + & 1 & & & & = 0 \\
 & & & & & & & & & & 1 & - & 5 & + & 10 & - & 10 & + & 5 & - & 1 & & = 0
 \end{array}$$

## Theorem

$$\text{For } n > 0, \quad \sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

## Theorem

$$\text{For } n > 0, \quad \sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

- For odd  $n$ , follows immediately from the symmetry property

## Theorem

$$\text{For } n > 0, \quad \sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

- For odd  $n$ , follows immediately from the symmetry property
- In general, can be shown by using the sum pattern of the triangle (each internal element is equal to the sum of the two elements above it)

# Combinatorial Proof

- Need to show that

$$\binom{n}{1} + \binom{n}{3} + \cdots = \binom{n}{0} + \binom{n}{2} + \cdots$$

# Combinatorial Proof

- Need to show that

$$\binom{n}{1} + \binom{n}{3} + \cdots = \binom{n}{0} + \binom{n}{2} + \cdots$$

- Combinatorial meaning: the number of odd size subsets is the same as the number of even size subsets

# Combinatorial Proof

- Need to show that

$$\binom{n}{1} + \binom{n}{3} + \dots = \binom{n}{0} + \binom{n}{2} + \dots$$

- Combinatorial meaning: the number of odd size subsets is the same as the number of even size subsets
- To prove this, we'll construct a **one-to-one correspondence** between odd size subsets and even size subsets

# One-to-one Correspondence

- Fix any element  $x$  (can do this since  $n > 0$ )



# One-to-one Correspondence

- Fix any element  $x$  (can do this since  $n > 0$ )
- Partition all subsets into pairs  $(A, B)$  where  $A = B + x$  (more formally,  $A = B \cup \{x\}$ )

# One-to-one Correspondence

- Fix any element  $x$  (can do this since  $n > 0$ )
- Partition all subsets into pairs  $(A, B)$  where  $A = B + x$  (more formally,  $A = B \cup \{x\}$ )
- One of  $A, B$  has odd size, the other one has even size

# Example

$$S = \{a, b, c, d\}$$

## Even size subsets

$\emptyset$

$\{a, b\}$

$\{a, c\}$

$\{a, d\}$

$\{b, c\}$

$\{b, d\}$

$\{c, d\}$

$\{a, b, c, d\}$

## Odd size subsets

$\{a\}$

$\{b\}$

$\{c\}$

$\{d\}$

$\{a, b, c\}$

$\{a, b, d\}$

$\{a, c, d\}$

$\{b, c, d\}$


# Outline

Pascal's Triangle

Symmetries

Row Sums

**Binomial Theorem**



$$(a + b)^2 = a^2 + 2ab + b^2$$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

$$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$$

# Binomial Theorem

$$(a+b)^n = \binom{n}{0} a^n + \binom{n}{1} a^{n-1} b + \dots + \binom{n}{k} a^{n-k} b^k + \dots + \binom{n}{n} b^n$$

# Binomial Theorem

$$(a+b)^n = \binom{n}{0} a^n + \binom{n}{1} a^{n-1} b + \dots + \binom{n}{k} a^{n-k} b^k + \dots + \binom{n}{n} b^n$$

Equivalently,

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

# Binomial Theorem

$$(a+b)^n = \binom{n}{0} a^n + \binom{n}{1} a^{n-1} b + \dots + \binom{n}{k} a^{n-k} b^k + \dots + \binom{n}{n} b^n$$

Equivalently,

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

Can be shown by expanding the expression

$$(a+b)(a+b)\dots(a+b)$$



# Proof by Induction

$$\begin{aligned}(a + b)^4 &= (a + b)^3(a + b) \\&= (a^3 + 3a^2b + 3ab^2 + b^3)(a + b) \\&= a^4 + 3a^3b + 3a^2b^2 + ab^3 + \\&\quad + a^3b + 3a^2b + 3ab^3 + b^4 \\&= a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4\end{aligned}$$

# Example

$$\begin{aligned}(2a - b)^4 &= ((2a) + (-b))^4 \\&= (2a)^4 + 4(2a)^3(-b) + 6(2a)^2(-b)^2 + 4(2a)(-b)^3 + (-b)^4 \\&= 16a^4 - 32a^3b + 24a^2b^2 - 8ab^3 + b^4\end{aligned}$$

# Consequences

- Set  $a = b = 1$ . The number of subsets is  $2^n$ :

$$2^n = \sum_{k=0}^n \binom{n}{k}$$

# Consequences

- Set  $a = b = 1$ . The number of subsets is  $2^n$ :

$$2^n = \sum_{k=0}^n \binom{n}{k}$$

- Set  $a = 1, b = -1$ . The number of odd size subsets is the same as the number of even size subsets:

$$0 = \sum_{k=0}^n (-1)^k \binom{n}{k}$$

## Another Consequence

- Set  $a = 1, b = 2$ :

$$3^n = \binom{n}{0} + \binom{n}{1}2 + \binom{n}{2}2^2 + \cdots + \binom{n}{n}2^n$$

## Another Consequence

- Set  $a = 1, b = 2$ :

$$3^n = \binom{n}{0} + \binom{n}{1}2 + \binom{n}{2}2^2 + \cdots + \binom{n}{n}2^n$$

- Combinatorial proof:

## Another Consequence

- Set  $a = 1, b = 2$ :

$$3^n = \binom{n}{0} + \binom{n}{1}2 + \binom{n}{2}2^2 + \cdots + \binom{n}{n}2^n$$

- Combinatorial proof:
  - $3^n$  is the number of words of length  $n$  over the alphabet  $\{x, y, z\}$

## Another Consequence

- Set  $a = 1$ ,  $b = 2$ :

$$3^n = \binom{n}{0} + \binom{n}{1}2 + \binom{n}{2}2^2 + \cdots + \binom{n}{n}2^n$$

- Combinatorial proof:
  - $3^n$  is the number of words of length  $n$  over the alphabet  $\{x, y, z\}$
  - $\binom{n}{0}$  is the number of words consisting of the letter  $x$  only



## Another Consequence

- Set  $a = 1, b = 2$ :

$$3^n = \binom{n}{0} + \binom{n}{1}2 + \binom{n}{2}2^2 + \cdots + \binom{n}{n}2^n$$

- Combinatorial proof:
  - $3^n$  is the number of words of length  $n$  over the alphabet  $\{x, y, z\}$
  - $\binom{n}{0}$  is the number of words consisting of the letter  $x$  only
  - $\binom{n}{1}2$  is the number of words with exactly  $n - 1$  letters  $x$

## Another Consequence

- Set  $a = 1, b = 2$ :

$$3^n = \binom{n}{0} + \binom{n}{1}2 + \binom{n}{2}2^2 + \cdots + \binom{n}{n}2^n$$

- Combinatorial proof:
  - $3^n$  is the number of words of length  $n$  over the alphabet  $\{x, y, z\}$
  - $\binom{n}{0}$  is the number of words consisting of the letter  $x$  only
  - $\binom{n}{1}2$  is the number of words with exactly  $n - 1$  letters  $x$
  - $\binom{n}{2}2^2$  is the number of words with exactly  $n - 2$  letters  $x$