

PROPERTIES OF HYPERHOLOMORPHIC FUNCTIONS ON DUAL SEDENION NUMBERS

JI EUN KIM, SU JIN HA, AND KWANG HO SHON*

Abstract. The aim of this paper is to define hyperholomorphic functions with dual sedenion variables on $\mathcal{S} \times \mathcal{S}$, where $\mathcal{S} \cong \mathbb{C}^8$. By the condition of harmonicity, we research properties of hyperholomorphic functions of dual sedenion variables in Clifford analysis.

1. Introduction

The dual numbers extended the real numbers by adjoining one new element ε with the property $\varepsilon^2 = 0$. Every dual number has the form $z = x + \varepsilon y$ with x and y uniquely determined real numbers. The collection of dual numbers forms a particular two-dimensional commutative unital associative algebra over the real numbers. Hörmander [1] and Krantz [6] introduced the concepts of several complex variables and theories to complex analysis in several variables. K. Nôno [8, 9] researched the properties of hyperholomorphic functions of quaternion variable and octonion variable in the 1980s. We [2, 7, 10] found the theorem about hyperholomorphic functions of dual quaternion in $\mathbb{C}^2 \times \mathbb{C}^2$ and investigated the extensions of quaternions. Also, we [3, 4, 5] researched properties of the regularity functions on a variety of quaternionic variables and the corresponding Cauchy-Riemann system on each several types.

2. Preliminaries

The multiplication of these unit sedenions follows:

Received November 19, 2014. Accepted November 27, 2014.

2010 Mathematics Subject Classification. 32A99, 30G35, 11E88.

Key words and phrases. Clifford Analysis, hyperholomorphic function, sedenion variables, dual number system, the condition of harmonicity.

*Corresponding author

\times	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8	e_9	e_{10}	e_{11}	e_{12}	e_{13}	e_{14}	e_{15}
e_0	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8	e_9	e_{10}	e_{11}	e_{12}	e_{13}	e_{14}	e_{15}
e_1	e_1	-1	e_3	$-e_2$	e_5	$-e_4$	e_7	$-e_6$	e_9	$-e_8$	e_{11}	$-e_{10}$	e_{13}	$-e_{12}$	e_{15}	$-e_{14}$
e_2	e_2	$-e_3$	-1	e_1	e_6	$-e_7$	$-e_4$	e_5	e_{10}	$-e_{11}$	$-e_8$	e_9	e_{14}	$-e_{15}$	e_{13}	e_{13}
e_3	e_3	e_2	$-e_1$	-1	e_7	e_6	$-e_5$	$-e_4$	e_{11}	e_{10}	$-e_9$	$-e_8$	e_{15}	e_{14}	$-e_{13}$	$-e_{12}$
e_4	e_4	$-e_5$	$-e_6$	$-e_7$	-1	e_1	e_2	e_3	e_{12}	e_{13}	e_{14}	e_{15}	$-e_8$	$-e_9$	$-e_{10}$	$-e_{11}$
e_5	e_5	e_4	e_7	$-e_6$	$-e_1$	-1	e_3	$-e_2$	e_{13}	$-e_{12}$	e_{15}	$-e_{14}$	e_9	$-e_8$	e_{11}	$-e_{10}$
e_6	e_6	$-e_7$	e_4	e_5	$-e_2$	$-e_3$	-1	e_1	e_{14}	$-e_{15}$	$-e_{12}$	e_{13}	e_{10}	$-e_{11}$	$-e_8$	e_9
e_7	e_7	e_6	$-e_5$	e_4	$-e_3$	e_2	$-e_1$	-1	e_{15}	e_{14}	$-e_{13}$	$-e_{12}$	e_{11}	e_{10}	$-e_9$	$-e_8$
e_8	e_8	$-e_9$	$-e_{10}$	$-e_{11}$	$-e_{12}$	$-e_{13}$	$-e_{14}$	$-e_{15}$	-1	e_1	e_2	e_3	e_4	e_5	e_6	e_7
e_9	e_9	e_8	e_{11}	$-e_{10}$	$-e_{13}$	e_{12}	e_{15}	$-e_{14}$	$-e_1$	-1	e_3	$-e_2$	$-e_5$	e_4	e_7	$-e_6$
e_{10}	e_{10}	$-e_{11}$	e_8	e_9	$-e_{14}$	$-e_{15}$	e_{12}	e_{13}	$-e_2$	$-e_3$	-1	e_1	$-e_6$	$-e_7$	e_4	e_5
e_{11}	e_{11}	e_{10}	$-e_9$	e_8	$-e_{15}$	e_{14}	$-e_{13}$	e_{12}	$-e_3$	e_2	$-e_1$	-1	$-e_7$	e_6	$-e_5$	e_4
e_{12}	e_{12}	$-e_{13}$	$-e_{14}$	$-e_{15}$	e_8	$-e_9$	$-e_{10}$	$-e_{11}$	$-e_4$	e_5	e_6	e_7	-1	e_1	e_2	e_3
e_{13}	e_{13}	e_{12}	e_{15}	$-e_{14}$	e_9	e_8	e_{11}	$-e_{10}$	$-e_5$	$-e_4$	e_7	$-e_6$	$-e_1$	-1	e_3	$-e_2$
e_{14}	e_{14}	$-e_{15}$	e_{12}	e_{13}	e_{10}	$-e_{11}$	e_8	e_9	$-e_6$	$-e_7$	$-e_4$	e_5	$-e_2$	$-e_3$	-1	e_1
e_{15}	e_{15}	e_{14}	$-e_{13}$	e_{12}	e_{11}	e_{10}	$-e_9$	e_8	$-e_7$	e_6	$-e_5$	$-e_4$	$-e_3$	$-e_2$	$-e_1$	-1

The algebra of dual numbers is a ring that is a local ring since the principal ideal generated by ε is its only maximal ideal. Dual numbers form the coefficients of dual quaternions. Using matrices, dual numbers can be represented as

$$\varepsilon = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad a + b\varepsilon = \begin{pmatrix} a & b \\ 0 & a \end{pmatrix}.$$

The sum and product of dual numbers are then calculated with ordinary matrix addition and matrix multiplication; both operations are commutative and associative within the algebra of dual numbers.

For the field \mathcal{S} of sedenions, we give hyperholomorphic functions and a harmonic function with dual sedenion variables on $\mathcal{S} \times \mathcal{S}$ and research properties of functions of dual sedenion variables.

The element e_0 is the identity of \mathcal{S} and e_1 identifies the imaginary unit $\sqrt{-1}$ in the \mathbb{C} -field of complex numbers. A sedenion z is given by

$$(2.1) \quad z = x_0 + \sum_{i=0}^{15} e_i x_i, \quad (x_i \in \mathbb{R}, \quad i = 0, 1, \dots, 15)$$

is an sixteen dimensional non-commutative and non-associative \mathbb{R} -field generated by sixteen base elements $e_l (l = 0, 1, \dots, 15)$ with the following non-commutative multiplication rules:

$$e_i^2 = -1, \quad e_i e_j = -e_j e_i, \quad e_i e_j e_k = e_i (e_j e_k) \quad (i \neq j \neq k, i \neq 0, j \neq 0, k \neq 0).$$

An sedenion z given by (2.1) is regarded as $z = z_1 + z_2 e_2 + z_3 e_4 + z_4 e_6 + z_5 e_8 + z_6 e_{10} + z_7 e_{12} + z_8 e_{14} \in \mathcal{S}$, where $z_1 := x_0 + e_1 x_1$, $z_2 := x_2 + e_1 x_3$, $z_3 := x_4 + e_1 x_5$, $z_4 := x_6 + e_1 x_7$, $z_5 := x_8 + e_1 x_9$, $z_6 := x_{10} + e_1 x_{11}$, $z_7 := x_{12} + e_1 x_{13}$ and $z_8 := x_{14} + e_1 x_{15}$ are complex numbers in \mathcal{S} . Thus, we identify \mathcal{S} with \mathbb{C}^8 .

We write the sedenion $z = \sum_{l=0}^{15} e_l x_l$ and the sedenion conjugate $z^* = x_0 - \sum_{l=1}^{15} e_l x_l$. Also, the absolute value $|z|$ of z and an inverse z^{-1} of z in \mathcal{S} are defined by

$$|z| = \sqrt{\sum_{l=1}^8 |z_l|^2}, \quad z^{-1} = \frac{z^*}{|z|^2} \quad (z \neq 0).$$

Thus, the sedenion $z \in \mathcal{S}$ have the following forms:

$$\begin{aligned} z &= \sum_{l=0}^{15} e_l x_l = z_1 + z_2 e_2 + z_3 e_4 + z_4 e_6 + z_5 e_8 + z_6 e_{10} + z_7 e_{12} + z_8 e_{14} \\ &= Z_1 + Z_2 e_4 + Z_3 e_8 + Z_4 e_{12} = P_1 + P_2 e_8 \in \mathcal{S} \end{aligned}$$

and

$$\begin{aligned}
 z^* &= x_0 - \sum_{l=1}^{15} e_l x_l \\
 &= \bar{z}_1 - z_2 e_2 - z_3 e_4 - z_4 e_6 - z_5 e_8 - z_6 e_{10} - z_7 e_{12} - z_8 e_{14} \\
 &= Z_1^* - Z_2 e_4 - Z_3 e_8 - Z_4 e_{12} = P_1^* - P_2 e_8 \in \mathcal{S},
 \end{aligned}$$

where $Z_1 = z_1 + z_2 e_2$, $Z_2 = z_3 + z_4 e_2$, $Z_3 = z_5 + z_6 e_2$ and $Z_4 = z_7 + z_8 e_2$ are quaternion numbers in \mathbb{C}^2 , $P_1 = Z_1 + Z_2 e_4$ and $P_2 = Z_3 + Z_4 e_4$ are octonion numbers in \mathbb{C}^4 .

We use the following differential operators:

$$\begin{aligned}
 \frac{\partial}{\partial Z_1} &:= \frac{\partial}{\partial z_1} - e_2 \frac{\partial}{\partial z_2}, & \frac{\partial}{\partial Z_1^*} &= \frac{\partial}{\partial \bar{z}_1} + e_2 \frac{\partial}{\partial z_2}, \\
 \frac{\partial}{\partial Z_2} &:= \frac{\partial}{\partial z_3} - e_2 \frac{\partial}{\partial z_4}, & \frac{\partial}{\partial Z_2^*} &= \frac{\partial}{\partial \bar{z}_3} + e_2 \frac{\partial}{\partial z_4}, \\
 \frac{\partial}{\partial Z_3} &:= \frac{\partial}{\partial z_5} - e_2 \frac{\partial}{\partial z_6}, & \frac{\partial}{\partial Z_3^*} &= \frac{\partial}{\partial \bar{z}_5} + e_2 \frac{\partial}{\partial z_6}, \\
 \frac{\partial}{\partial Z_4} &:= \frac{\partial}{\partial z_7} - e_2 \frac{\partial}{\partial z_8}, & \frac{\partial}{\partial Z_4^*} &= \frac{\partial}{\partial \bar{z}_7} + e_2 \frac{\partial}{\partial z_8},
 \end{aligned}$$

where $\partial/\partial z_l$, $\partial/\partial \bar{z}_l$ ($l = 1, 2, \dots, 8$) are usual differential operators used in complex analysis. And, we use the following differential operators:

$$\begin{aligned}
 \frac{\partial}{\partial P_1} &:= \frac{\partial}{\partial Z_1} - e_4 \frac{\partial}{\partial Z_2}, & \frac{\partial}{\partial P_1^*} &= \frac{\partial}{\partial Z_1^*} + e_4 \frac{\partial}{\partial Z_2}, \\
 \frac{\partial}{\partial P_2} &:= \frac{\partial}{\partial Z_3} - e_4 \frac{\partial}{\partial Z_4}, & \frac{\partial}{\partial P_2^*} &= \frac{\partial}{\partial Z_3^*} + e_4 \frac{\partial}{\partial Z_4}.
 \end{aligned}$$

Also, we use the following sedenion differential operators:

$$D := \frac{\partial}{\partial P_1} - e_8 \frac{\partial}{\partial P_2}, \quad D^* = \frac{\partial}{\partial P_1^*} + e_8 \frac{\partial}{\partial P_2^*}.$$

The operator

$$DD^* = \sum_{l=1}^8 \frac{\partial^2}{\partial z_l \partial \bar{z}_l} = \frac{1}{4} \sum_{l=0}^{15} \frac{\partial^2}{\partial x_l^2}$$

is the usual complex Laplacian Δ .

3. Regular function with sedenion variables

Let $\Omega = \Omega_1 \times \Omega_2$ be an open set in $\mathcal{S} \times \mathcal{S}$. The function $f(z)$ is defined by the following form in Ω with valued in \mathcal{S} :

$$f : \Omega \rightarrow \mathcal{S}, (z, w) \in \Omega \mapsto f(z, w) = \phi_1(z, w) + \phi_2(z, w)e_8.$$

In detail,

$$f(z, w) = \sum_{l=0}^{15} u_l e_l = \sum_{m=1}^8 f_m e_{2m-1} = \sum_{n=0}^3 F_n e_{4n} = \phi_1 + \phi_2 e_8 \in \mathcal{S},$$

where u_l ($l = 0, 1, \dots, 15$) are real-valued functions, $f_m = u_{2m-2} + u_{2m-1}e_1$ ($m = 1, 2, \dots, 8$) are complex-valued functions, $F_n = f_{2n-1} + f_{2n}e_2$ ($n = 1, 2, 3, 4$) are quaternion-valued functions, $\phi_1 = F_1 + F_2e_4$ and $\phi_2 = F_3 + F_4e_4$ are octonion-valued functions in \mathcal{S} .

We use the following differential operators:

$$\begin{aligned} \frac{\partial}{\partial Z_1} &:= \frac{\partial}{\partial z_1} - e_2 \frac{\partial}{\partial z_2}, & \frac{\partial}{\partial Z_1^*} &= \frac{\partial}{\partial \bar{z}_1} + e_2 \frac{\partial}{\partial \bar{z}_2}, \\ \frac{\partial}{\partial Z_2} &:= \frac{\partial}{\partial z_3} - e_2 \frac{\partial}{\partial z_4}, & \frac{\partial}{\partial Z_2^*} &= \frac{\partial}{\partial \bar{z}_3} + e_2 \frac{\partial}{\partial \bar{z}_4}, \\ \frac{\partial}{\partial Z_3} &:= \frac{\partial}{\partial z_5} - e_2 \frac{\partial}{\partial z_6}, & \frac{\partial}{\partial Z_3^*} &= \frac{\partial}{\partial \bar{z}_5} + e_2 \frac{\partial}{\partial \bar{z}_6}, \\ \frac{\partial}{\partial Z_4} &:= \frac{\partial}{\partial z_7} - e_2 \frac{\partial}{\partial z_8}, & \frac{\partial}{\partial Z_4^*} &= \frac{\partial}{\partial \bar{z}_7} + e_2 \frac{\partial}{\partial \bar{z}_8}, \\ \frac{\partial}{\partial W_1} &:= \frac{\partial}{\partial w_1} - e_2 \frac{\partial}{\partial w_2}, & \frac{\partial}{\partial W_1^*} &= \frac{\partial}{\partial \bar{w}_1} + e_2 \frac{\partial}{\partial \bar{w}_2}, \\ \frac{\partial}{\partial W_2} &:= \frac{\partial}{\partial w_3} - e_2 \frac{\partial}{\partial w_4}, & \frac{\partial}{\partial W_2^*} &= \frac{\partial}{\partial \bar{w}_3} + e_2 \frac{\partial}{\partial \bar{w}_4}, \\ \frac{\partial}{\partial W_3} &:= \frac{\partial}{\partial w_5} - e_2 \frac{\partial}{\partial w_6}, & \frac{\partial}{\partial W_3^*} &= \frac{\partial}{\partial \bar{w}_5} + e_2 \frac{\partial}{\partial \bar{w}_6}, \\ \frac{\partial}{\partial W_4} &:= \frac{\partial}{\partial w_7} - e_2 \frac{\partial}{\partial w_8}, & \frac{\partial}{\partial Z_4^*} &= \frac{\partial}{\partial \bar{w}_7} + e_2 \frac{\partial}{\partial \bar{w}_8}, \end{aligned}$$

where $\partial/\partial z_l, \partial/\partial \bar{z}_l$ ($l = 1, 2, \dots, 8$) are usual differential operators used in complex analysis. And, we use the following differential operators:

$$\begin{aligned}
\frac{\partial}{\partial P_1} &:= \frac{\partial}{\partial Z_1} - e_4 \frac{\partial}{\partial Z_2}, & \frac{\partial}{\partial P_1^*} &= \frac{\partial}{\partial Z_1^*} + e_4 \frac{\partial}{\partial Z_2}, \\
\frac{\partial}{\partial P_2} &:= \frac{\partial}{\partial Z_3} - e_4 \frac{\partial}{\partial Z_4}, & \frac{\partial}{\partial P_2^*} &= \frac{\partial}{\partial Z_3^*} + e_4 \frac{\partial}{\partial Z_4}, \\
\frac{\partial}{\partial Q_1} &:= \frac{\partial}{\partial W_1} - e_4 \frac{\partial}{\partial W_2}, & \frac{\partial}{\partial Q_1^*} &= \frac{\partial}{\partial W_1^*} + e_4 \frac{\partial}{\partial W_2}, \\
\frac{\partial}{\partial Q_2} &:= \frac{\partial}{\partial W_3} - e_4 \frac{\partial}{\partial W_4}, & \frac{\partial}{\partial Q_2^*} &= \frac{\partial}{\partial W_3^*} + e_4 \frac{\partial}{\partial W_4}.
\end{aligned}$$

Also, we use the following sedenion differential operators:

$$\begin{aligned}
D &:= \frac{\partial}{\partial P_1} - e_8 \frac{\partial}{\partial P_2}, & D^* &= \frac{\partial}{\partial P_1^*} + e_8 \frac{\partial}{\partial P_2}, \\
E &:= \frac{\partial}{\partial Q_1} - e_8 \frac{\partial}{\partial Q_2}, & E^* &= \frac{\partial}{\partial Q_1^*} + e_8 \frac{\partial}{\partial Q_2}, \\
H &:= D + \varepsilon E, & H^* &:= D^* + \varepsilon E^*.
\end{aligned}$$

Definition 3.1. Let Ω be an open set in $\mathcal{S} \times \mathcal{S}$. A function $f(z, w) = \phi_1(z, w) + \phi_2(z, w)e_8$ is said to be bihyperholomorphic in Ω if the following two conditions are satisfied:

- (1) $\phi_r(z, w)$ ($r = 1, 2$) are continuously differential functions in Ω .
- (2) The function f satisfies

$$(3.1) \quad D^* f = 0, \quad f E^* = 0$$

in Ω .

Theorem 3.1. Let G and T be functions with values in sedenions of class \mathcal{C}^∞ on Ω in $\mathcal{S} \times \mathcal{S}$. Then the Cousin 1 problem according to sedenion-valued functions on Ω has a solution.

Proof. Consider the inhomogeneous Cauchy-Riemann system of partial differential equations

$$(3.2) \quad D^* f = G, \quad f E^* = T.$$

If the system (3.1) has a solution f with values in sedenions, then

$$GE^* = D^* f E^* = D^* T.$$

For the system (3.2), the solvability condition

$$(3.3) \quad GE^* = D^* T$$

is necessary for the existence of a solution f to the system (3.1). Let Ω be a product domain $\Omega_1 \times \Omega_2$ of simply connected domains in \mathcal{S} and \mathcal{S} , respectively, and let

$$Rf := (D^*f, fE^*).$$

Then $Rf = (G, T) = 0$ satisfies (3.3). For $Rf : \Omega \rightarrow \mathcal{S}$ of class \mathcal{C}^2 , if

$$A(G, T) = GE^* - D^*T,$$

then the Cousin 1 problem according to sedenion-valued functions on Ω has a solution. \square

Let $\Omega = \Omega_1 \times \Omega_2$ be an open set in $\mathcal{S} \times \mathcal{S}$. The function $F(\zeta)$ is defined by the following form in Ω with valued in $\mathcal{S} \times \mathcal{S}$:

$$\begin{aligned} F : \Omega &\rightarrow \mathcal{S} \times \mathcal{S}, \\ \zeta = z + \varepsilon w \in \Omega &\mapsto F(\zeta) = F(z + \varepsilon w) = \psi_1(z, w) + \varepsilon \psi_2(z, w) \\ &= (\phi_1(z, w) + \phi_2(z, w)e_8) + \varepsilon(\varphi_1(z, w) + \varphi_2(z, w)e_8), \end{aligned}$$

where

$$\psi_2(z, w) = \sum_{l=0}^{15} v_l e_l = \sum_{m=1}^8 g_m e_{2m-2} = \sum_{n=1}^4 G_n e_{4n-4} = \varphi_1 + \varphi_2 e_8 \in \mathcal{S}$$

with v_l ($l = 0, 1, \dots, 15$) are real-valued functions, $g_m = v_{2m-2} + v_{2m-1}e_1$ ($m = 1, 2, \dots, 8$) are complex-valued functions, $G_n = g_{2n-1} + g_{2n}e_2$ ($n = 1, 2, 3, 4$) are quaternion-valued functions, $\varphi_1 = G_1 + G_2e_4$ and $\varphi_2 = G_3 + G_4e_4$ are octonion-valued functions in \mathcal{S} .

Definition 3.2. Let Ω be an open set in $\mathcal{S} \times \mathcal{S}$. A function $F(\zeta)$ is said to be $L(R)$ -hyperholomorphic in Ω if the following two conditions are satisfied:

(1) $f_k(z)$, $g_k(w)$ ($k = 1, \dots, 8$) are continuously differential functions in Ω .

(2)

$$(3.4) \quad H^*F = 0 \quad (FH^* = 0) \text{ in } \Omega.$$

When we deal with a L -hyperholomorphic function $F(\zeta)$ in $\Omega \subset \mathcal{S} \times \mathcal{S}$, for simplicity, we often say that $F(\zeta)$ is a hyperholomorphic function in $\Omega \subset \mathcal{S} \times \mathcal{S}$.

The Equation (3.4) is applied to $F(\zeta)$ as follows:

$$\begin{aligned}
H^*F &= (D^* + \varepsilon E^*)(\psi_1(z, w) + \varepsilon \psi_2(z, w)) \\
&= (D^* \psi_1(z, w) + \varepsilon(E^* \psi_1(z, w) + D^* \psi_2(z, w))) \\
&= (D^*(F_1(z) + F_2(z)e_4 + F_3(z)e_8 + F_4(z)e_{12}) \\
&\quad + \varepsilon(E^*(F_1(z) + F_2(z)e_4 + F_3(z)e_8 + F_4(z)e_{12}) \\
&\quad + D^*(G_1(w) + G_2(w)e_4 + G_3(w)e_8 + G_4(w)e_{12}))) \\
&= \left(\frac{\partial f_1}{\partial P_1^*} - \frac{\partial \bar{f}_5}{\partial P_2^*} \right) + \left(\frac{\partial f_2}{\partial P_1^*} + \frac{\partial \bar{f}_6}{\partial P_2^*} \right) e_2 + \left(\frac{\partial f_3}{\partial P_1^*} + \frac{\partial \bar{f}_7}{\partial P_2^*} \right) e_4 \\
&\quad + \left(\frac{\partial f_4}{\partial P_1^*} + \frac{\partial \bar{f}_8}{\partial P_2^*} \right) e_6 + \left(\frac{\partial f_5}{\partial P_1^*} + \frac{\partial \bar{f}_1}{\partial P_2^*} \right) e_8 + \left(\frac{\partial f_6}{\partial P_1^*} - \frac{\partial \bar{f}_2}{\partial P_2^*} \right) e_{10} \\
&\quad + \left(\frac{\partial f_7}{\partial P_1^*} - \frac{\partial \bar{f}_3}{\partial P_2^*} \right) e_{12} + \left(\frac{\partial f_8}{\partial P_1^*} - \frac{\partial \bar{f}_4}{\partial P_2^*} \right) e_{14} \\
&\quad + \varepsilon \left(\left(\frac{\partial f_1}{\partial Q_1^*} - \frac{\partial \bar{f}_5}{\partial Q_2^*} + \frac{\partial g_1}{\partial P_1^*} - \frac{\partial \bar{g}_5}{\partial P_2^*} \right) + \left(\frac{\partial f_2}{\partial Q_1^*} + \frac{\partial \bar{f}_6}{\partial Q_2^*} + \frac{\partial g_2}{\partial P_1^*} + \frac{\partial \bar{g}_6}{\partial P_2^*} \right) e_2 \right. \\
&\quad + \left(\frac{\partial f_3}{\partial Q_1^*} + \frac{\partial \bar{f}_7}{\partial Q_2^*} + \frac{\partial g_3}{\partial P_1^*} + \frac{\partial \bar{g}_7}{\partial P_2^*} \right) e_4 + \left(\frac{\partial f_4}{\partial Q_1^*} + \frac{\partial \bar{f}_8}{\partial Q_2^*} + \frac{\partial g_4}{\partial P_1^*} + \frac{\partial \bar{g}_8}{\partial P_2^*} \right) e_6 \\
&\quad + \left(\frac{\partial f_5}{\partial Q_1^*} + \frac{\partial \bar{f}_1}{\partial Q_2^*} + \frac{\partial g_5}{\partial P_1^*} + \frac{\partial \bar{g}_1}{\partial P_2^*} \right) e_8 + \left(\frac{\partial f_6}{\partial Q_1^*} - \frac{\partial \bar{f}_2}{\partial Q_2^*} + \frac{\partial g_6}{\partial P_1^*} - \frac{\partial \bar{g}_2}{\partial P_2^*} \right) e_{10} \\
&\quad \left. + \left(\frac{\partial f_7}{\partial Q_1^*} - \frac{\partial \bar{f}_3}{\partial Q_2^*} + \frac{\partial g_7}{\partial P_1^*} - \frac{\partial \bar{g}_3}{\partial P_2^*} \right) e_{12} + \left(\frac{\partial f_8}{\partial Q_1^*} - \frac{\partial \bar{f}_4}{\partial Q_2^*} + \frac{\partial g_8}{\partial P_1^*} - \frac{\partial \bar{g}_4}{\partial P_2^*} \right) e_{14} \right).
\end{aligned}$$

If the following equations

$$\begin{aligned}
\frac{\partial f_1}{\partial P_1^*} &= \frac{\partial \bar{f}_5}{\partial P_2^*}, \quad \frac{\partial f_2}{\partial P_1^*} = -\frac{\partial \bar{f}_6}{\partial P_2^*}, \quad \frac{\partial f_3}{\partial P_1^*} = -\frac{\partial \bar{f}_7}{\partial P_2^*}, \quad \frac{\partial f_4}{\partial P_1^*} = -\frac{\partial \bar{f}_8}{\partial P_2^*}, \\
\frac{\partial f_5}{\partial P_1^*} &= -\frac{\partial \bar{f}_1}{\partial P_2^*}, \quad \frac{\partial f_6}{\partial P_1^*} = \frac{\partial \bar{f}_2}{\partial P_2^*}, \quad \frac{\partial f_7}{\partial P_1^*} = \frac{\partial \bar{f}_3}{\partial P_2^*}, \quad \frac{\partial f_8}{\partial P_1^*} = \frac{\partial \bar{f}_4}{\partial P_2^*}, \\
\frac{\partial g_1}{\partial P_1^*} + \frac{\partial f_1}{\partial Q_1^*} &= \frac{\partial \bar{g}_5}{\partial P_2^*} + \frac{\partial \bar{f}_5}{\partial Q_2^*}, \quad \frac{\partial g_2}{\partial P_1^*} + \frac{\partial f_2}{\partial Q_1^*} = -\frac{\partial \bar{g}_6}{\partial P_2^*} - \frac{\partial \bar{f}_6}{\partial Q_2^*}, \\
\frac{\partial g_3}{\partial P_1^*} + \frac{\partial f_3}{\partial Q_1^*} &= -\frac{\partial \bar{g}_7}{\partial P_2^*} - \frac{\partial \bar{f}_7}{\partial Q_2^*}, \quad \frac{\partial g_4}{\partial P_1^*} + \frac{\partial f_4}{\partial Q_1^*} = -\frac{\partial \bar{g}_8}{\partial P_2^*} - \frac{\partial \bar{f}_8}{\partial Q_2^*}, \\
\frac{\partial g_5}{\partial P_1^*} + \frac{\partial f_5}{\partial Q_1^*} &= -\frac{\partial \bar{g}_1}{\partial P_2^*} - \frac{\partial \bar{f}_1}{\partial Q_2^*}, \quad \frac{\partial g_6}{\partial P_1^*} + \frac{\partial f_6}{\partial Q_1^*} = \frac{\partial \bar{g}_2}{\partial P_2^*} + \frac{\partial \bar{f}_2}{\partial Q_2^*}, \\
(3.5) \quad \frac{\partial g_7}{\partial P_1^*} + \frac{\partial f_7}{\partial Q_1^*} &= \frac{\partial \bar{g}_3}{\partial P_2^*} + \frac{\partial \bar{f}_3}{\partial Q_2^*}, \quad \frac{\partial g_8}{\partial P_1^*} + \frac{\partial f_8}{\partial Q_1^*} = \frac{\partial \bar{g}_4}{\partial P_2^*} + \frac{\partial \bar{f}_4}{\partial Q_2^*},
\end{aligned}$$

are satisfied, the function $F(\zeta)$ is a hyperholomorphic function on Ω . These are the corresponding s-Cauchy-Riemann equations on $\mathcal{S} \times \mathcal{S}$.

Definition 3.3. Let Ω be an open set in $\mathcal{S} \times \mathcal{S}$ and let

$$F = (f_1, f_2, \dots, f_8, g_1, g_2, \dots, g_8) : \Omega \rightarrow \mathcal{S} \times \mathcal{S}.$$

This mapping is said to be harmonic if all its components f_j and g_j ($j = 1, 2, \dots, 8$) of F are harmonic on Ω . The system (3.5) is called a generalized Cauchy-Riemann system if every solution $F(\zeta)$ has only harmonic components f_j and g_j ($j = 1, 2, \dots, 8$).

Remark 3.1. We redefine the Equations (3.5) as follows:

$$(3.6) \quad \begin{aligned} \frac{\partial f_1}{\partial \bar{z}_1} &= \frac{\partial \bar{f}_5}{\partial \bar{z}_5}, & \frac{\partial f_1}{\partial z_2} &= \frac{\partial \bar{f}_5}{\partial z_6}, & \frac{\partial f_1}{\partial z_3} &= \frac{\partial \bar{f}_5}{\partial z_7}, & \frac{\partial f_1}{\partial z_4} &= \frac{\partial \bar{f}_5}{\partial z_8}, \\ \frac{\partial f_2}{\partial \bar{z}_1} &= -\frac{\partial \bar{f}_6}{\partial \bar{z}_5}, & \frac{\partial f_2}{\partial z_2} &= -\frac{\partial \bar{f}_6}{\partial z_6}, & \frac{\partial f_2}{\partial z_3} &= -\frac{\partial \bar{f}_6}{\partial z_7}, & \frac{\partial f_2}{\partial z_4} &= -\frac{\partial \bar{f}_6}{\partial z_8}, \\ \frac{\partial f_3}{\partial \bar{z}_1} &= -\frac{\partial \bar{f}_7}{\partial \bar{z}_5}, & \frac{\partial f_3}{\partial z_2} &= -\frac{\partial \bar{f}_7}{\partial z_6}, & \frac{\partial f_3}{\partial z_3} &= -\frac{\partial \bar{f}_7}{\partial z_7}, & \frac{\partial f_3}{\partial z_4} &= -\frac{\partial \bar{f}_7}{\partial z_8}, \\ \frac{\partial f_4}{\partial \bar{z}_1} &= -\frac{\partial \bar{f}_8}{\partial \bar{z}_5}, & \frac{\partial f_4}{\partial z_2} &= -\frac{\partial \bar{f}_8}{\partial z_6}, & \frac{\partial f_4}{\partial z_3} &= -\frac{\partial \bar{f}_8}{\partial z_7}, & \frac{\partial f_4}{\partial z_4} &= -\frac{\partial \bar{f}_8}{\partial z_8}, \\ \frac{\partial f_5}{\partial \bar{z}_1} &= -\frac{\partial \bar{f}_1}{\partial \bar{z}_5}, & \frac{\partial f_5}{\partial z_2} &= -\frac{\partial \bar{f}_1}{\partial z_6}, & \frac{\partial f_5}{\partial z_3} &= -\frac{\partial \bar{f}_1}{\partial z_7}, & \frac{\partial f_5}{\partial z_4} &= -\frac{\partial \bar{f}_1}{\partial z_8}, \\ \frac{\partial f_6}{\partial \bar{z}_1} &= \frac{\partial \bar{f}_2}{\partial \bar{z}_5}, & \frac{\partial f_6}{\partial z_2} &= \frac{\partial \bar{f}_2}{\partial z_6}, & \frac{\partial f_6}{\partial z_3} &= \frac{\partial \bar{f}_2}{\partial z_7}, & \frac{\partial f_6}{\partial z_4} &= \frac{\partial \bar{f}_2}{\partial z_8}, \\ \frac{\partial f_7}{\partial \bar{z}_1} &= \frac{\partial \bar{f}_3}{\partial \bar{z}_5}, & \frac{\partial f_7}{\partial z_2} &= \frac{\partial \bar{f}_3}{\partial z_6}, & \frac{\partial f_7}{\partial z_3} &= \frac{\partial \bar{f}_3}{\partial z_7}, & \frac{\partial f_7}{\partial z_4} &= \frac{\partial \bar{f}_3}{\partial z_8}, \\ \frac{\partial f_8}{\partial \bar{z}_1} &= \frac{\partial \bar{f}_4}{\partial \bar{z}_5}, & \frac{\partial f_8}{\partial z_2} &= \frac{\partial \bar{f}_4}{\partial z_6}, & \frac{\partial f_8}{\partial z_3} &= \frac{\partial \bar{f}_4}{\partial z_7}, & \frac{\partial f_8}{\partial z_4} &= \frac{\partial \bar{f}_4}{\partial z_8}. \end{aligned}$$

We call that the Equations (3.6) are the condition of harmonicity of the hyperholomorphic function $F(\zeta)$ on Ω in $\mathcal{S} \times \mathcal{S}$.

Remark 3.2. We redefine the Equations (3.5) as follows:

$$\begin{aligned} \frac{\partial f_1}{\partial \bar{z}_1} &= \frac{\partial \bar{f}_3}{\partial \bar{z}_3}, & \frac{\partial f_1}{\partial \bar{z}_2} &= -\frac{\partial \bar{f}_3}{\partial \bar{z}_4}, & \frac{\partial f_3}{\partial \bar{z}_1} &= -\frac{\partial \bar{f}_1}{\partial \bar{z}_3}, & \frac{\partial f_3}{\partial \bar{z}_2} &= \frac{\partial \bar{f}_1}{\partial \bar{z}_4}, \\ \frac{\partial f_2}{\partial \bar{z}_1} &= \frac{\partial \bar{f}_4}{\partial \bar{z}_3}, & \frac{\partial f_2}{\partial \bar{z}_2} &= -\frac{\partial \bar{f}_4}{\partial \bar{z}_4}, & \frac{\partial f_4}{\partial \bar{z}_1} &= -\frac{\partial \bar{f}_2}{\partial \bar{z}_3}, & \frac{\partial f_4}{\partial \bar{z}_2} &= \frac{\partial \bar{f}_2}{\partial \bar{z}_4}, \\ \frac{\partial g_1}{\partial \bar{z}_1} + \frac{\partial f_1}{\partial \bar{w}_1} &= \frac{\partial \bar{g}_3}{\partial \bar{z}_3} + \frac{\partial \bar{f}_3}{\partial \bar{w}_3}, & \frac{\partial g_1}{\partial \bar{z}_2} + \frac{\partial f_1}{\partial \bar{w}_2} &= -\frac{\partial \bar{g}_3}{\partial \bar{z}_4} - \frac{\partial \bar{f}_3}{\partial \bar{w}_4}, \end{aligned}$$

$$\begin{aligned}
\frac{\partial g_3}{\partial \bar{z}_1} + \frac{\partial f_3}{\partial \bar{w}_1} &= -\frac{\partial \bar{g}_1}{\partial z_3} - \frac{\partial \bar{f}_1}{\partial w_3}, & \frac{\partial g_3}{\partial \bar{z}_2} + \frac{\partial f_3}{\partial \bar{w}_2} &= \frac{\partial \bar{g}_1}{\partial \bar{z}_4} + \frac{\partial \bar{f}_1}{\partial \bar{w}_4}, \\
\frac{\partial g_2}{\partial \bar{z}_1} + \frac{\partial f_2}{\partial \bar{w}_1} &= \frac{\partial \bar{g}_4}{\partial z_3} + \frac{\partial \bar{f}_4}{\partial w_3}, & \frac{\partial g_2}{\partial \bar{z}_2} + \frac{\partial f_2}{\partial \bar{w}_2} &= -\frac{\partial \bar{g}_4}{\partial \bar{z}_4} - \frac{\partial \bar{f}_4}{\partial \bar{w}_4}, \\
\frac{\partial g_4}{\partial \bar{z}_1} + \frac{\partial f_4}{\partial \bar{w}_1} &= -\frac{\partial \bar{g}_2}{\partial z_3} - \frac{\partial \bar{f}_2}{\partial w_3}, & \frac{\partial g_4}{\partial \bar{z}_2} + \frac{\partial f_4}{\partial \bar{w}_2} &= \frac{\partial \bar{g}_2}{\partial \bar{z}_4} + \frac{\partial \bar{f}_2}{\partial \bar{w}_4}.
\end{aligned}$$

Now, we need the following:

$$(3.7) \quad \begin{aligned}
\frac{\partial g_1}{\partial \bar{z}_1} &= \frac{\partial \bar{g}_3}{\partial z_3}, & \frac{\partial g_1}{\partial \bar{z}_2} &= -\frac{\partial \bar{g}_3}{\partial \bar{z}_4}, & \frac{\partial g_3}{\partial \bar{z}_1} &= -\frac{\partial \bar{g}_1}{\partial z_3}, & \frac{\partial g_3}{\partial \bar{z}_2} &= \frac{\partial \bar{g}_1}{\partial \bar{z}_4}, \\
\frac{\partial g_2}{\partial \bar{z}_1} &= \frac{\partial \bar{g}_4}{\partial z_3}, & \frac{\partial g_2}{\partial \bar{z}_2} &= -\frac{\partial \bar{g}_4}{\partial \bar{z}_4}, & \frac{\partial g_4}{\partial \bar{z}_1} &= -\frac{\partial \bar{g}_2}{\partial z_3}, & \frac{\partial g_4}{\partial \bar{z}_2} &= \frac{\partial \bar{g}_2}{\partial \bar{z}_4}.
\end{aligned}$$

We call that the Equations (3.6) and (3.7) are the condition of harmonicity.

Theorem 3.2. *If the hyperholomorphic function*

$$\begin{aligned}
F(\zeta) &= f_1(z) + f_2(z)e_2 + f_3(z)e_4 + f_4(z)e_6 + f_5(z)e_8 + f_6(z)e_{10} \\
&\quad + f_7(z)e_{12} + f_8(z)e_{14} + \varepsilon(g_1(w) + g_2(w)e_2 + g_3(w)e_4 \\
&\quad + g_4(w)e_6 + g_5(w)e_8 + g_6(w)e_{10} + g_7(w)e_{12} + g_8(w)e_{14})
\end{aligned}$$

satisfies the condition of harmonicity (3.6) and (3.7) in an open set Ω in $\mathcal{S} \times \mathcal{S}$, then the functions $f_k(z)$, $g_k(w)$ ($k = 1, 2, \dots, 8$) are harmonic in Ω .

Proof. We have

$$\begin{aligned}
HH^* f_1 &= (D + \varepsilon E)(D^* + \varepsilon E^*) f_1 = (DD^* + \varepsilon(DE^* + D^*E)) f_1 \\
&= \frac{\partial^2 f_1}{\partial z_1 \partial \bar{z}_1} + \frac{\partial^2 f_1}{\partial z_2 \partial \bar{z}_2} + \frac{\partial^2 f_1}{\partial z_3 \partial \bar{z}_3} + \frac{\partial^2 f_1}{\partial z_4 \partial \bar{z}_4} + \frac{\partial^2 f_1}{\partial z_5 \partial \bar{z}_5} + \frac{\partial^2 f_1}{\partial z_6 \partial \bar{z}_6} \\
&\quad + \frac{\partial^2 f_1}{\partial z_7 \partial \bar{z}_7} + \frac{\partial^2 f_1}{\partial z_8 \partial \bar{z}_8} + \varepsilon \left(\frac{\partial^2 f_1}{\partial z_1 \partial \bar{w}_1} + \frac{\partial^2 f_1}{\partial z_2 \partial \bar{w}_2} + \frac{\partial^2 f_1}{\partial z_3 \partial \bar{w}_3} + \frac{\partial^2 f_1}{\partial z_4 \partial \bar{w}_4} \right. \\
&\quad + \frac{\partial^2 f_1}{\partial z_5 \partial \bar{w}_5} + \frac{\partial^2 f_1}{\partial z_6 \partial \bar{w}_6} + \frac{\partial^2 f_1}{\partial z_7 \partial \bar{w}_7} + \frac{\partial^2 f_1}{\partial z_8 \partial \bar{w}_8} + \frac{\partial^2 f_1}{\partial \bar{z}_1 \partial w_1} + \frac{\partial^2 f_1}{\partial \bar{z}_2 \partial w_2} \\
&\quad \left. + \frac{\partial^2 f_1}{\partial \bar{z}_3 \partial w_3} + \frac{\partial^2 f_1}{\partial \bar{z}_4 \partial w_4} + \frac{\partial^2 f_1}{\partial \bar{z}_5 \partial w_5} + \frac{\partial^2 f_1}{\partial \bar{z}_6 \partial w_6} + \frac{\partial^2 f_1}{\partial \bar{z}_7 \partial w_7} + \frac{\partial^2 f_1}{\partial \bar{z}_8 \partial w_8} \right)
\end{aligned}$$

$$\begin{aligned}
 &= \frac{\partial}{\partial z_1} \left(\frac{\partial \bar{f}_5}{\partial \bar{z}_5} \right) + \frac{\partial}{\partial \bar{z}_2} \left(\frac{\partial \bar{f}_5}{\partial z_6} \right) + \frac{\partial}{\partial \bar{z}_3} \left(\frac{\partial \bar{f}_5}{\partial z_7} \right) + \frac{\partial}{\partial \bar{z}_4} \left(\frac{\partial \bar{f}_5}{\partial z_8} \right) \\
 &\quad + \frac{\partial}{\partial \bar{z}_5} \left(-\frac{\partial \bar{f}_5}{\partial z_1} \right) + \frac{\partial}{\partial z_6} \left(-\frac{\partial \bar{f}_5}{\partial \bar{z}_2} \right) + \frac{\partial}{\partial z_7} \left(-\frac{\partial \bar{f}_5}{\partial \bar{z}_3} \right) + \frac{\partial}{\partial z_8} \left(-\frac{\partial \bar{f}_5}{\partial \bar{z}_4} \right) \\
 &\quad + \varepsilon \left(\left(\frac{\partial}{\partial z_1} \left(\frac{\partial \bar{f}_5}{\partial \bar{w}_5} \right) + \frac{\partial}{\partial \bar{z}_2} \left(\frac{\partial \bar{f}_5}{\partial w_6} \right) + \frac{\partial}{\partial \bar{z}_3} \left(\frac{\partial \bar{f}_5}{\partial w_7} \right) + \frac{\partial}{\partial \bar{z}_4} \left(\frac{\partial \bar{f}_5}{\partial w_8} \right) \right. \right. \\
 &\quad \left. \left. + \frac{\partial}{\partial \bar{z}_5} \left(-\frac{\partial \bar{f}_5}{\partial w_1} \right) + \frac{\partial}{\partial z_6} \left(-\frac{\partial \bar{f}_5}{\partial \bar{w}_2} \right) + \frac{\partial}{\partial z_7} \left(-\frac{\partial \bar{f}_5}{\partial \bar{w}_3} \right) + \frac{\partial}{\partial z_8} \left(-\frac{\partial \bar{f}_5}{\partial \bar{w}_4} \right) \right) \right) \\
 &\quad + \left(\frac{\partial}{\partial w_1} \left(\frac{\partial \bar{f}_5}{\partial \bar{z}_5} \right) + \frac{\partial}{\partial \bar{w}_2} \left(\frac{\partial \bar{f}_5}{\partial z_6} \right) + \frac{\partial}{\partial \bar{w}_3} \left(\frac{\partial \bar{f}_5}{\partial z_7} \right) + \frac{\partial}{\partial \bar{w}_4} \left(\frac{\partial \bar{f}_5}{\partial z_8} \right) \right. \\
 &\quad \left. + \frac{\partial}{\partial \bar{w}_5} \left(-\frac{\partial \bar{f}_5}{\partial z_1} \right) + \frac{\partial}{\partial w_6} \left(-\frac{\partial \bar{f}_5}{\partial \bar{z}_2} \right) + \frac{\partial}{\partial w_7} \left(-\frac{\partial \bar{f}_5}{\partial \bar{z}_3} \right) + \frac{\partial}{\partial w_8} \left(-\frac{\partial \bar{f}_5}{\partial \bar{z}_4} \right) \right) \\
 &= 0,
 \end{aligned}$$

and the functions $f_k(z), g_k(w)$ ($k = 1, 2, \dots, 8$) are proved by the similar method as in the proof of the case of f_1 . \square

Proposition 3.1. *Let Ω be a domain in $\mathcal{S} \times \mathcal{S}$ and $F(\zeta)$ be a L -hyperholomorphic functions in Ω . If $HH^*F(\zeta) = 0$, then the system (3.5) is a generalized Cauchy-Riemann system in Ω .*

Proof. If $HH^*F = 0$, then $\sum_{j=1}^8 \left(\frac{\partial^2 f}{\partial z_j \partial \bar{z}_j} + \frac{\partial^2 f}{\partial w_j \partial \bar{w}_j} \right) = 0$. Therefore, the components f_j, g_j ($j = 1, 2, \dots, 8$) are harmonic in Ω . That is, the system (3.5) is a generalized Cauchy-Riemann system on Ω . \square

References

- [1] L. Hörmander, *An introduction to complex analysis in several variables*, North-Holland, Amsterdam, 1966.
- [2] H. S. Jung, S. J. Ha, K. H. Lee, S. M. Lim and K. H. Shon, *Structures of hyperholomorphic functions on dual quaternion numbers*, Honam Mathematical J. **35** (2013), 809-817.
- [3] J. E. Kim, S. J. Lim and K. H. Shon, *Regular functions with values in ternary number system on the complex Clifford analysis*, Abstract and Applied Analysis. (2013) Article ID 136120, 7 Pages.
- [4] J. E. Kim, S. J. Lim and K. H. Shon, *Regularity of functions on the reduced quaternion field in Clifford analysis*, Abstract and Applied Analysis. (2014) Article ID 654798, 8 Pages.
- [5] J. E. Kim and K. H. Shon, *The regularity functionson dual split quaternions in Clifford analysis*, Abstract and Applied Analysis. (2014) Article ID 369430, 8 Pages.

- [6] S. G. Krantz, *Funtion theory of several complex variables*, Am. Math. Soc, Providence. (2001).
- [7] S. J. Lim and K. H. Shon, *Hyperholomorphic functions and hyperconjugate harmonic functions of octonion variables*, J. Inequ. Appl. **77** (2013), 1-8.
- [8] K. Nôno, *Hyperholomorphic functions of a quaternion variable*, Bull. Fukuoka Univ. Ed. **32** (1983), 21-37.
- [9] K. Nôno, *On the octonionic linearization of Laplacian and octonionic function theory*, Bull. Fukuoka Univ. Ed. **37** (1988), 1-15.
- [10] S. H. Park and K. H. Shon, *Sedenion functions of hypercomplex variables in the sense of Clifford analysis*, East Asian Math. J. **29(5)** (2013), 521-527.

Ji Eun Kim

Department of Mathematics, Pusan National University,
Busan 609-735, Korea.

E-mail: jeunkim@pusan.ac.kr

Su Jin Ha

Department of Mathematics, Pusan National University,
Busan 609-735, Korea.

E-mail: adorm84@naver.com

Kwang Ho Shon

Department of Mathematics, Pusan National University,
Busan 609-735, Korea.

E-mail: khshon@pusan.ac.kr