

# 1 Introduction

*I would like to expand this section and flesh out the examples a bit more to make them clearer.*

The basic idea underpinning the game semantics for nondeterministic programming languages is to remove the constraint of *determinism* from the usual definition of a strategy (see, for example, [AJ13]) that says that the player may have at most one reply to each move made by the opponent. The determinism constraint is of little technical value and its main purpose is to provide a fully abstract denotational semantics for deterministic programming languages; if we remove it in order to study nondeterministic languages, then much of the technical apparatus of game semantics carries over immediately.

One problem that does arise, however, is that the resulting semantics no longer adequately captures the possibility of divergence in the interaction. In deterministic game semantics, divergence is represented by the absence of a reply to an opponent's move, but this is not adequate in the nondeterministic case because it gives us no way to capture a *possible* divergence. For example, if we represent divergence by the absence of a reply then Program A, which constantly returns the numeral 1, and Program B, which makes a nondeterministic choice and then either returns 1 or diverges, have the same semantics, despite exhibiting different behaviour [HM99].

We therefore need a new way to keep track of divergence. This was done by Harmer and McCusker [HM99], who resolved the problem by adding extra data to the definition of a strategy that records which opponent positions allow the possibility of divergence. Using this device, Harmer and McCusker were able to provide a fully abstract game semantics for PCF with finite nondeterminism.

There is, however, a technical subtlety that arises when we try to use the same technique to model PCF with *unbounded* nondeterminism; that is, PCF with the addition of a nondeterministic natural number generator that never diverges. Indeed, if we naively relax the Harmer-McCusker model to allow infinite branching then composition of strategies is no longer associative [Har99].

Ultimately, the reason why unbounded nondeterminism introduces problems is that it makes the language fine-grained enough to distinguish between computation that is *divergent*; i.e., a computation that carries on for infinitely many steps and does not terminate, and computation that is convergent, but may take an unbounded number of steps to do so. For example, using recursion, we may define a function that, at each step, makes a nondeterministic choice whether to call its argument or to return. If we have access to unbounded nondeterminism then we may write a function that nondeterministically chooses a natural number  $n$ , calls its argument  $n$  times and then returns. Observationally, these programs are different: the first may diverge, while the second always converges.

In the usual model of strategies as sets of positions, the strategies defined by these two programs are the same.

At first glance, it appears that the Harmer-McCusker model is fine-grained enough to distinguish these two strategies, and this is true if we define the strategies directly: the strategy denoting the first program ought to designate the opening move as divergent, whereas the strategy denoting the second program should not designate any positions as divergent. However, if we want to define these strategies compositionally, it turns out that both strategies are designated divergent, since the model treats an infinite increasing sequence of positions as a divergence occurring from livelock (the ‘infinite chattering’ in the game semantics). This inconsistency between the direct definition of the strategy and the compositional one is what leads to the failure of composition to be associative.

We need, then, some way to distinguish between computation that takes infinitely many steps and computation that takes a finite, but arbitrarily large, number of steps. One approach, taken by Tsukada and Ong [TO15] is to represent strategies as trees, rather than sets of positions; here, there is a distinction between a tree of infinite height and an infinite family of trees of finite but arbitrarily large height. We take a different approach, one suggested by the work of Roscoe and Levy ([ROS93] and [Lev08], cited in [TO15]), and by Laird’s sequential algorithms model for PCF with unbounded nondeterminism [Lai15], and enlarge our strategies to include information about the infinite positions that may occur in them.

## 1.1 Related Work

A more detailed overview of previous work, particularly [Lai15].

# 2 PCF with Countable Nondeterminism

As in [Lai15].

## 2.1 Type Theory

## 2.2 Operational Semantics

## 2.3 Must Testing

We are interested in soundness and adequacy for the ‘must-converge’ relation, since the ‘may-converge’ relation can be adequately modelled using finite non-determinism.

### 3 Game Semantics

#### 3.1 Games

**Definition 3.1.** Given a set  $X$ , we write  $\overline{X^*}$  for the set of all (finite and infinite) sequences of elements of  $X$  of length  $\leq \omega$ .

A *game* is given by a tuple

$$A = (M_A, \lambda_A, \zeta_A, P_A)$$

where

- $M_A$  is a set of moves.
- $\lambda_A: M_A \rightarrow \{O, P\}$  designates each move as an  $O$ -move (played by Player  $O$ ) or a  $P$ -move (played by Player  $P$ ).
- $P_A \subseteq \overline{M_A^*}$  is a non-empty prefix-closed set of sequences of moves from  $M_A$  of length  $\leq \omega$ , called *positions*. We call a sequence  $s \in P_A$  *infinite* if  $\text{length}(s) = \omega$ . Otherwise, we say that  $s$  is *finite*.
- $\zeta_A: P_A \rightarrow \{O, P\}$  designates each position in  $P_A$  as a  $P$ -position or an  $O$ -position.

The set  $P_A$  and the functions  $\lambda_A, \zeta_A$  are required to satisfy certain additional axioms:

**Compactness** If  $s \in M_A^\omega$  is an infinite sequence of moves such that  $t \in P_A$  for all finite prefixes  $t \sqsubset s$ , then  $s \in P_A$ .

**Consistency** If  $a \in M_A$  and  $sa \in P_A$  then  $\zeta_A(sa) = \lambda_A(a)$ .

**Alternation** If  $a \in M_A$  and  $sa \in P_A$  then  $\zeta_A(s) = \neg \zeta_A(sa)$ .

**Negativity**  $\zeta_A(\epsilon) = P$ , where  $\epsilon$  is the empty position.

Note that the negativity rule and the alternation rule between them imply that all starting moves are made by player  $O$ .

We shall write  $P_A^P = \zeta_A^{-1}(\{P\})$  for the set of  $P$ -positions in  $A$  and  $P_A^O = \zeta_A^{-1}(\{O\})$  for the set of  $O$ -positions in  $A$ .

#### 3.2 Strategies

A nondeterministic *strategy* for a game  $A$  will be given by a pair  $\sigma = (T_\sigma, D_\sigma)$ , where

- $T_\sigma$  is the set of *traces* of the strategy; i.e., the set of positions that can occur when player  $P$  plays according to the strategy. That is,  $T_\sigma$  is a

non-empty prefix-closed subset of  $P_A$  that is closed under  $O$ -replies to  $P$ -positions: whenever  $s \in T_\sigma$  is a  $P$ -position and  $a \in M_A$  is an  $O$ -move such that  $sa \in P_A$ , then  $sa \in T_\sigma$ .

- $D_\sigma \subseteq T_\sigma$  denotes the set of *divergences* in  $T_\sigma$ ; that is, the set of those  $O$ -positions in  $T_\sigma$  for which Player  $P$  may elect to diverge rather than respond. Accordingly,  $\zeta_A(s) = O$  for all minimal  $s \in D_\sigma$ . If a trace has a divergent prefix, then it is divergent, so we shall follow Roscoe and insist that  $D_\sigma$  be postfix-closed in  $P_A$  ([ROS93], cited in [HM99]).

**Definition 3.2.** Let  $A = (M_A, \lambda_A, \zeta_A, P_A)$  be a game. Then a (nondeterministic) *strategy* for  $A$  is a pair  $\sigma = (T_\sigma, D_\sigma)$  such that

- $T_\sigma \subseteq P_A$  is a prefix-closed subset such that whenever  $s \in T_\sigma$  is a  $P$ -position and  $sa \in P_A$  for some move  $a$ , then  $sa \in T_\sigma$ .
- $D_\sigma \subseteq P_A$  is a postfix-closed subset such that all of the minimal positions in  $D_\sigma$  are  $O$ -positions contained in  $T_\sigma$ . In other words, if  $s \in D_\sigma$ , then there is some  $O$ -position  $t \sqsubseteq s$  such that  $t \in T_\sigma \cap D_\sigma$ .

$T_\sigma$  and  $D_\sigma$  are required to satisfy the following totality axiom:

**Diverge-or-reply** If  $s \in T_\sigma$  is an  $O$ -position then either  $sa \in T_\sigma$  for some  $P$ -move  $a$  or  $s \in D_\sigma$ .

The diverge-or-reply rule tells us that a function must either return a value or go into an infinite loop, and corresponds to the absence of an error constant in our language.

A consequence of the diverge-or-reply rule is that if  $s \in \sigma$  is an infinite  $O$ -position, then  $s \in D_\sigma$  (since it is impossible for player  $P$  to reply to it, positions of length  $\omega + 1$  or greater being disallowed). This works well for our purposes; it captures our intuition that divergence should arise from some infinite behaviour, but it excludes those cases where Player  $O$  is ‘to blame’ for the infinite behaviour (passing, for example, a divergent function into the non-divergent strategy  $\lambda f.f$ ).

We will need one extra condition on the definition of a strategy in order to make sure that the composition of two strategies satisfies the diverge-or-reply rule.

**Definition 3.3.** We say a strategy  $\sigma$  is *compact* if it satisfies the compactness criterion given above for games: if  $s \in M_A^\omega$  is an infinite sequence such that  $t \in T_\sigma$  for all finite prefixes  $t \sqsubset s$ , then  $t \in T_\sigma$ .

Compactness is too strong a condition for our purposes: we shall later see that it is not preserved by composition. We shall use the following local version instead:

**Definition 3.4.** We say a strategy  $\sigma$  is *locally compact* if whenever  $s \in T_\sigma$  there is some compact strategy  $\sigma'$  such that  $s \in T_{\sigma'} \subseteq T_\sigma$  and  $D_\sigma \subseteq D_{\sigma'}$ .

From now on, we shall use the word *strategy* to mean *locally compact strategy*.

### 3.3 Multiplicatives

Let  $A, B$  be games. If  $s \in \overline{(M_A \sqcup M_B)^*}$  then we write  $s|_A$  for the subsequence of  $s$  consisting of all moves from  $A$  and  $s|_B$  for the subsequence of  $s$  consisting of all moves from  $B$ . We write  $P_A \parallel P_B$  for the set of all such sequences  $s$  such that  $s|_A \in P_A$  and  $s|_B \in P_B$ .

**Definition 3.5.** Let  $A, B$  be games. Then the *tensor product* of  $A$  and  $B$  is the game  $A \otimes B = (M_{A \otimes B}, \lambda_{A \otimes B}, \zeta_{A \otimes B}, P_{A \otimes B})$  where we define  $M_{A \otimes B}$ ,  $\lambda_{A \otimes B}$ ,  $\zeta_{A \otimes B}$ ,  $P_{A \otimes B}$  as follows:

- $M_{A \otimes B} = M_A \sqcup M_B$
- $\lambda_{A \otimes B} = \lambda_A \sqcup \lambda_B$
- For  $s \in P_A \parallel P_B$ ,  $\zeta_{A \otimes B}(s) = \zeta_A(s|_A) \wedge \zeta_B(s|_B)$ .
- $P_{A \otimes B}$  is the set of all sequences  $s \in P_A \parallel P_B$  such that  $s$  satisfies the alternating condition with respect to  $\zeta_{A \otimes B}$ : i.e., for all  $t, ta \sqsubseteq s$ ,  $\zeta_{A \otimes B}(t) = \neg \zeta_{A \otimes B}(ta)$ .

**Definition 3.6.** Let  $A, B$  be games. Then the *linear implication* of  $A$  and  $B$  is the game  $A \multimap B = (M_{A \multimap B}, \lambda_{A \multimap B}, \zeta_{A \multimap B}, P_{A \multimap B})$  where we define  $M_{A \multimap B}$ ,  $\lambda_{A \multimap B}$ ,  $\zeta_{A \multimap B}$ ,  $P_{A \multimap B}$  as follows:

- $M_{A \multimap B} = M_A \sqcup M_B$
- $\lambda_{A \multimap B} = (\neg \circ \lambda_A) \sqcup \lambda_B$
- For  $s \in P_A \parallel P_B$ ,  $\zeta_{A \multimap B}(s) = \zeta_A(s|_A) \wedge \zeta_B(s|_B)$ .
- $P_{A \multimap B}$  is the set of all sequences  $s \in P_A \parallel P_B$  such that  $s$  satisfies the alternating condition with respect to  $\zeta_{A \multimap B}$ .

We want to prove that these are well formed games.

**Proposition 3.7.** *If  $A, B$  are games then  $A \otimes B$  and  $A \multimap B$  are games.*

First, we will prove a useful lemma.

**Lemma 3.8.** *Let  $A, B$  be games. Given a position  $s \in P_A \parallel P_B$ , we define the sign profile of  $s$  to be given by the signs of its two components:*

| $\zeta_A(s _A)$ | $\zeta_B(s _B)$ | sign profile |
|-----------------|-----------------|--------------|
| $P$             | $P$             | $PP$         |
| $O$             | $P$             | $OP$         |
| $P$             | $O$             | $PO$         |
| $O$             | $O$             | $OO$         |

If  $s \in P_{A \otimes B}$  is a finite position, then  $s$  cannot have sign profile  $OO$ . If  $s \in P_{A \multimap B}$  is a finite position then  $s$  cannot have sign profile  $OP$ .

*Proof.* By negativity, the empty sequence  $\epsilon$  always has sign profile  $PP$ . Otherwise,  $s = s'a$  for some sequence  $s'$  and some move  $a$ . If  $s \in P_{A \otimes B}$  and  $s$  has sign profile  $OO$ , then  $s$  is an  $O$ -position in  $A \otimes B$ . Therefore,  $s'$  is a  $P$ -position in  $A \otimes B$  and so  $s'$  must have sign profile  $PP$ , by the definition of  $\zeta_{A \otimes B}$ . But this is impossible, as the move  $a$  can only change the sign profile in one of the two components.

The proof for the linear implication is entirely similar.  $\square$

*Proof of Proposition 3.7.* Alternation is part of the definition and negativity is implied by the equations  $P \wedge P = P$  and  $P \Rightarrow P = P$ . The only tricky part is proving consistency of the  $\lambda$  and  $\zeta$  functions. Since consistency applies only to positions with a last move, we may ignore the infinite plays in  $A$  and  $B$ .

Suppose  $sa \in P_{A \otimes B}$ . We wish to show that  $\zeta_A(sa) = \lambda_A(a)$ . In the presence of alternation, this is equivalent to showing that  $\zeta_A(s) = \neg \lambda_A(a)$ . We show this by cases on the sign profile of  $s$ . By Lemma 3.8,  $s$  has sign profile  $PP$ ,  $OP$  or  $PO$ . If  $s$  has sign profile  $PP$  then  $s$  is a  $P$ -position. Since  $s|_A$  and  $s|_B$  are both  $P$ -positions,  $a$  must be an  $O$ -move, whether it occurs in  $A$  or  $B$ .

If instead  $s$  has sign profile  $OP$  or  $PO$ , then  $s$  is an  $O$ -position. By alternation,  $sa$  must be a  $P$ -position, so it must have sign profile  $PP$ . By consistency and alternation for the games  $A, B$ , the move  $a$  must be a  $P$ -move.

For compactness, suppose that  $s \in M_{A \otimes B}^\omega$  is an infinite sequence of moves such that  $t \in P_{A \otimes B}$  for all finite prefixes  $t \sqsubset s$ . Then  $s$  is certainly alternating. Moreover, for all finite prefixes  $t \sqsubset s|_A$ , we have  $t \in P_A$ , and so therefore  $s|_A \in P_A$  by compactness for  $A$ . Similarly,  $s|_B \in P_B$ , and therefore,  $s \in P_{A \otimes B}$ .

The proof for the linear implication is similar.  $\square$

Looking at the proof above, we may deduce the following important fact:

**Proposition 3.9.** *In the tensor product  $A \otimes B$ , only player  $O$  may switch games. In the linear implication  $A \multimap B$ , only player  $P$  may switch games.*

### 3.4 Composition of Strategies

Let  $A, B, C$  be games, let  $\sigma$  be a strategy for  $A \multimap B$  and let  $\tau$  be a strategy for  $B \multimap C$ . We shall construct a strategy  $\tau \circ \sigma$  for  $A \multimap C$ .

In order to define  $T_{\tau \circ \sigma}$ , we first define the set  $\sigma \parallel \tau$ , given by:

$$\sigma \parallel \tau = \left\{ s \in \overline{(M_A \sqcup M_B \sqcup M_C)^*} : s|_{A,B} \in \sigma, s|_{B,C} \in \tau \right\}$$

We then define

$$T_{\tau \circ \sigma} = \{\mathfrak{s}|_{A,C} : \mathfrak{s} \in \sigma \parallel \tau\}$$

We need to prove that  $T_{\tau \circ \sigma} \subseteq P_{A \multimap B}$ . If  $\mathfrak{s} \in \sigma \parallel \tau$ , then  $\mathfrak{s}|_{A,C}|_A = \mathfrak{s}|_A = \mathfrak{s}|_{A,B}|_A \in P_A$ , and similarly  $\mathfrak{s}|_{A,C} \in P_C$ , so certainly  $\mathfrak{s}|_{A,C} \in P_A \parallel P_C$ . It will suffice, then, to show that  $\mathfrak{s}|_{A,C}$  is alternating with respect to  $\zeta_{A \multimap C}$ .

**Proposition 3.10.** *Let  $A, B, C$  be games, let  $\sigma$  be a strategy for  $A \multimap B$  and let  $\tau$  be a strategy for  $B \multimap C$ . Suppose  $s, sa \in \tau \circ \sigma$ . Then  $\zeta_{A \multimap C}(s) = \neg \lambda_{A \multimap C}(a)$ .*

*Proof.* Write  $sa = \mathfrak{s}a|_{A,C}$ , where  $\mathfrak{s}a \in \sigma \parallel \tau$ . Then  $\mathfrak{s}|_{A,C} = s$ .  $s, sa$  must be finite, so, by Lemma 3.8, neither  $\mathfrak{s}|_{A,B}$  nor  $\mathfrak{s}|_{B,C}$  may have sign profile  $OP$ , so we get four possible sign profiles for  $\mathfrak{s}$  giving us the following values for  $\zeta_{A \multimap C}(\mathfrak{s}|_{A,C})$ :

| $\zeta_A(\mathfrak{s} _A)$ | $\zeta_B(\mathfrak{s} _B)$ | $\zeta_C(\mathfrak{s} _C)$ | $\zeta_{A \multimap C}(s _{A,C})$ |
|----------------------------|----------------------------|----------------------------|-----------------------------------|
| $P$                        | $P$                        | $P$                        | $P$                               |
| $P$                        | $P$                        | $O$                        | $O$                               |
| $P$                        | $O$                        | $O$                        | $O$                               |
| $O$                        | $O$                        | $O$                        | $P$                               |

Suppose that  $a$  is an  $O$ -move in  $A \multimap C$ : so  $a$  is a  $P$ -move in  $A$  or an  $O$ -move in  $C$ . By alternation in the component games, playing the move  $a$  either turns an  $O$ -position in  $A$  into a  $P$ -position in  $A$  or turns a  $P$ -position in  $C$  into an  $O$ -position in  $C$ . That is, we must be switching from sign profile  $PPP$  to sign profile  $PPO$  or from sign profile  $OOO$  to sign profile  $POO$ . In either case,  $s = \mathfrak{s}|_{A,C}$  must be a  $P$ -position.

Similarly, if  $a$  is a  $P$ -move in  $A \multimap C$ , then switching from  $\mathfrak{s}$  to  $\mathfrak{s}a$  must move us from sign profile  $PPO$  to sign profile  $PPP$  or from sign profile  $POO$  to sign profile  $OOO$ ; in either case,  $s$  must be an  $O$ -position.  $\square$

To complete the proof that  $T_{\tau \circ \sigma}$  is a legal trace set for a strategy, we need to show that  $T_{\tau \circ \sigma}$  is closed under opponent replies.

**Proposition 3.11.** *Let  $A, B, C$  be games, let  $\sigma$  be a strategy for  $A \multimap B$  and let  $\tau$  be a strategy for  $B \multimap C$ . Suppose  $s \in \tau \circ \sigma$  is a  $P$ -position such that  $sa \in P_{A \multimap C}$  for some  $O$ -move  $a$ . Then  $sa \in \tau \circ \sigma$ .*

*Proof.* Write  $s = \mathfrak{s}|_{A,C}$ , for  $\mathfrak{s} \in \sigma \parallel \tau$ . We show that  $\mathfrak{s}a \in \sigma \parallel \tau$ : so  $sa = \mathfrak{s}a|_{A,C} \in \tau \circ \sigma$ .

Since  $s$  is a  $P$ -position, the sign profile of  $\mathfrak{s}$  must be  $PPP$  or  $OOO$ . It follows that  $\mathfrak{s}|_{A,B} \in \sigma$  and  $\mathfrak{s}|_{B,C} \in \tau$  are both  $P$ -positions. Since  $a$  is a  $P$ -move in  $A \multimap C$ , it must be either a  $P$ -move in  $A \multimap B$  or a  $P$ -move in  $B \multimap C$ , by straightforward analysis of the definitions of the  $\lambda$  functions. It follows that either  $s|_{A,B}a \in \sigma$  (if  $a$  is a move in  $A$ ) or that  $s|_{B,C}a \in \tau$  (if  $a$  is a move in  $C$ ). In either case, we have  $\mathfrak{s}a \in \sigma \parallel \tau$ , as desired.  $\square$

Before we consider the definition of  $D_{\tau \circ \sigma}$ , we show that composition is associative at the level of trace sets.

**Proposition 3.12.** *Let  $A, B, C$  be games, let  $\sigma$  be a strategy for  $A \multimap B$ , let  $\tau$  be a strategy for  $B \multimap C$  and let  $v$  be a strategy for  $C \multimap D$ . Then*

$$T_{v \circ (\tau \circ \sigma)} = T_{(v \circ \tau) \circ \sigma}$$

*Proof.* As in [AJ13]. Define

$$\sigma \parallel \tau \parallel v = \left\{ \mathfrak{s} \in \overline{(M_A \sqcup M_B \sqcup M_C \sqcup M_D)^*} : \mathfrak{s}|_{A,B} \in \sigma, \mathfrak{s}|_{B,C} \in \tau, \mathfrak{s}|_{C,D} \in v \right\}$$

We shall show that

$$T_{v \circ (\tau \circ \sigma)} = \{ \mathfrak{s}|_{A,D} : \mathfrak{s} \in \sigma \parallel \tau \parallel v \}$$

The same claim will follow for  $T_{(v \circ \tau) \circ \sigma}$  by symmetry, completing the proof.

Let  $\mathfrak{s} \in \sigma \parallel \tau \parallel v$ . Then  $\mathfrak{s}|_{A,B,C} \in \sigma \parallel \tau$  and so  $\mathfrak{s}|_{A,C} \in \tau \circ \sigma$ . Now we see that  $\mathfrak{s}|_{A,C,D} \in (\tau \circ \sigma) \parallel v$  and so  $\mathfrak{s}|_{A,D} \in v \circ (\tau \circ \sigma)$ . We have shown that

$$\{ \mathfrak{s}|_{A,D} : \mathfrak{s} \in \sigma \parallel \tau \parallel v \} \subseteq T_{v \circ (\tau \circ \sigma)}$$

To show the reverse inclusion, suppose that  $s \in v \circ (\tau \circ \sigma)$ . Let  $\mathfrak{s} \in (\tau \circ \sigma) \parallel v$  be such that  $\mathfrak{s}|_{A,D} = s$  and let  $\mathfrak{t} \in \sigma \parallel \tau$  be such that  $\mathfrak{t}|_{A,C} = \mathfrak{s}|_{A,C}$ . Our goal is to interleave the sequences  $\mathfrak{s}$  and  $\mathfrak{t}$  in order to give rise to a sequence  $\mathfrak{u} \in \sigma \parallel \tau \parallel v$  such that  $\mathfrak{u}|_{A,D} = s$ .

Note that since  $\mathfrak{s}|_{A,C} = \mathfrak{t}|_{A,C}$ , we may decompose  $\mathfrak{s}$  and  $\mathfrak{t}$  into blocks:

$$\begin{aligned} \mathfrak{s} &= \mathfrak{s}_0 D_0 \mathfrak{s}_1 D_1 \dots \\ \mathfrak{t} &= \mathfrak{s}_0 B_0 \mathfrak{s}_1 B_1 \dots \end{aligned}$$

where  $\mathfrak{s}_1, \mathfrak{s}_2, \dots$  are sequences of moves from either  $A$  or  $C$ ,  $B_0, B_1 \dots$  are (possibly empty) sequences of moves from  $B$  and  $D_0, D_1, \dots$  are (possibly empty) sequences of moves from  $D$ .

We then form the sequence  $\mathfrak{u}$  as:

$$\mathfrak{u} = \mathfrak{s}_0 B_0 D_0 \mathfrak{s}_1 B_1 D_1 \dots$$

It might look as if we have made an arbitrary choice in placing the  $B_i$  before the  $D_i$ , but in fact it turns out that there is no choice to make: an examination of sign profiles, for example, will show that  $B_i$  is non-empty only if  $D_i$  is empty and vice versa: if  $\mathfrak{s}_i$  can be followed by a move from  $B$  then it cannot be followed by a move from  $D$ .

In any case, we have

$$\begin{aligned} \mathfrak{u}|_{A,C,D} &= \mathfrak{s}_0 D_0 \mathfrak{s}_1 D_1 \dots = \mathfrak{s} \\ \mathfrak{u}|_{A,B,C} &= \mathfrak{s}_0 B_0 \mathfrak{s}_1 B_1 \dots = \mathfrak{t} \end{aligned}$$



It follows that  $u \in \sigma \parallel \tau \parallel v$ :

$$\begin{aligned} u|_{A,B} &= t|_{A,B} \in \sigma \\ u|_{B,C} &= t|_{B,C} \in \tau \\ u|_{C,D} &= s|_{C,D} \in v \end{aligned}$$

Lastly, we have  $u|_{A,D} = s|_{A,D} = s$ , completing the proof.  $\square$

At the moment, we have an associative composition at the level of trace sets. We now define composition at the level of divergences.

Given a game  $A$  and a subset  $S \subseteq P_A$ , we write  $\text{pocl}_A(S)$  for the postfix closure of  $S$  in  $P_A$ ; i.e.:

$$\text{pocl}_A(S) = \{s \in P_A : t \sqsubseteq s \text{ for some } t \in S\}$$

**Definition 3.13.** Let  $A, B, C$  be games, let  $\sigma$  be a strategy for  $A \multimap B$  and let  $\tau$  be a strategy for  $B \multimap C$ . We define the divergences of  $\tau \circ \sigma$  as follows: first we define

$$\sigma \not\prec \tau = \left\{ s \in \overline{(M_A \sqcup M_B \sqcup M_C)}^* \mid \begin{array}{l} \text{either } s|_{A,B} \in D_\sigma \text{ and } s|_{B,C} \in T_\tau \\ \text{or } s|_{A,B} \in T_\sigma \text{ and } s|_{B,C} \in D_\tau \end{array} \right\}$$

$$D_{\tau \circ \sigma} = \text{pocl}_{A \multimap C}(\{\mathfrak{s}|_{A,C} : \mathfrak{s} \in \sigma \not\prec \tau\})$$

**Proposition 3.14.**  $D_{\tau \circ \sigma}$  is a well-defined divergence set.

*Proof.*  $D_{\tau \circ \sigma}$  alternating by the same argument we used for  $T_{\tau \circ \sigma}$ , and is postfix-closed by definition. Now let  $s \in D_{\tau \circ \sigma}$  and suppose that  $\mathfrak{s}|_{A,C} \sqsubseteq s$ , where  $\mathfrak{s} \in \sigma \not\prec \tau$ . We need to show that there is some  $O$ -position  $t \sqsubseteq s$  such that  $t \in T_{\tau \circ \sigma} \cap D_{\tau \circ \sigma}$ .

Suppose first that  $\mathfrak{s}|_{A,B} \in D_\sigma$  and  $\mathfrak{s}|_{B,C} \in T_\tau$ . Then there is some  $O$ -position  $r \sqsubseteq \mathfrak{s}|_{A,B}$  such that  $r \in T_\sigma \cap D_\sigma$ . Let  $\mathfrak{t} \sqsubseteq \mathfrak{s}$  be some prefix such that  $\mathfrak{t}|_{A,B} = r$ . Then  $\mathfrak{t} \in \sigma \parallel \tau \cap \sigma \not\prec \tau$  and so  $t := \mathfrak{t}|_{A,C} \in T_{\tau \circ \sigma} \cap D_{\tau \circ \sigma}$ . Certainly  $t \sqsubseteq s$ , and  $t$  is an  $O$ -position since  $r$  is, by examination of sign profiles.

The case where  $\mathfrak{s}|_{A,B} \in T_\sigma$  and  $\mathfrak{s}|_{B,C} \in D_\tau$  is similar.  $\square$

We need to show that this composition satisfies the diverge-or-reply and local compactness rules.

In order to prove that the composition of locally compact strategies is locally compact, we are going to need some facts about deterministic strategies.

**Definition 3.15.** We say a strategy  $\sigma$  is *deterministic* if whenever  $s \in \sigma$  is an  $O$ -position and  $a, b$  are moves such that  $sa, sb \in \sigma$ , then  $a = b$ .

Deterministic strategies have the following nice technical property:

**Proposition 3.16.** *Let  $A, B, C$  be games, let  $\sigma$  be a deterministic strategy for  $A \multimap B$  and let  $\tau$  be a deterministic strategy for  $B \multimap C$ . Then  $\tau \circ \sigma$  is a deterministic strategy. Moreover, for any  $s \in \tau \circ \sigma$ , there is a unique minimal  $\mathfrak{s} \in \sigma \parallel \tau$  such that  $\mathfrak{s}|_{A,C} = s$ .*

*Proof.* The idea behind the proof is that determinism of  $\sigma$  and  $\tau$  mean that the play in the intermediate game  $B$  is completely determined (since every move played in  $B$  is either a  $P$ -move in  $A \multimap B$  or a  $P$ -move in  $B \multimap C$ ). For full details, see [AJ13].  $\square$

We can use this to prove the following:

**Proposition 3.17.** *Let  $A, B, C$  be games and let  $\sigma: A \multimap B$  and  $\tau: B \multimap C$  be compact deterministic strategies. Then  $\tau \circ \sigma$  is compact.*

*Proof.* Let  $s \in M_{A \multimap C}^\omega$  be an infinite sequence such that  $t \in \tau \circ \sigma$  for all finite prefixes  $t \sqsubset s$ . For each finite prefix  $t$  of  $s$ , there is a unique minimal sequence  $\mathfrak{t} \in \sigma \parallel \tau$  such that  $\mathfrak{t}|_{A,C} = t$ . Writing  $s$  as the limit of its finite prefixes

$$t_1 \sqsubseteq t_2 \sqsubseteq t_3 \sqsubseteq \dots$$

we build up a sequence

$$\mathfrak{t}_1 \sqsubseteq \mathfrak{t}_2 \sqsubseteq \mathfrak{t}_3 \sqsubseteq \dots$$

where  $\mathfrak{t}_i$  is the unique minimal sequence in  $\sigma \parallel \tau$  such that  $\mathfrak{t}_i|_{A,C} = t_i$ . We have  $\mathfrak{t}_i \sqsubseteq \mathfrak{t}_{i+1}$  because if we remove the last move from  $\mathfrak{t}_{i+1}$  then the resulting sequence  $\mathfrak{u}_i$  satisfies  $\mathfrak{u}_i|_{A,C} = t_i$  and therefore  $\mathfrak{t}_i \sqsubseteq \mathfrak{u}_i \sqsubseteq \mathfrak{t}_{i+1}$  by minimality of  $\mathfrak{t}_i$ .

Then the limit of the sequences  $\mathfrak{t}_i|_{A,B}$  must be contained in  $\sigma$ , since  $\sigma$  is compact, and similarly the limit of the sequences  $\mathfrak{t}_i|_{B,C}$  must be contained in  $\tau$ , since  $\tau$  is compact. It follows that the limit  $\mathfrak{t}$  of the sequences  $\mathfrak{t}_i$  is contained in  $\sigma \parallel \tau$ , and therefore that  $\mathfrak{t}|_{A,C} = s \in \tau \circ \sigma$ .  $\square$

Note that the composition of compact nondeterministic strategies need not be compact.

**Example 3.18.** Let the games  $I$  and  $\Sigma^*$  be defined as before, and let the game  $\Sigma$  be the game with a unique opponent starting move and a unique player reply:

$$\Sigma = (\{q, a\}, \{q \mapsto o, a \mapsto p\}, \{\epsilon \mapsto p, q \mapsto o, qa \mapsto p\}, \{\epsilon, q, qa\})$$

Let  $\mathbb{N}$  be the game where the opponent asks the player to choose a natural number:

$$\mathbb{N} = (\{q\} \cup \mathbb{N}, \{q \mapsto O, n \mapsto P\}, \{\epsilon \mapsto P, q \mapsto O, qn \mapsto P\}, \{\epsilon, q\} \cup \{qn : n \in \mathbb{N}\})$$

We have a (deterministic) strategy  $\sigma$  for  $\mathbb{N} \multimap (\Sigma^* \multimap \Sigma)$  in which the player asks the opponent for a number  $n$  and then calls the argument of  $\Sigma^*$   $n$  times

before returning. For example, a play in  $\sigma$  might look like this:

$$\begin{array}{c}
\mathbb{N} \quad \Sigma^* \quad \Sigma \\
\qquad \qquad \qquad q \\
q \\
2 \\
\qquad \qquad \qquad q \\
\qquad \qquad \qquad a \\
\qquad \qquad \qquad q \\
\qquad \qquad \qquad a \\
\qquad \qquad \qquad a
\end{array}$$

$\sigma$  is compact because there are no infinite chains of positions; indeed, the length of a play in  $\sigma$  is already determined by the third move.

We have a nondeterministic strategy  $?$  for  $\mathbb{N} \cong I \multimap \mathbb{N}$  where the player nondeterministically chooses any number - so  $? = P_{\mathbb{N}}$  as sets of positions.  $?$  is compact because all plays in  $?$  have length at most 2.

When we compose these strategies, the resulting strategy  $\sigma \circ ?$  contains the infinite chain of positions  $q, qqa, qqqa, qqqaqa, \dots$ , but does not contain any limiting plays. So it is not compact. It is, however, locally compact, since it is the union of compact strategies  $\sigma_n$ , where  $\sigma_n$  has the unique maximal play

$$q \underbrace{qa \dots qa}_{n \text{ times}} a$$

**Proposition 3.19.** *Let  $A, B, C$  be games and let  $\sigma: A \multimap B$  and  $\tau: B \multimap C$  be (nondeterministic, locally compact) strategies. Then  $\tau \circ \sigma$  is locally compact.*

*Proof.* Let  $s \in T_{\tau \circ \sigma}$ . Then there is some sequence  $\mathfrak{s} \in \sigma \parallel \tau$  such that  $\mathfrak{s}|_{A,C} = s$ . We have  $\mathfrak{s}|_{A,B} \in \sigma$  and  $\mathfrak{s}|_{B,C} \in \tau$ , so, by local compactness for  $\sigma$  and  $\tau$ , there are compact strategies  $\sigma': A \multimap B$  and  $\tau': B \multimap C$  such that  $\mathfrak{s}|_{A,B} \in \sigma'$  and  $\mathfrak{s}|_{B,C} \in \tau'$ . After removing positions from (and possibly adding divergences to)  $\sigma'$  and  $\tau'$ , we may assume that they are deterministic (and still compact). Then  $\tau' \circ \sigma'$  is compact and we have  $\mathfrak{s}|_{A,C} \in \tau' \circ \sigma' \subseteq \tau \circ \sigma$  and  $D_{\tau \circ \sigma} \subseteq D_{\tau' \circ \sigma'}$ .  $\square$

Unlike [HM99], we do not need to change the definition of  $D_{\tau \circ \sigma}$  to reflect divergence arising from infinite chattering. The following example shows that it arises naturally from existing definition of  $D_{\tau \circ \sigma}$ .

**Example 3.20** (Infinite chattering). Let  $\Sigma^*$  denote the game where each player has exactly one move available in every position and the resulting infinite position is a  $P$ -position:

$$\Sigma^* = (\{q, a\}, \{q \mapsto O, a \mapsto P\}, \{\epsilon \mapsto P, sq \mapsto O, sa \mapsto P, qaqa \dots \mapsto P\}, \{s \sqsubseteq qaqa \dots\})$$

Let  $I$  denote the empty game with no moves, and let  $\perp$  denote the game with a single opponent move and no player reply:

$$\begin{aligned} I &= (\emptyset, \emptyset, \{\epsilon \mapsto P\}, \{\epsilon\}) \\ \perp &= (\{*\}, \{*\mapsto O\}, \{\epsilon \mapsto P, *\mapsto O\}, \{\epsilon, *\}) \end{aligned}$$

There are natural strategies  $\sigma$  for  $I \multimap \Sigma^*$  and  $\tau$  for  $\Sigma^* \multimap \perp$  given at the level of traces by:

$$\begin{aligned} \sigma &= \{s \sqsubseteq qaqa \dots\} \\ \tau &= \{s \sqsubseteq *qaqa \dots\} \end{aligned}$$

Player  $P$  always has a reply to an  $O$ -position in  $\sigma$ , and we set  $D_\sigma = \emptyset$ . In  $\tau$ , the infinite sequence  $*qaqa \dots$  satisfies  $\zeta_{\Sigma^*}(*qaqa \dots |_{\Sigma^*}) = \zeta_{\Sigma^*}(qaqa \dots) = P$  and  $\zeta_\perp(*qaqa \dots |_\perp) = \zeta_\perp(*) = O$  and so it is an  $O$ -position. As in the remark at the end of the definition of a strategy, this  $O$ -position can have no  $P$ -reply, so it (or some prefix) must be designated divergent. We set

$$D_\tau = \{*qaqa \dots\}$$

Now, when we compose the strategies  $\sigma$  and  $\tau$ , at the level of traces we have

$$\tau \circ \sigma = \{\epsilon, *\}$$

The sequence  $*$  is the restriction to  $I, \perp$  of the infinite sequence

$$*qaqa \dots \in \sigma \parallel \tau$$

We now have that  $*qaqa \dots |_{\Sigma^*, \perp} \in D_\tau$  and therefore the sequence  $*$   $\in \tau \circ \sigma$  must be designated divergent. We have

$$D_{\tau \circ \sigma} = \{*\}$$

as desired.

Note that if we modify the definition of  $\Sigma^*$  so that the infinite position is an  $O$ -position, then we still get a divergence – this time, because the sequence  $*qaqa \dots |_{I, \sigma^*}$  must be contained in  $D_\sigma$ . If we leave out the infinite position from  $\sigma$  or  $\tau$  altogether, then that strategy is no longer locally compact, so it is not a valid strategy according to our definition.

This example is at the heart of the proof that the composition of strategies satisfies the diverge-or-reply rule. This is the point at which we need local compactness.

**Proposition 3.21.** *Let  $A, B, C$  be games, let  $\sigma$  be a strategy for  $A \multimap B$  and let  $\tau$  be a strategy for  $B \multimap C$ . Then  $T_{\tau \circ \sigma}, D_{\tau \circ \sigma}$  satisfy the diverge-or-reply rule.*

*Proof.* Suppose that  $s \in \tau \circ \sigma$  is an  $O$ -position and that  $s \notin D_{\tau \circ \sigma}$ . Let  $\mathfrak{s} \in \sigma \parallel \tau$  be such that  $\mathfrak{s}|_{A,C} = s$ . By local compactness of  $\sigma$  and  $\tau$ , there must be compact strategies  $\sigma' : A \multimap B$  and  $\tau' : B \multimap C$  such that  $\mathfrak{s}|_{A,C} \in \sigma'$  and  $\mathfrak{s}|_{B,C} \in \tau'$ .

Let  $\mathfrak{s}' \in \sigma' \parallel \tau'$  be maximal such that  $\mathfrak{s}'|_{A,C} = s$ . We know that such a maximal sequence must exist: if  $\mathfrak{s}_i$  are a chain of sequences from  $\sigma' \parallel \tau'$  such that  $\mathfrak{s}_i|_{A,C} = s$  then their limit  $\mathfrak{s}$  (which is contained in  $\sigma' \parallel \tau'$  by compactness of  $\sigma'$  and  $\tau'$ ) must satisfy  $\mathfrak{s}|_{A,C} = s$ . Since  $s \notin D_{\tau \circ \sigma}$ , we must have  $\mathfrak{s}|_{A,B} \notin D_\sigma \subseteq D_{\sigma'}$  and  $\mathfrak{s}|_{B,C} \notin D_\tau \subseteq D_{\tau'}$ .

Since  $s$  is an  $O$ -position, one of  $\mathfrak{s}|_{A,B}$  and  $\mathfrak{s}|_{B,C}$  must be an  $O$ -position (by transitivity of  $\Rightarrow$ ). Suppose without loss of generality that  $\mathfrak{s}|_{A,B}$  is an  $O$ -position. Since  $\mathfrak{s}|_{A,B} \notin D_{\sigma'}$ , there must be some move  $x$  such that  $\mathfrak{s}x|_{A,B} \in \sigma'$ , since  $\sigma'$  satisfies the diverge-or-reply rule. Then we have  $\mathfrak{s}x|_{A,B} \in \sigma'$ . Moreover, Lemma 3.8 tells us that  $\mathfrak{s}|_{B,C}$  must be a  $P$ -position, so  $\mathfrak{s}x|_{B,C} \in \tau'$ , since  $\tau'$  is closed under opponent replies. Therefore,  $\mathfrak{s}x \in \sigma' \parallel \tau'$ . By maximality of  $\mathfrak{s}$ ,  $x$  must be a move in  $A$  and therefore we have  $sx \in \tau' \circ \sigma' \subseteq \tau \circ \sigma$ .  $\square$

We now have a working definition of composition. It remains to prove that it is associative at the level of divergences:

**Proposition 3.22.** *Let  $A, B, C, D$  be games, let  $\sigma$  be a strategy for  $A \multimap B$ , let  $\tau$  be a strategy for  $B \multimap C$  and let  $v$  be a strategy for  $C \multimap D$ . Then*

$$D_{v \circ (\tau \circ \sigma)} = D_{(v \circ \tau) \circ \sigma}$$

*Proof.* Let

$$D = \left\{ \mathfrak{s} \in \overline{(M_A \sqcup M_B \sqcup M_C \sqcup M_D)^*} \mid \begin{array}{l} \text{either } \mathbf{1} \ \mathfrak{s}|_{A,B} \in D_\sigma, \mathfrak{s}|_{B,C} \in T_\tau \text{ and } \mathfrak{s}|_{C,D} \in T_v \\ \text{or } \mathbf{2} \ \mathfrak{s}|_{A,B} \in T_\sigma, \mathfrak{s}|_{B,C} \in D_\tau \text{ and } \mathfrak{s}|_{C,D} \in T_v \\ \text{or } \mathbf{3} \ \mathfrak{s}|_{A,B} \in T_\sigma, \mathfrak{s}|_{B,C} \in T_\tau \text{ and } \mathfrak{s}|_{C,D} \in D_v \end{array} \right\}$$

We claim that  $D_{v \circ (\tau \circ \sigma)} = \text{pocl}_{A \multimap D}(\{\mathfrak{s}|_{A,D} : \mathfrak{s} \in D\})$ . The same claim will hold for  $D_{(v \circ \tau) \circ \sigma}$  by symmetry, completing the proof.

Let  $\mathfrak{s} \in D$ . In cases **1** and **2**, we must have  $\mathfrak{s}|_{A,C} \in D_{\tau \circ \sigma}$  and  $\mathfrak{s}|_{C,D} \in T_v$ . In case **3**, we have  $\mathfrak{s}|_{C,D} \in D_v$  and  $\mathfrak{s}|_{A,C} \in T_{\tau \circ \sigma}$ . In either case, we must have  $\mathfrak{s}|_{A,D} \in D_{v \circ (\tau \circ \sigma)}$ .

Conversely, suppose that  $s \in D_{v \circ (\tau \circ \sigma)}$ . Then there must be some  $\mathfrak{t} \in \overline{(M_A \sqcup M_C \sqcup M_D)^*}$  such that  $\mathfrak{t}|_{A,D} \sqsubseteq s$  and either

$$\mathfrak{t}|_{A,C} \in D_{\tau \circ \sigma} \text{ and } \mathfrak{t}|_{C,D} \in T_v$$

or

$$\mathfrak{t}|_{A,C} \in T_{\tau \circ \sigma} \text{ and } \mathfrak{t}|_{C,D} \in D_v$$

In the first case, we then get a sequence  $\mathfrak{u} \in \overline{(M_A \sqcup M_B \sqcup M_C)^*}$  such that  $\mathfrak{u}|_{A,C} \sqsubseteq \mathfrak{t}|_{A,C}$  and either

$$\mathfrak{u}|_{A,B} \in D_\sigma \text{ and } \mathfrak{u}|_{B,C} \in T_\tau$$

or

$$\mathbf{u}|_{A,B} \in T_\sigma \text{ and } \mathbf{u}|_{B,C} \in D_\tau$$

By passing to a suitable prefix of  $\mathbf{t}$ , we may assume that  $\mathbf{u}|_{A,C} = \mathbf{t}|_{A,C}$  (since  $T_v$  is prefix-closed).

In the second case, we get a sequence  $\mathbf{u} \in \sigma \parallel \tau$  such that  $\mathbf{u}|_{A,C} = \mathbf{t}|_{A,C}$ .

By the argument from Proposition 3.12, we may interleave these two sequences  $\mathbf{t}$  and  $\mathbf{u}$  to form a sequence  $\mathbf{s} \in (M_A \sqcup M_B \sqcup M_C \sqcup M_D)^*$  such that  $\mathbf{s}|_{A,B,C} = \mathbf{u}$  and  $\mathbf{s}|_{A,C,D} = \mathbf{t}$ . Then it is clear that  $\mathbf{s} \in D$  in all cases and we have  $\mathbf{s}|_{A,D} = \mathbf{t}|_{A,D} \sqsubseteq s$ .  $\square$

In order to show that we have a category, it remains to define the identity morphisms. These are given by the usual copycat strategies:

$$\text{id}_A = \{s \in P_{A_1 \multimap A_2} : \text{for all even length } t \sqsubseteq s, t|_{A_1} = t|_{A_2}\}$$

(Here, we consider infinite positions to have even length).

The divergence set  $D_{\text{id}_A}$  is empty: note that infinite positions in  $\text{id}_A$  are always  $P$ -positions.

**Theorem 3.23.** *There is a category  $\mathcal{G}$  where the objects are games as defined above, morphisms from a game  $A$  to a game  $B$  are given by nondeterministic strategies  $(T_\sigma, D_\sigma)$  for  $A \multimap B$  and the composition and identity morphisms are as defined above.*

*Proof.* All that is left to do is to show that  $\text{id}_A$  really is an identity morphism. We shall show that if  $A$  and  $B$  are games and  $\sigma$  is a strategy for  $A \multimap B$  then  $\sigma \circ \text{id}_A = \sigma$ ; it will follow by a symmetrical argument that  $\text{id}_B \circ \sigma = \sigma$ .

We first show that  $T_{\sigma \circ \text{id}_A} = T_\sigma$ . Because of closure under opponent replies, it suffices to show that  $\sigma$  and  $\sigma \circ \text{id}_A$  contain the same  $P$ -positions. If  $s$  is a  $P$ -position in  $\sigma \circ \text{id}_A$ , then there is some sequence  $\mathbf{s} \in \text{id}_A \parallel \sigma$  such that  $\mathbf{s}|_{A_1,B} = s$ . Since  $\mathbf{s}|_{A_1,B}$  is a  $P$ -position, it must have sign profile  $PPP$  or  $OOO$  by Lemma 3.8, and therefore  $\mathbf{s}|_{A_1,A_2}$  is a  $P$ -position, so has even length. Therefore,  $\mathbf{s}|_{A_1} = \mathbf{s}|_{A_2}$  and so  $s = \mathbf{s}|_{A_1,B} = \mathbf{s}|_{A_2,B} \in \sigma$ .

Conversely, suppose that  $s \in \sigma$  is a  $P$ -position. There is an obvious sequence  $s^2 \in \text{id}_A$ , given by copying the moves of  $s|_A$  between copies of  $A$ , such that  $s^2|_{A_1} = s^2|_{A_2} = s|_A$ . Then  $s^2|_{A_2} = s|_{A_2}$ , and so the sequences  $s^2$  and  $s$  may be interleaved as in the proof of Proposition 3.12 to yield a sequence  $\mathbf{s} \in \text{id}_A \parallel \sigma$  such that  $\mathbf{s}|_{A_1,B} = \mathbf{s}|_{A_2,B} = s$ .

Now we show that  $D_{\sigma \circ \text{id}_A} = D_\sigma$ . Suppose  $s \in D_{\sigma \circ \text{id}_A}$ . Since  $D_{\text{id}_A} = \emptyset$ , there is a sequence  $\mathbf{s} \in (M_{A_1} \sqcup M_{A_2} \sqcup M_B)^*$  such that  $\mathbf{s}|_{A_1,B} \sqsubseteq s$ ,  $\mathbf{s}|_{A_1,B_1} \in \text{id}_A$  and  $\mathbf{s}|_{A_2,B} \in D_\sigma$ . In particular,  $\mathbf{s}|_{A_2,B}$  is an  $O$ -position in  $A \multimap B$  and so  $\mathbf{s}|_{A_2}$  is a  $P$ -position in  $A$ . It follows by Lemma 3.8 that  $\mathbf{s}|_{A_1,A_2}$  is a  $P$ -position in  $A \multimap A$ ;

hence, it is even-length and so  $\mathfrak{s}|_{A_1} = \mathfrak{s}|_{A_2}$ . It follows that  $\mathfrak{s}|_{A_1, B} = \mathfrak{s}|_{A_2, B} \in D_\sigma$ , so  $s \in D_\sigma$ , since  $D_\sigma$  is postfix-closed. Therefore,  $D_{\sigma \text{oid}_A} \subseteq D_\sigma$ .

Conversely, suppose that  $s \in D_\sigma$ . We know already that there is some  $\mathfrak{s} \in \text{id}_A \parallel \sigma$  such that  $\mathfrak{s}|_{A_1, A_2} \in \text{id}_A$  and  $\mathfrak{s}|_{A_2, B} \in \sigma$ . Then  $\mathfrak{s}|_{A, C} \in D_{\sigma \text{oid}_A}$  by definition.  $\square$

### 3.5 Symmetric monoidal closed structure

We make the category  $\mathcal{G}$  into a symmetric monoidal closed category with tensor product given by  $\otimes$  and linear implication given by  $\multimap$ . At the level of morphisms, the action of  $\_ \otimes \_$  is defined as follows: if  $A, B, C, D$  are games,  $\sigma$  is a strategy for  $A \multimap C$  and  $\tau$  is a strategy for  $B \multimap D$ , then  $\sigma \otimes \tau$  is the strategy for  $(A \otimes B) \multimap (C \otimes D)$  given by playing the strategy  $\sigma$  in  $A$  and  $C$  and playing the strategy  $\tau$  in  $B$  and  $D$ :

$$\sigma \otimes \tau = \{s \in P_{(A \otimes B) \multimap (C \otimes D)} : s|_{A, C} \in \sigma, s|_{B, D} \in \tau\}$$

The divergences for  $\sigma \otimes \tau$  are defined using the divergences for  $\sigma$  and  $\tau$ :

$$D_{\sigma \otimes \tau} = \text{pocl} \left( \left\{ s \in P_{(A \otimes B) \multimap (C \otimes D)} \mid \begin{array}{l} \text{either } s|_{A, C} \in D_\sigma \text{ and } s|_{B, D} \in T_\tau \\ \text{or } s|_{A, C} \in T_\sigma \text{ and } s|_{B, D} \in D_\tau \end{array} \right\} \right)$$

**Proposition 3.24.**  $\sigma \otimes \tau$  is a valid strategy for  $(A \otimes B) \multimap (C \otimes D)$

*Proof.*  $\sigma \otimes \tau$  is non-empty and prefix-closed, since  $\sigma$  and  $\tau$  are. To show that it is a valid trace-set, we need to show that it is closed under opponent replies.

Let  $s \in \sigma \otimes \tau$  be a  $P$ -position and suppose that  $sx \in P_{(A \otimes B) \multimap (C \otimes D)}$  for some move  $x$ . We start with some brief analysis of sign profiles.

By Lemma 3.8,  $s|_{A, B}$  and  $s|_{C, D}$  must either both be  $O$ -positions or both be  $P$ -positions. If they are  $P$ -positions, then the only possibility for the sign profile of  $s$  is:

$$\frac{\zeta_A(s|_A) \quad \zeta_B(s|_B) \quad \zeta_C(s|_C) \quad \zeta_D(s|_D)}{P \quad P \quad P \quad P}$$

Otherwise, if they are both  $O$ -positions, then each of  $s|_{A, B}$  and  $s|_{C, D}$  may either have sign profile  $OP$  or sign profile  $PO$ , by Lemma 3.8. But now, since  $s|_{A, C} \in \sigma \subseteq P_{A \multimap C}$  and since  $s|_{B, D} \in \tau \subseteq P_{B \multimap D}$ , we also see that neither  $s|_{A, C}$  nor  $s|_{B, D}$  may have sign profile  $OP$ . We therefore have two more possibilities for the sign profile of  $s$ :

$$\frac{\zeta_A(s|_A) \quad \zeta_B(s|_B) \quad \zeta_C(s|_C) \quad \zeta_D(s|_D)}{P \quad O \quad P \quad O}$$

$$\frac{\zeta_A(s|_A) \quad \zeta_B(s|_B) \quad \zeta_C(s|_C) \quad \zeta_D(s|_D)}{O \quad P \quad O \quad P}$$

In all three cases,  $s|_{A, C}$  and  $s|_{B, D}$  are both  $P$ -positions. Since  $\sigma$  and  $\tau$  are both closed under opponent replies, it follows that one of them must contain

the move  $x$  as a reply to the current position, and therefore that  $sx \in \sigma \otimes \tau$ , as desired.

$D_{\sigma \otimes \tau}$  is postfix-closed by definition. If  $s \in D_{\sigma \otimes \tau}$ , we need to find some  $O$ -position  $t \in T_{\sigma \otimes \tau} \cap D_{\sigma \otimes \tau}$  such that  $t \sqsubseteq s$ . By definition, we know there is some  $s' \sqsubseteq s$  such that either  $s'|_{A,C} \in D_\sigma$  and  $s'|_{B,D} \in T_\tau$  or  $s'|_{A,C} \in T_\sigma$  and  $s'|_{B,D} \in D_\tau$ . In the first case, we know that there is some  $r \sqsubseteq s'|_{A,C}$  such that  $r$  is an  $O$ -move in  $T_\sigma \cap D_\sigma$ ; passing to an appropriate prefix  $s''$  of  $s'$ , we get that  $s''|_{A,C} = r$  and  $s''|_{B,D} \in T_\tau$ . Therefore,  $s'' \in T_{\sigma \otimes \tau} \cap D_{\sigma \otimes \tau}$ . Moreover,  $s''$  is an  $O$ -position, since  $s''|_{A,C}$  is, and the claim follows since  $s'' \sqsubseteq s' \sqsubseteq s$ . The proof in the other case is similar.

Now we need to show that  $\sigma \otimes \tau$  is locally compact. We first show that if  $\sigma$  and  $\tau$  are compact strategies then  $\sigma \otimes \tau$  is compact. Indeed, suppose that  $s \in \sigma \otimes \tau$  is an infinite sequence such that  $t \in \sigma \otimes \tau$  for all finite prefixes  $t \sqsubset s$ . Then  $\sigma$  must contain all finite prefixes of  $s|_{A,C}$ , so it must contain  $s|_{A,C}$ , since  $\sigma$  is compact. Similarly,  $\tau$  must contain  $s|_{B,D}$ . Therefore,  $s \in \sigma \otimes \tau$ . Since  $s$  was arbitrary, it follows that  $\sigma \otimes \tau$  is compact.

Now suppose that  $\sigma$  and  $\tau$  are locally compact. Let  $s \in \sigma \otimes \tau$ . Then there exists a compact strategy  $\sigma'$  for  $A \multimap C$  and  $\tau': B \multimap D$  such that  $s|_{A,C} \in \sigma' \subseteq \sigma$  and  $D_\sigma \subseteq D_{\sigma'}$  and a compact strategy  $\tau'$  for  $B \multimap C$  such that  $s|_{B,D} \in \tau' \subseteq \tau$  and  $D_\tau \subseteq D_{\tau'}$ . Then  $s \in \sigma' \otimes \tau'$ , which is compact, and we have  $\sigma' \otimes \tau' \subseteq \sigma \otimes \tau$  and  $D_{\sigma \otimes \tau} \subseteq D_{\sigma' \otimes \tau'}$ .

Lastly, we need to show that  $\sigma \otimes \tau$  satisfies diverge-or-reply. Indeed, suppose that  $s \in \sigma \otimes \tau$  is an  $O$ -position and that  $s \notin D_{\sigma \otimes \tau}$ . Then  $s|_{A,C} \notin D_\sigma$  and  $s|_{B,D} \notin D_\tau$ . Using Lemma 3.8, we see that  $s$  must have one of the following sign profiles:

$$\begin{array}{cccc} \zeta_A(s|_A) & \zeta_B(s|_B) & \zeta_C(s|_C) & \zeta_D(s|_D) \\ \hline P & P & P & O \\ P & P & O & P \end{array}$$

It follows that one of  $s|_{A,C}$  and  $s|_{B,D}$  must be an  $O$ -position. Suppose without loss of generality that  $s|_{A,C}$  is an  $O$ -position. Since  $s|_{A,C} \notin D_\sigma$ , there must be some move  $x$  such that  $sx \in \sigma$ , since  $\sigma$  satisfies diverge-or-reply. Then we have  $sx \in \sigma \otimes \tau$ .  $\square$

We also need to show that the tensor product respects composition. This is easy.

**Proposition 3.25.** *Let  $A'', A', A, B'', B', B$  be games and let  $\sigma': A'' \rightarrow A'$ ,  $\sigma: A' \rightarrow A$ ,  $\tau': B'' \rightarrow B'$ ,  $\tau: B' \rightarrow B$  be strategies. Then*

$$(\sigma \otimes \tau) \circ (\sigma' \otimes \tau') = (\sigma \circ \sigma') \otimes (\tau \circ \tau')$$

*Proof.* At the level of trace sets, we have:

$$(\sigma \circ \sigma') \otimes (\tau \circ \tau') = \{s \in P_{(A'' \otimes B'') \multimap (A \otimes B)} : s|_{A'', A} \in \sigma \circ \sigma', s|_{B'', B} \in \tau \circ \tau'\}$$



Meanwhile:

$$\begin{aligned}
(\sigma' \otimes \tau') \parallel (\sigma \otimes \tau) &= \left\{ \mathfrak{s} \in \overline{(M_{A'' \otimes B''} \sqcup M_{A' \otimes B'} \sqcup M_{A \otimes B})^*} \mid \begin{array}{l} \mathfrak{s}|_{A'' \otimes B'', A' \otimes B'} \in \sigma' \otimes \tau' \\ \mathfrak{s}|_{A' \otimes B', A \otimes B} \in \sigma \otimes \tau \end{array} \right\} \\
&= \left\{ \mathfrak{s} \in \overline{(M_{A'' \otimes B''} \sqcup M_{A' \otimes B'} \sqcup M_{A \otimes B})^*} \mid \begin{array}{l} \mathfrak{s}|_{A'', A'} \in \sigma' \\ \mathfrak{s}|_{A', A} \in \sigma \\ \mathfrak{s}|_{B'', B'} \in \tau' \\ \mathfrak{s}|_{B', B} \in \tau \end{array} \right\} \\
&= \left\{ \mathfrak{s} \in \overline{(M_{A'' \otimes B''} \sqcup M_{A' \otimes B'} \sqcup M_{A \otimes B})^*} \mid \begin{array}{l} \mathfrak{s}|_{A'', A', A} \in \sigma' \parallel \sigma \\ \mathfrak{s}|_{B'', B', B} \in \tau' \parallel \tau \end{array} \right\}
\end{aligned}$$

It follows that

$$(\sigma \otimes \tau) \circ (\sigma' \otimes \tau') = \{\mathfrak{s}|_{A'' \otimes B'', A \otimes B} : \mathfrak{s} \in (\sigma' \otimes \tau') \parallel (\sigma \otimes \tau)\} = (\sigma \circ \sigma') \otimes (\tau \circ \tau')$$

We can show that

$$D_{(\sigma \otimes \tau) \circ (\sigma' \otimes \tau')} = D_{(\sigma \circ \sigma') \otimes (\tau \circ \tau')}$$

using a similar argument.  $\square$

The associators, unitors and braiding for this tensor product are the usual (deterministic, compact, non-divergent) copycat strategies. It is straightforward to show that they are natural transformations, using the same argument we used to show that  $\text{id}_A$  is an identity, and that the appropriate coherence diagrams are satisfied.

Closedness follows immediately from the structural isomorphism

$$(A \otimes B) \multimap C \cong A \multimap (B \multimap C)$$

for all games  $A, B, C$ , together with the definition of morphisms as strategies on the linear implication.

### 3.6 Exponentials

Let  $A$  be a game. We define a game  $!A$  in which the opponent may at any point open a new copy of the game  $A$ :

$$!A = (M_{!A}, \lambda_{!A}, \zeta_{!A}, P_{!A})$$

where

- $M_{!A} = \omega \times M_A$
- $\lambda_{!A} = \lambda_A \circ \text{pr}_2$

Given some sequence  $s \in \overline{M_{!A}^*}$ , we write  $s|_n$  for the projection on to the second component of the subsequence of  $s$  consisting of those moves of the form  $(n, a)$ . We define

$$!P_A = \left\{ s \in \overline{M_{!A}^*} : s|_n \in P_A \text{ for all } n \in \omega \right\}$$

We define  $\zeta_{!A} : !P_A \rightarrow \{O, P\}$  by

$$\zeta_{!A}(s) = \bigwedge_{n \in \omega} \zeta_A(s|_n)$$

Lastly, we define  $P_{!A}$  to be the subset of  $!P_A$  consisting of all those terms that are alternating with respect to  $\zeta_{!A}$  and that satisfy the following criterion on the order in which copies of  $A$  may be opened:

**Ordering criterion** If the move  $(n+1, a)$  occurs in  $s$ , for  $n \geq 0$ , then there must be some move  $b$  such that  $(n, b)$  occurs earlier in the sequence  $s$  than  $(n+1, a)$ .

This criterion says that player  $O$  must open the copies of  $A$  in sequence: he is not allowed to play in game  $n+1$  till he has made a move in game  $n$ .

**Proposition 3.26.**  *$!A$  is a well formed game.*

*Proof.* Exactly the same argument as in Proposition 3.7. We need to prove a version of Lemma 3.8 which will tell us that if  $s \in P_{!A}$  then either  $\zeta_A(s|_n) = O$  for at most one value of  $n$ .  $\square$

Just like the tensor product, the exponential  $!A$  has the property that only player  $O$  may switch games.

The exponential connective gives rise to a functor  $\mathcal{G} \rightarrow \mathcal{G}$ , where the operation on morphisms  $\sigma : A \multimap B$  is given by:

$$!\sigma = \{s \in P_{!A \multimap !B} : s|_{A_n, B_n} \in \sigma \text{ for all } n \in \omega\}$$

with divergences

$$D_{!\sigma} = \text{pocl}_{!A} \left( \left\{ s \in !\sigma \mid \begin{array}{l} \text{there exists } n \in \omega \text{ such that } s|_n \in D_\sigma \\ \text{and for all } m \neq n, s|_m \in T_\sigma \end{array} \right\} \right)$$

The proof that  $!(\cdot)$  preserves composition is the same as for the tensor product.

$!(\cdot)$  gives rise to a *linear exponential comonad* on  $\mathcal{G}$  (see [Hyl97], for example). The relevant natural transformations, including the comultiplication  $\text{mult} : !A \rightarrow !!A$ , which opens a copy of  $A$  on the left for every copy of  $A$  opened on the right, are all deterministic, compact, non-divergent strategies. We only have to show that they are natural transformations. For illustration, we shall do this in the case of the comultiplication.

**Proposition 3.27.** *Let  $A, B$  be games, and let  $\sigma$  be a strategy for  $A \multimap B$ . Then the following diagram commutes:*

$$\begin{array}{ccc} !A & \xrightarrow{\text{mult}} & !!A \\ !\sigma \downarrow & & \downarrow !!\sigma \\ !B & \xrightarrow{\text{mult}} & !!B \end{array}$$

*Proof.* Both branches of the diagram are strategies that play as  $\sigma$  in each copy of  $B$  in  $!!B$ , opening a copy of  $A$  in  $!A$  as required. The ordering condition means that there is only one possible order in which to open the copies of  $A$  in  $!A$ , so both strategies must behave the same way.

A divergence occurs in either branch if and only if a divergence arises in  $\sigma$  in one of the  $A - B$  pairs. So both branches have the same divergence sets.  $\square$

We may now form the co-Kleisli category  $\mathcal{G}^!$ . The objects of  $\mathcal{G}^!$  are games  $A$  and a morphism from  $A$  to  $B$  is given by a strategy for

$$A \Longrightarrow B := !A \multimap B$$

We compose two morphisms  $\sigma: A \Longrightarrow B$  and  $\tau: B \Longrightarrow C$  by setting:

$$\tau \circ \sigma = !A \xrightarrow{\text{mult}} !!A \xrightarrow{!\sigma} !B \xrightarrow{\tau} C$$

where composition on the right is in the category  $\mathcal{G}$ .

A standard property of the linear exponential comonad is:

**Theorem 3.28.** *The category  $\mathcal{G}^!$  is Cartesian closed with  $\otimes$  as the Cartesian product and  $\Longrightarrow$  as the internal Hom.*

## 4 Denotational Semantics for Fair PCF

Our base PCF type  $\mathbb{N}$  will be denoted by the game  $\mathbb{N}$  defined above, while the product and function types will be interpreted by the connectives  $\otimes$  and  $\Longrightarrow$ . It remains to give the denotation of each of the terms of Fair PCF as a strategy for the appropriate game.

The numerals  $\underline{n}$  will each be interpreted as the strategy for  $\mathbb{N}$  that always returns the value  $n$ . The nondeterministic oracle  $?$  will be interpreted as the strategy  $?$  given at the level of trace sets by  $? = P_{\mathbb{N}}$ . None of these strategies have any divergent positions.

The other PCF terms all have obvious strategies that they correspond to. The only difficult one is the combinator  $\mathbf{Y}_T$ .

As usual, we shall use fixed point theorems from domain theory to define the denotation of  $\mathbf{Y}$ . The main ideas come from [Lai15].

## 4.1 The stable order on strategies

We start by defining some orders on strategies. Our main ordering, which we shall call the *stable ordering* (following [Lai15]) is the order given in [HM99].

**Definition 4.1** (Extensional order). Let  $A$  be a game and let  $\sigma, \tau$  be strategies for  $A$ . We say that  $\sigma \sqsubseteq \tau$  if:

- 1)  $T_\sigma \subseteq T_\tau$
- 2)  $D_\sigma \subseteq D_\tau$

The extensional order is simple to define, but it is not preserved by composition (see [HM99] for an example). Instead, we shall prefer the following order:

**Definition 4.2** (Stable order). Let  $A$  be a game and let  $\sigma, \tau$  be strategies for  $A$ . We say that  $\sigma \leq_s \tau$  if:

- 1)  $T_\sigma \subseteq T_\tau$
- 2)  $D_\tau \subseteq D_\sigma$
- 3)  $T_\tau \subseteq T_\sigma \cup D_\sigma$

To show that this is an order, the only non-trivial part is to show that property (3) is transitive. Suppose that  $\sigma \leq_s \tau \leq_s \nu$  are strategies and that  $s \in T_\nu \setminus T_\tau$ . If  $s \in T_\tau$ , then there exists  $s \in D_\sigma$ , since  $\sigma \leq_s \tau$  and if  $s \notin T_\tau$ , then  $s \in D_\tau \subseteq D_\sigma$ , since  $\tau \leq_s \nu$ . Therefore,  $\sigma \leq_s \nu$ .

**Proposition 4.3.** Let  $A, B, C$  be games, let  $\sigma, \sigma'$  be strategies for  $A \multimap B$  and let  $\tau, \tau'$  be strategies for  $B \multimap C$ . Suppose that  $\sigma \leq_s \sigma'$  and  $\tau \leq_s \tau'$ . Then:

$$\tau \circ \sigma \leq_s \tau' \circ \sigma'$$

*Proof.* First suppose that  $\tau = \tau'$ .

Since  $T_\sigma \subseteq T_{\sigma'}$ , we have  $\sigma \parallel \tau \subseteq \sigma' \parallel \tau$  and therefore  $T_{\tau \circ \sigma} \subseteq T_{\tau \circ \sigma'}$ .

Now let  $e \in D_{\tau \circ \sigma'}$ . Suppose  $e \sqsubseteq \mathfrak{e}$ , where  $\mathfrak{e} \in \sigma' \not\leq \tau$  and suppose first that  $\mathfrak{e}|_{A,B} \in D_{\sigma'}$  and  $\mathfrak{e}|_{B,C} \in T_\tau$ . Since  $D_{\sigma'} \subseteq D_\sigma$ , we have  $\mathfrak{e}|_{A,B} \in D_\sigma$ , so  $\mathfrak{e} \in \sigma \not\leq \tau$  and therefore  $e$ , which is a postfix of  $\mathfrak{e}|_{A,C}$ , is contained in  $D_{\tau \circ \sigma}$ .

Now suppose that  $\mathfrak{e}|_{A,B} \in T_{\sigma'}$  and  $\mathfrak{e}|_{B,C} \in D_{\tau'}$ . Since  $\sigma \leq_s \sigma'$ , we have  $\mathfrak{e}|_{A,B} \in T_\sigma \cup D_\sigma$ . If  $\mathfrak{e}|_{B,C} \in T_\sigma$ , then we have  $\mathfrak{e} \in \sigma \not\leq \tau$  and so  $\mathfrak{e}|_{A,C} \in D_{\tau \circ \sigma}$  and therefore  $e \in D_{\tau \circ \sigma}$ , since  $D_{\tau \circ \sigma}$  is postfix-closed.

Otherwise,  $\mathfrak{e}|_{B,C} \in D_\sigma$ . Then  $\mathfrak{e}|_{A,B}$  and  $\mathfrak{e}|_{B,C}$  have prefixes  $q \sqsubseteq \mathfrak{e}|_{A,B}$ ,  $r \sqsubseteq \mathfrak{e}|_{B,C}$  such that  $q$  and  $r$  are  $O$ -moves,  $q \in T_\sigma \cap D_\sigma$  and  $r \in T_\tau \cap D_\tau$ . Passing to an appropriate prefix  $\mathfrak{e}'$  of  $\mathfrak{e}$ , we have either  $\mathfrak{e}'|_{A,B} = q$  and  $r \sqsubseteq \mathfrak{e}'|_{B,C}$  or  $q \sqsubseteq \mathfrak{e}'|_{A,B}$  and  $\mathfrak{e}'|_{B,C} = r$ . In either case,  $\mathfrak{e}' \in \sigma \not\leq \tau$  and so  $\mathfrak{e}'|_{A,C} \in D_{\tau \circ \sigma}$ . Then we have  $\mathfrak{e}'|_{A,C} \sqsubseteq \mathfrak{e}|_{A,C} \sqsubseteq e$ , and so  $e \in D_{\tau \circ \sigma}$ .

Lastly, suppose that  $s \in T_{\tau' \circ \sigma'} \setminus T_{\tau \circ \sigma}$ . Write  $s = \mathfrak{s}$ , where  $\mathfrak{s} \in \sigma' \parallel \tau'$ . Since  $\mathfrak{s} \notin T_{\tau \circ \sigma}$ , we must have  $\mathfrak{s}|_{A,B} \in \sigma' \setminus \sigma$ . It follows that  $\mathfrak{s}|_{A,B} \in D_\sigma$  and therefore  $\mathfrak{s} \in \sigma \not\leq \tau$ . Therefore,  $s = \mathfrak{s}|_{A,B} \in D_{\tau \circ \sigma}$ .

The result follows by a similar argument in the case that  $\sigma = \sigma'$ . Therefore, the full result follows from transitivity of  $\leq_s$ : if  $\sigma \leq_s \sigma'$  and  $\tau \leq_s \tau'$ , then

$$\tau \circ \sigma \leq_s \tau \circ \sigma' \leq_s \tau' \circ \sigma'$$

□

We want to get a similar result for composition in the category  $\mathcal{G}^!$ . To do that it will suffice to prove the following result:

**Proposition 4.4.** *Let  $A, B$  be games and let  $\sigma, \tau$  be strategies for  $A \multimap B$  such that  $\sigma \leq_s \tau$ . Then  $!\sigma \leq_s !\tau$  as strategies for  $!A \multimap !B$ .*

*Proof.* If  $s \in !\sigma$ , then  $s|_{A_n, B_n} \in \sigma$  for all  $n$  and therefore  $s|_{A_n, B_n} \in \sigma'$  for all  $n$ . It follows that  $!\sigma \subseteq !\tau$ .

If  $e \in D_{!\tau}$  then there exists some  $e' \sqsubseteq e$  and some  $n \in \omega$  such that  $e'|_n \in D_\tau$  and  $e'|_m \in T_\tau$  for all  $m \neq n$ .

Since  $\sigma \leq_s \tau$ , we have  $D_\tau \subseteq D_\sigma$  and  $T_\tau \subseteq T_\sigma \cup D_\sigma$ . For each  $m$ , define a subsequence  $d_n$  of  $e'$  as follows: if  $e'|_m \in T_\sigma$  then  $d_m = e'|_m$ . If  $e'|_m \in D_\sigma$ , then there is some prefix of  $e'|_m$  that is an  $O$ -position in  $T_\sigma \cap D_\sigma$ . Let  $d_m$  be the shortest such prefix of  $e'|_m$ .

We know that  $e'|_n \in D_\tau \subseteq D_\sigma$ , so  $d_n \in D_\sigma$ . Let  $d$  be the shortest prefix of  $e'$  such that there exists some  $m$  such that  $d_m \in D_\sigma$  and  $d_m \sqsubseteq d$ . We claim that  $d|_k \in T_\sigma$  for all  $k \neq n$ . If  $d|_k \sqsubseteq d_k$ , then  $d|_k \in T_\sigma$ , since  $d_k \in T_\sigma$ . Otherwise,  $d_k \not\sqsubseteq d|_k$ . But this contradicts the minimality of  $d$ .

Therefore, we have  $d \in D_{!\sigma}$ . It follows that  $e \in D_{!\sigma}$ , since  $D_{!\sigma}$  is postfix-closed.

Lastly, if  $s \in !\tau \setminus !\sigma$ , then there exists some  $s' \sqsubseteq s$  and some  $n \in \omega$  such that  $s'|_{A_n, B_n} \in \tau \setminus \sigma$ . Therefore, if we construct the  $d_n$  as before, we have  $d_n \in D_\sigma$  for at least one  $n$  and it follows that we may construct  $d \in D_{!\sigma}$  as before such that  $d \sqsubseteq s'$ . It follows that  $s \in D_{!\sigma}$ . □

If  $A$  is a game and  $S \subseteq P_A$  is some set, we say that  $s \in S$  is *minimal* if  $t \not\sqsubseteq s$  for all proper prefixes  $t \sqsubset s$ . Let  $A$  be a game and let  $F$  be a non-empty set of strategies for  $A$  that is directed under the extensional order: so if  $\sigma, \tau \in F$  then there is some  $v \in F$  such that  $\sigma, \tau \sqsubseteq v$ . Write  $D(F) = \bigcap_{\sigma \in F} D_\sigma$ . We define a strategy  $\bigsqcup F$  for  $F$  by setting

$$\begin{aligned} T_{\bigsqcup F} &= \bigcup_{\sigma \in F} T_\sigma \cup \{s \in D(F) : s \text{ is an infinite and minimal } O\text{-position in } D(F)\} \\ D_{\bigsqcup F} &= D(F) \setminus \{s \in D(F) : s \text{ is an infinite and minimal } P\text{-position in } D(F)\} \end{aligned}$$

**Proposition 4.5.** *If  $A$  is a game and  $F$  is a set of strategies for  $A$  that is directed under the extensional order then  $\sqcup F$  is a strategy for  $A$ .*

*Proof.* Certainly  $T_{\sqcup F}$  is non-empty. To show that  $T_{\sqcup F}$  is prefix-closed, suppose  $s \in T_{\sqcup F}$ . If  $s \in T_\sigma$  for some  $\sigma \in F$ , then  $t \in T_{\sqcup F}$  for all prefixes  $t \sqsubseteq s$ , since  $T_\sigma$  is prefix closed. Otherwise, suppose  $s \in D(F)$  is infinite and that there is some finite prefix  $t \sqsubset s$  such that  $t \notin T_{\sqcup F}$ . We shall show that  $s$  is not minimal in  $D(F)$ .

We have  $s \in D_\sigma$  for all  $\sigma \in F$ . So for each  $\sigma \in F$ , there exists some prefix  $t_\sigma \sqsubseteq s$  such that  $t_\sigma \in T_\sigma \cap D_\sigma$ . If  $t \sqsubseteq t_\sigma$ , then  $t \in T_\sigma$ , which contradicts  $t$  not being in  $T_{\sqcup F}$ , so we must have  $t_\sigma \sqsubseteq t$ . Therefore, the  $t_\sigma$  are finite and have length bounded by the length of  $t$ . Let  $t_\tau$  be the longest of the  $t_\sigma$ . We have  $t_\sigma \sqsubseteq t_\tau$  for all  $\sigma \in F$ , and therefore  $t_\tau \in D(F)$ . So  $s$  is not minimal, proving that  $T_{\sqcup F}$  is prefix-closed.

Now we show that  $D_{\sqcup F}$  is postfix-closed. Let  $s \in D_{\sqcup F}$ . Then  $s \in D_\sigma$  for all  $\sigma$ . Suppose that  $s \sqsubset t$  for some  $t$ . Then  $t \in D_\sigma$  for all  $\sigma$ , so  $t \in D(F)$ . Moreover,  $t$  is certainly not minimal in  $D(F)$ , since it has  $s$  as a proper prefix. Therefore,  $t \in D_{\sqcup F}$ .

Suppose  $s \in T_{\sqcup F}$  is a  $P$ -position and that  $sa \in P_A$ . Then  $s$  is certainly finite, and so  $s \in T_\sigma$  for some  $\sigma$ . It follows that  $sa \in T_\sigma$ , so  $sa \in T_{\sqcup F}$ .

Now suppose that  $d \in D_{\sqcup F}$ . If  $d$  is infinite minimal, then  $d \in T_{\sqcup F} \cap D_{\sqcup F}$ . Moreover,  $d$  must be an  $O$ -position. Otherwise,  $d$  must have some finite prefix  $d'$  such that  $d' \in D(F)$  (if  $d$  itself is finite, we can take  $d' = d$ ). Now  $d' \in D_\sigma$  for all  $\sigma \in F$ , so for each  $\sigma \in F$  there is some  $O$ -position  $d_\sigma \sqsubseteq d'$  such that  $d_\sigma \in T_\sigma \cap D_\sigma$ . The length of the  $d_\sigma$  is bounded by the length of  $d'$ , so there is some  $\tau$  such that  $d_\sigma \sqsubseteq d_\tau$  for all  $\sigma \in F$ . Then  $d_\tau \in D_\sigma$  for all  $\sigma \in F$ , so  $d_\tau \in T_{\sqcup F} \cap D_{\sqcup F}$ .

Next we show that  $\sqcup F$  satisfies diverge-or-reply. Suppose that  $s \in T_{\sqcup F}$  is an  $O$ -position. If  $s$  is an infinite and minimal  $O$ -position in  $D(F)$ , then in particular  $s$  is in  $D(F)$  and is not a  $P$ -position, so  $s \in D_{\sqcup F}$ . Otherwise,  $s \in T_\sigma$  for some  $\sigma$ . Now suppose that  $s \notin D_{\sqcup F}$ . Since  $s$  is an  $O$ -position, this means that  $s \notin D_\tau$  for some  $\tau \in F$ . Since  $F$  is directed under the extensional order, it means that there exists some  $v \in F$  such that  $\sigma, \tau \sqsubseteq v$ . In particular,  $T_\sigma \subseteq T_v$  and  $D_v \subseteq D_\tau$ . It follows that  $s \in T_v$  but  $s \notin D_v$ ; therefore, there exists some  $a$  such that  $sa \in T_v$  and so  $sa \in T_{\sqcup F}$ .

Lastly, we need to show that  $\sqcup F$  is locally compact. Let  $s \in T_{\sqcup F}$ . If  $s \in T_\sigma$  for some  $\sigma$  then there exists  $\sigma'$  such that  $\sigma' \sqsubseteq \sigma$  and  $s \in T_{\sigma'}$ . Then certainly we have  $\sigma' \sqsubseteq \sqcup F$ . Otherwise,  $s$  is an infinite and minimal  $O$ -position in  $D(F)$ . We may define a strategy  $\sigma_s$  by:

$$\begin{aligned} T_{\sigma_s} &= \{t \in P_A : \text{for all } P\text{-positions } u \sqsubseteq t, u \sqsubseteq s\} \\ D_{\sigma_s} &= P_A \setminus \{\epsilon\} \end{aligned}$$

This is the strategy where player  $P$  always tries to play according to the sequence  $s$  and diverges if player  $O$  ever deviates from that sequence. It is clear that  $\sigma_s$  is a valid strategy. We just need to show that  $\sigma_s \sqsubseteq \bigsqcup F$ . Suppose  $t \in P_A$  is a  $P$ -position such that  $t \sqsubseteq s$  and suppose for a contradiction that  $t \notin T_{\bigsqcup F}$ . Since  $s$  is an  $O$ -position,  $t$  must be finite. For each  $\sigma \in F$ , we know that there is some  $O$ -position  $t_\sigma \sqsubseteq s$  such that  $t_\sigma \in T_\sigma \cap D_\sigma$ . Then for each  $\sigma$ , we cannot have  $t \sqsubseteq t_\sigma$ ; otherwise,  $t$  would be contained in  $T_{\bigsqcup F}$ . Therefore,  $u_\sigma \sqsubseteq t$  for all  $\sigma$  and so the  $u_\sigma$  are finite and have bounded length. It follows that there is some  $u_\tau$  such that  $u_\tau \in T_\sigma$  for all  $\sigma \in F$ , contradicting minimality of  $s$ . It follows that  $t \in T_{\bigsqcup F}$ . We then see that  $T_{\sigma_f} \subseteq T_{\bigsqcup F}$ , by closure under  $O$ -replies.

Lastly, we certainly have  $D_{\bigsqcup F} \subseteq D_{\sigma_s}$ , since  $D_{\bigsqcup F}$  may only contain non-empty positions.  $\square$

**Proposition 4.6.** *Let  $A$  be a game and let  $F$  be a set of strategies that is directed under the extensional order  $\sqsubseteq$ . Then  $\bigsqcup F$  is an extensional least upper bound for  $F$ .*

*Proof.* We certainly have  $T_\sigma \subseteq T_{\bigsqcup F}$  and  $D_{\bigsqcup F} \subseteq D_\sigma$  for all  $\sigma \in F$ . Now suppose that  $\alpha$  is some other strategy for  $A$  such that  $T_\sigma \subseteq T_\alpha$  and  $D_\alpha \subseteq D_\sigma$  for all  $\sigma \in F$ . Then we have

$$\bigcup_{\sigma \in F} T_\sigma \subseteq T_\alpha$$

$$D_\alpha \subseteq \bigcap_{\sigma \in F} D_\sigma$$

Now suppose that  $s$  is an infinite and minimal  $O$ -position in  $D(F)$ . We show that  $s \in T_\alpha$ . Indeed, if  $s \in D(F)$  then  $s \in D_\sigma$  for all  $\sigma \in F$ . Then for each  $\sigma \in F$  there exists some  $s_\sigma \sqsubseteq s$  such that  $s_\sigma \in T_\sigma \cap D_\sigma$ . Since  $s$  is minimal in  $D(F)$ , either one of the  $s_\sigma$  is equal to  $s$  or the  $s_\sigma$  are finite but unbounded in length.  $\square$

## 4.2 Soundness

## 4.3 Adequacy

I don't think I'll have a full abstraction result yet, but I could add a section explaining how this model gives us some progress towards full abstraction.

## References

- [AJ13] Samson Abramsky and Radha Jagadeesan. Games and full completeness for multiplicative linear logic. *CoRR*, abs/1311.6057, 2013.

- [Har99] Russell S. Harmer. Games and full abstraction for nondeterministic languages. Technical report, 1999.
- [HM99] R. Harmer and G. McCusker. A fully abstract game semantics for finite nondeterminism. In *Proceedings. 14th Symposium on Logic in Computer Science (Cat. No. PR00158)*, pages 422–430, 1999.
- [Hyl97] Martin Hyland. Game semantics. *Semantics and logics of computation*, 14:131, 1997.
- [Lai15] J. Laird. Sequential algorithms for unbounded nondeterminism. *Electronic Notes in Theoretical Computer Science*, 319:271 – 287, 2015.
- [Lev08] Paul Blain Levy. Infinite trace equivalence. *Annals of Pure and Applied Logic*, 151(2):170 – 198, 2008.
- [ROS93] A. W. ROSCOE. Unbounded non-determinism in CSP. *Journal of Logic and Computation*, 3(2):131, 1993.
- [TO15] Takeshi Tsukada and C. H. Luke Ong. Nondeterminism in game semantics via sheaves. In *Proceedings of the 2015 30th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, LICS '15, pages 220–231, Washington, DC, USA, 2015. IEEE Computer Society.