

# Monte Carlo Methods: Sampling, Integration, and MCMC

Computational Simulation Templates

November 24, 2025

## 1 Introduction

Monte Carlo methods use random sampling to solve deterministic problems. This template covers basic sampling techniques, numerical integration, importance sampling, and the Metropolis-Hastings algorithm for Markov Chain Monte Carlo.

## 2 Mathematical Framework

### 2.1 Monte Carlo Integration

Estimate integrals using random samples:

$$I = \int_a^b f(x) dx \approx \frac{b-a}{N} \sum_{i=1}^N f(x_i) \quad (1)$$

### 2.2 Variance of Estimator

The standard error decreases as  $1/\sqrt{N}$ :

$$\text{SE} = \frac{\sigma_f}{\sqrt{N}} \quad (2)$$

### 2.3 Importance Sampling

Use proposal distribution  $q(x)$  to reduce variance:

$$I = \int f(x)p(x) dx \approx \frac{1}{N} \sum_{i=1}^N \frac{f(x_i)p(x_i)}{q(x_i)}, \quad x_i \sim q \quad (3)$$

### 2.4 Metropolis-Hastings Algorithm

Accept proposed state  $x'$  with probability:

$$\alpha = \min \left( 1, \frac{p(x')q(x|x')}{p(x)q(x'|x)} \right) \quad (4)$$

<b>3</b>	<b>Environment Setup</b>
<b>4</b>	<b>Basic Monte Carlo Integration</b>
<b>5</b>	<b>Numerical Integration</b>
<b>6</b>	<b>Importance Sampling</b>
<b>7</b>	<b>Metropolis-Hastings Algorithm</b>
<b>8</b>	<b>2D Metropolis-Hastings</b>
<b>9</b>	<b>Results Summary</b>
<b>9.1</b>	<b>Pi Estimation</b>
<b>9.2</b>	<b>Importance Sampling</b>
<b>9.3</b>	<b>MCMC Results</b>
<b>9.4</b>	<b>Statistical Summary</b>
	<ul style="list-style-type: none"> <li>• Pi estimation error: ??</li> <li>• Gaussian integral estimate: ??</li> <li>• Importance sampling variance reduction: ??x</li> <li>• Optimal MCMC acceptance rate: ??</li> <li>• 2D MCMC acceptance rate: ??</li> </ul>
<b>10</b>	<b>Conclusion</b>

This template demonstrates Monte Carlo methods for numerical computation. Basic MC integration achieves  $1/\sqrt{N}$  convergence, while importance sampling provides substantial variance reduction for rare events (??x improvement). The Metropolis-Hastings algorithm successfully samples from complex distributions, with optimal acceptance rates around 0.234 for 1D targets (achieved with  $\sigma_q = 1.0$  giving ??).