

Quantum Theory of Solids

Lecture notes - Spring 2020

March 30, 2020

Digitalized lecture notes for the course “TFY4210 - Quantum Theory of Many-Particle Systems” held by Prof. Asle Sudbø spring 2020. These notes follow the pdf containing the hand written lecture notes, which are based upon the lecture notes for the course “FY8302 - Quantum Theory of Solids”

Website: <https://www.ntnu.edu/studies/courses/TFY4210>

Contributors

- Karl Kristian Ladegård Lockert

Contents

1	Introduction	2
2	Many-particle states, fermions	3
2.1	N-particle vacuum state	3
2.2	Completeness relation	3
2.3	Operators	3
3	Quasi-particles in interacting electron-systems. Fermi-liquids.	4
3.1	Fermi-liquids	4
3.2	Screening of the Coulomb-interaction	4
3.3	Phonon mediated electron-electron interaction	4
3.4	Magnon mediated electron-electron interaction	5
4	The Cooper-problem	5

1 Introduction

2 Many-particle states, fermions

2.1 N-particle vacuum state

2.2 Completeness relation

2.3 Operators

3 Quasi-particles in interacting electron-systems. Fermi-liquids.

3.1 Fermi-liquids

3.2 Screening of the Coulomb-interaction

3.3 Phonon mediated electron-electron interaction

Due to the electron-phonon coupling depicted in **TODO: SETT INN DIAGRAM**, we will get an effective phonon-mediated interaction between electrons.

This is an exchange of a virtual phonon. The above diagram is the effective interaction to second order in g_q if we regard the wavy line $\sim\sim\sim$ as a bare phonon Green's function. We could also imagine that we replaced this line by **TODO: SETT INN DIAGRAM** which would include an effective interaction computed correctly up to order $\mathcal{O}(g^4)$. In fact, we might replace $D_0 + D_0 \Pi D_0 + \dots$ by D ! Thus computing the effective interaction up to infinite order in g . Another, often used approach would be to replace $\sim\sim\sim$ by **TODO: SETT INN DIAGRAM**

Here, we have resummed a subset of diagrams to infinite order in g in order to get an effective interaction between electrons. Under the assumption that g is weak, we will keep terms only to $\mathcal{O}(g^2)$.

$$V_{\text{eff}}(q, \omega) = |g_q|^2 \frac{2\omega_q}{\omega^2 - \omega_q^2} \quad (3.1)$$

Thus, the interaction part of the Hamiltonian becomes

$$\mathcal{H} = \sum_{\substack{k, k', q \\ \sigma, \sigma'}} V_{\text{tot}}(q, \omega) c_{k+q, \sigma}^\dagger c_{k', \sigma'}^\dagger c_{k', \sigma'} c_{k, \sigma} \quad (3.2)$$

$$V_{\text{tot}}(q, \omega) = \frac{e^2}{4\pi\epsilon q^2} + V_{\text{eff}}(q, \omega), \quad (3.3)$$

where the first term is the Coulomb-interaction. Furthermore, ω is the energy transfer between scattering electrons when they exchange a phonon

$$\omega = \varepsilon_{k+q} - \varepsilon_k \quad (3.4)$$

TODO: Sett inn figur

Note the singularities in V_{tot} when $|\omega| \rightarrow \omega_q$. In particular, note the negative singularity when $|\omega| \rightarrow \omega_q^-$. This singularity persists when Coulomb-repulsion is included. For most frequencies, the Coulomb-interaction completely dominates. However, in a narrow ω -region close to ω_q , the extremely weak electron-phonon coupling will always beat the Coulomb-interaction! This frequency is slightly smaller than ω_q . For small ω , V_{tot} is repulsive. For large ω , V_{tot} is repulsive. For $|\omega| \lesssim \omega_q$, V_{tot} is attractive.

Let us try to give a physical picture for this: When an electron moves past an ion, they interact. The electron pulls slightly on the heavy, positively charged ion. Electrons are light, and move much faster than the heavy ions. The electron moves quickly out of the scattering zone, while the ion relaxes slowly back to its equilibrium position. The ion in its out-of-equilibrium position represents excessive positive charge in that position, which can pull another electron towards it. This is effectively a charge-dipole interaction. If the second electron “waits” a little for the first electron to get away (thus reducing Coulomb-repulsion) but does not wait for too long (such that the ion has relaxed back to its equilibrium position), then the second electron can be attracted to the scattering region. Effectively, the second electron is attracted to the scattering region because the first electron was there. This is an effective electron-electron attraction. It only works if the electron waits a little, but not for too long. A minimum time corresponds to a maximum frequency, while a maximum time corresponds to a minimum frequency. This implies that V_{tot} is attractive if $\omega_{\text{min}} < \omega < \omega_{\text{max}}$, as depicted in **TODO: Sett in figurer og referanse til figuren (s.7)** We may view the effective electron-electron attraction as a result of an electron locally deforming an elastic medium. Think of a rubber membrane that you put a little metal sphere on. The membrane is stretched, dipping down where you put the first sphere. If you put another little sphere on the membrane, it will fall into the dip, i.e. it will be attracted to the first particle. This is also how gravity works: A mass deforms space-time (an elastic medium) and thus attracts another mass.

Disclaimer: The above two analogs are classical. There will be an important quantum effect coming into play here, which we will come back to. Here, it will suffice to note that, classically, one can keep adding particles to the dip, such that all particles will be gathered in the same one, forming a large heavy object. This is not how it works quantum mechanically with fermions. Note also that in V_{tot} , and the two different simplified models for \bar{V} , they are only attractive up to a maximum ω , i.e. only after a minimum amount of time. The second particle has to wait a minimum amount of time for the interaction to be attractive. This is called retardation.

The electrons avoid the Coulomb-interactions by avoiding each other, not in space, but in time.

3.4 Magnon mediated electron-electron interaction

4 The Cooper-problem